Jelena B JoviÄ

List of Publications by Year
in descending order

Source: https://exaly.com/author-pdf/6043433/publications.pdf

Version: 2024-02-01

65 papers	1,072 citations	430874 18 h-index	477307 29 g-index
65 all docs	65 docs citations	65 times ranked	839 citing authors

#	Article	IF	Citations
1	Genetic Diversity of Flavescence DorÃ©e Phytoplasmas in Vineyards of Serbia: From the Widespread Occurrence of Autochthonous Map-M51 to the Emergence of Endemic Map-FD2 (Vectotype II) and New Map-FD3 (Vectotype III) Epidemic Genotypes. Agronomy, 2022, 12, 448.	3.0	6
2	Symptomatology, (Co)occurrence and Differential Diagnostic PCR Identification of â€˜Ca. Phytoplasma solaniâ€™ and â€˜Ca. Phytoplasma convolvuliâ€™ in Field Bindweed. Pathogens, 2021, 10, 160.	2.8	4
3	The â€œcode redâ€•for Balkan vineyards: occurrence of Orientus ishidae (Matsumura, 1902) (Hemiptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 147 Td	1.1	2
4	Diversity of phytoplasmas identified in the polyphagous leafhopper Euscelis incisus (Cicadellidae,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 Td European Journal of Plant Pathology, 2020, 156, 201-221.	1.7	14
5	When a Palearctic bacterium meets a Nearctic insect vector: Genetic and ecological insights into the emergence of the grapevine Flavescence dorÃ©e epidemics in Europe. PLoS Pathogens, 2020, 16, e1007967.	4.7	55
6	Role of plant-specialized Hyalesthes obsoletus associated with Convolvulus arvensis and Crepis foetida in the transmission of â€˜Candidatus Phytoplasma solaniâ€™-inflicted bois noir disease of grapevine in Serbia. European Journal of Plant Pathology, 2019, 153, 183-195.	1.7	31
7	Vector Role of Cixiids and Other Planthopper Species. , 2019, , 79-113.		6
8	Resource allocation in response to herbivory and gall formation in Linaria vulgaris. Plant Physiology and Biochemistry, 2019, 135, 224-232.	5.8	10
9	Twentyâ€five years after: postâ€introduction association of Mecinus janthinus s.l. with invasive host toadflaxes Linaria vulgaris and Linaria dalmatica in North America. Annals of Applied Biology, 2018, 173, 16-34.	2.5	11
10	Wolbachia infection in natural populations of Dictyophara europaea, an alternative vector of grapevine Flavescence dorÃ©e phytoplasma: effects and interactions. Annals of Applied Biology, 2018, 172, 47-64.	2.5	20
11	Occurrence and Epidemiological Aspects of Phytoplasmas in Cereals. , 2018, , 67-89.		5
12	Widespread plant specialization in the polyphagous planthopper Hyalesthes obsoletus (Cixiidae), a major vector of stolbur phytoplasma: Evidence of cryptic speciation. PLoS ONE, 2018, 13, e0196969.	2.5	20
13	Molecular and experimental evidence of multi-resistance of Cercospora beticola field populations to MBC, DMI and QoI fungicides. European Journal of Plant Pathology, 2017, 149, 895-910.	1.7	19
14	Divergent evolution of life span associated with mitochondrial DNA evolution. Evolution; International Journal of Organic Evolution, 2017, 71, 160-166.	2.3	12
15	Dictyophara europaea (Hemiptera: Fulgoromorpha: Dictyopharidae): description of immatures, biology and host plant associations. Bulletin of Entomological Research, 2016, 106, 395-405.	1.0	15
16	Comparative analysis of phenolic profiles of ovipositional fluid of Rhinusa pilosa (Mecinini,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 Td 2016, 10, 311-322.	1.1	4
17	Molecular tracing of the transmission routes of bois noir in Mediterranean vineyards of Montenegro and experimental evidence for the epidemiological role of Vitex agnusâ€castus (Lamiaceae) and associated Hyalesthes obsoletus (Cixiidae). Plant Pathology, 2016, 65, 285-298.	2.4	65
18	Morphology versus DNA barcoding: two sides of the same coin. A case study of Ceutorhynchus erysimi and C. contractus identification. Insect Science, 2016, 23, 638-648.	3.0	10

#	Article	IF	Citations
19	â€˜Candidatus phytoplasma solaniâ€™ genotypes associated with potato stolbur in Serbia and the role of Hyalesthes obsoletus and Reptalus panzeri (hemiptera, cixiidae) as natural vectors. European Journal of Plant Pathology, 2016, 144, 619-630.	1.7	32
20	First Report of QoI Resistance in Botrytis cinerea Isolates Causing Gray Mold in Strawberry Fields in Serbia. Plant Disease, 2016, 100, 221-221.	1.4	1
21	First Report of â€˜Candidatus Phytoplasma solaniâ€™ Associated With Potato Stolbur Disease in Montenegro. Plant Disease, 2016, 100, 1775-1775.	1.4	3
22	Bioremediation Potential Assessment of Plant Growth-Promoting Autochthonous Bacteria: a Lignite Mine Case Study. Polish Journal of Environmental Studies, 2016, 25, 113-119.	1.2	3
23	Euscelis incisus(Cicadellidae, Deltocephalinae), a natural vector of 16SrIII-B phytoplasma causing multiple inflorescence disease ofCirsium arvense. Annals of Applied Biology, 2015, 167, 406-419.	2.5	8
24	Hostâ€associated genetic divergence and taxonomy in the Rhinusa pilosa<scp>G</scp>yllenhal species complex: an integrative approach. Systematic Entomology, 2015, 40, 268-287.	3.9	13
25	Occurrence of Cercospora beticola populations resistant to benzimidazoles and demethylation-inhibiting fungicides in SerbiaÂandÂtheir impact on disease management. Crop Protection, 2015, 75, 80-87.	2.1	18
26	The molecular epidemiology of bois noir grapevine yellows caused by â€˜Candidatus Phytoplasma solaniâ€™ in the Republic of Macedonia. European Journal of Plant Pathology, 2015, 142, 759-770.	1.7	34
27	First Report of Cercospora armoraciae, Causal Agent of Cercospora Leaf Spot, on Horseradish in Serbia. Plant Disease, 2015, 99, 1645-1645.	1.4	1
28	First Report of â€˜Candidatus Phytoplasma solaniâ€™ Infecting Garden Bean Phaseolus vulgaris in Serbia. Plant Disease, 2015, 99, 551-551.	1.4	6
29	First Report of Thielaviopsis thielavioides, A Causal Agent of Postharvest Blackening on Daucus carota in Serbia. Plant Disease, 2015, 99, 1274.	1.4	4
30	Potential Hemipteran vectors of â€œstolburâ€•phytoplasma in potato fields in Serbia. Phytopathogenic Mollicutes, 2015, 5, S49.	0.1	4
31	First Report of Cercospora violae Infecting the Garden Violet Viola odorata in Serbia. Plant Disease, 2015, 99, 1035.	1.4	0
32	First Report of Cercospora apii, Causal Agent of Cercospora Early Blight of Celery, in Serbia. Plant Disease, 2014, 98, 1157-1157.	1.4	6
33	First Report of Cercospora carotae, Causal Agent of Cercospora Leaf Spot of Carrot, in Serbia. Plant Disease, 2014, 98, 1153-1153.	1.4	3
34	Experimental and molecular evidence of <scp>R</scp>eptalus panzeri as a natural vector of bois noir. Plant Pathology, 2014, 63, 42-53.	2.4	92
35	Revision of <scp>M</scp>ecinus heydenii species complex (<scp>C</scp>urculionidae): integrative taxonomy reveals multiple species exhibiting host specialization. Zoologica Scripta, 2014, 43, 34-51.	1.7	11
36	First Report of Alder Yellows Phytoplasma Associated with Common Alder (Alnus glutinosa) in the Republic of Macedonia. Plant Disease, 2014, 98, 1268-1268.	1.4	11

#	Article	IF	Citations
37	First Report of Stolbur Phytoplasma Associated with Maize Redness Disease of Maize in Bosnia and Herzegovina. Plant Disease, 2014, 98, 418-418.	1.4	9
38	Crepis foetida L.: New host plant of cixiid planthopper Hyalesthes obsoletus Signoret 1865 (Hemiptera:) Tj ETQq0 0 0 rgBT /Overlock 10	0.2	4
39	Drosophila suzukii (Matsumura, 1931) (Siptera: Srosophilidae): A new invasive pest in Serbia. Zastita Bilja, 2014, 65, 99-104.	0.2	18
40	Morphological and molecular identification of Cercospora apii on celery in Serbia. Zastita Bilja, 2014, 65, 77-84.	0.2	0
41	PCR-RFLP-based method for reliable discrimination of cryptic species within Mecinus janthinus species complex (Mecinini, Curculionidae) introduced in North America for biological control of invasive toadflaxes. BioControl, 2013, 58, 563-573.	2.0	15
42	Characterisation of benzimidazole resistance of Cercospora beticola in Serbia using PCR-based detection of resistance-associated mutations of the Î²-tubulin gene. European Journal of Plant Pathology, 2013, 135, 889-902.	1.7	24
43	First Report of Alder Yellows Phytoplasma Infecting Common and Grey Alder (Alnus glutinosa) Tj ETQq1 1 0.784314 rgBT /Ove	1.4	11
44	Characterisation of a 16SrII phytoplasma strain associated with bushy stunt of hawkweed oxtongue (Picris hieracioides) in south-eastern Serbia and the role of the leafhopper Neoaliturus fenestratus (Deltocephalinae) as a natural vector. European Journal of Plant Pathology, 2012, 134, 647-660.	1.7	25
45	Non-persistently aphid-borne viruses infecting pumpkin and squash in Serbia and partial characterization of Zucchini yellow mosaic virus isolates. European Journal of Plant Pathology, 2012, 133, 935-947.	1.7	26
46	Biodegradation of methyl tert-butyl ether by Kocuria sp.. Hemijska Industrija, 2012, 66, 717-722.	0.7	5
47	Multigene sequence data and genetic diversity among â€˜Candidatus Phytoplasma ulmiâ€™ strains infecting Ulmus spp. in Serbia. Plant Pathology, 2011, 60, 356-368.	2.4	22
48	Morphological, molecular and biological evidence reveal two cryptic species inMecinus janthinusGermar (Coleoptera, Curculionidae), a successful biological control agent of Dalmatian toadflax,Linaria dalmatica(Lamiales, Plantaginaceae). Systematic Entomology, 2011, 36, 741-753.	3.9	46
49	First Report of the Occurrence of Cucurbit aphid-borne yellows virus on Oilseed Pumpkin in Serbia. Plant Disease, 2011, 95, 1035-1035.	1.4	5
50	First Report of Tomato spotted wilt virus on Gerbera hybrida in Serbia. Plant Disease, 2011, 95, 226-226.	1.4	13
51	First Report of Plasmopara obducens on Impatiens walleriana in Serbia. Plant Disease, 2011, 95, 491-491.	1.4	9
52	Tuta absoluta (Meyrick, 1917) (Lepidoptera, Gelechiidae): A new pest of tomato in Serbia. Pesticidi I Fitomedicina = Pesticides and Phytomedicine, 2011, 26, 197-204.	0.2	7
53	Plasmopara obducens: A new threat to the production of Impatiens Walleriana in Serbia. Pesticidi I Fitomedicina = Pesticides and Phytomedicine, 2011, 26, 43-53.	0.2	2
54	Phytophthora ramorum Occurrence in Ornamentals in Serbia. Plant Disease, 2010, 94, 703-708.	1.4	8

#	Article	IF	Citations
55	Host-associated genetic differentiation in a seed parasitic weevil Rhinusa antirrhini (Coleptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 2286-2300.	3.9	35
56	First report of Phytophthora ramorum on Rhododendron sp. in Serbia. Plant Pathology, 2009, 58, 804-804.	2.4	3
57	Molecular characteristics of phytoplasmas associated with Flavescence dorÃ©e in clematis and grapevine and preliminary results on the role of Dictyophara europaea as a vector. Plant Pathology, 2009, 58, 826-837.	2.4	74
58	Occurrence and Distribution of Grapevine Yellows Caused by Stolbur Phytoplasma in Montenegro. Journal of Phytopathology, 2009, 157, 682-685.	1.0	14
59	Stolbur Phytoplasma Transmission to Maize by Reptalus panzeri and the Disease Cycle of Maize Redness in Serbia. Phytopathology, 2009, 99, 1053-1061.	2.2	44
60	Incidence and Distribution of Iris yellow spot virus on Onion in Serbia. Plant Disease, 2009, 93, 976-982.	1.4	21
61	First report of alder yellows phytoplasma on common alder (Alnus glutinosa) in Serbia. Plant Pathology, 2008, 57, 773-773.	2.4	8
62	New strain of â€˜Candidatus Phytoplasma ulmiâ€™ infecting Ulmus minor and U. laevis in Serbia. Plant Pathology, 2008, 57, 1174-1174.	2.4	9
63	First Report of Iris yellow spot virus on Onion (Allium cepa) in Serbia. Plant Disease, 2008, 92, 1247-1247.	1.4	10
64	Roles of stolbur phytoplasma and Reptalus panzeri (Cixiinae, Auchenorrhyncha) in the epidemiology of Maize redness in Serbia. European Journal of Plant Pathology, 2007, 118, 85-89.	1.7	43
65	Framework for risk assessment of â€˜Candidatus Phytoplasma solaniâ€™ associated diseases outbreaks in agroecosystems in Serbia. Journal of Plant Pathology, 0, , 1.	1.2	3