

# Jing-Quan Yu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6044902/publications.pdf>

Version: 2024-02-01

185  
papers

16,000  
citations

12303

69  
h-index

19136

118  
g-index

185  
all docs

185  
docs citations

185  
times ranked

13762  
citing authors

#	ARTICLE	IF	CITATIONS
1	Brassinosteroids' regulation of plant architecture. , 2022, , 43-57.		1
2	Lightâ€dependent activation of HY5 promotes mycorrhizal symbiosis in tomato by systemically regulating strigolactone biosynthesis. <i>New Phytologist</i> , 2022, 233, 1900-1914.	3.5	30
3	Melatonin delays darkâ€induced leaf senescence by inducing <i>miR171b</i> expression in tomato. <i>Journal of Pineal Research</i> , 2022, 72, .	3.4	22
4	S-Nitrosoglutathione Reductase Contributes to Thermotolerance by Modulating High Temperature-Induced Apoplastic H <sub>2</sub> O <sub>2</sub> in <i>Solanum lycopersicum</i> . <i>Frontiers in Plant Science</i> , 2022, 13, 862649.	1.7	0
5	Exogenous Rosmarinic Acid Application Enhances Thermotolerance in Tomatoes. <i>Plants</i> , 2022, 11, 1172.	1.6	6
6	The <i>miR164a</i> - <i>NAM3</i> module confers cold tolerance by inducing ethylene production in tomato. <i>Plant Journal</i> , 2022, 111, 440-456.	2.8	35
7	High Nitric Oxide Concentration Inhibits Photosynthetic Pigment Biosynthesis by Promoting the Degradation of Transcription Factor HY5 in Tomato. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6027.	1.8	6
8	Glucose sensing by regulator of G protein signaling 1 ( <i>RGS1</i> ) plays a crucial role in coordinating defense in response to environmental variation in tomato. <i>New Phytologist</i> , 2022, 236, 561-575.	3.5	8
9	Brassinosteroids promote starch synthesis and the implication in low-light stress tolerance in <i>Solanum lycopersicum</i> . <i>Environmental and Experimental Botany</i> , 2022, 201, 104990.	2.0	5
10	High CO <sub>2</sub> and pathogen-driven expression of the carbonic anhydrase <i>CA3</i> confers basal immunity in tomato. <i>New Phytologist</i> , 2021, 229, 2827-2843.	3.5	26
11	Ethylene response factors 15 and 16 trigger jasmonate biosynthesis in tomato during herbivore resistance. <i>Plant Physiology</i> , 2021, 185, 1182-1197.	2.3	32
12	Nitrogen forms and metabolism affect plant defence to foliar and root pathogens in tomato. <i>Plant, Cell and Environment</i> , 2021, 44, 1596-1610.	2.8	37
13	Crosstalk between Brassinosteroid and Redox Signaling Contributes to the Activation of CBF Expression during Cold Responses in Tomato. <i>Antioxidants</i> , 2021, 10, 509.	2.2	16
14	The protein kinase CPK28 phosphorylates ascorbate peroxidase and enhances thermotolerance in tomato. <i>Plant Physiology</i> , 2021, 186, 1302-1317.	2.3	61
15	Brassinosteroid signaling integrates multiple pathways to release apical dominance in tomato. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	49
16	The phyâ€dependent induction of HY5 promotes iron uptake by systemically activating <i>FER</i> expression. <i>EMBO Reports</i> , 2021, 22, e51944.	2.0	37
17	ELONGATED HYPOCOTYL 5 mediates blue light-induced starch degradation in tomato. <i>Journal of Experimental Botany</i> , 2021, 72, 2627-2641.	2.4	21
18	Strigolactones positively regulate abscisic acid-dependent heat and cold tolerance in tomato. <i>Horticulture Research</i> , 2021, 8, 237.	2.9	47

#	ARTICLE	IF	CITATIONS
19	Rosmarinic Acid Delays Tomato Fruit Ripening by Regulating Ripening-Associated Traits. <i>Antioxidants</i> , 2021, 10, 1821.	2.2	14
20	BRASSINAZOLE RESISTANT 1 Mediates Brassinosteroid-Induced Calvin Cycle to Promote Photosynthesis in Tomato. <i>Frontiers in Plant Science</i> , 2021, 12, 811948.	1.7	10
21	Crosstalk of PIF4 and DELLA modulates CBF transcript and hormone homeostasis in cold response in tomato. <i>Plant Biotechnology Journal</i> , 2020, 18, 1041-1055.	4.1	65
22	Brassinosteroid-mediated reactive oxygen species are essential for tapetum degradation and pollen fertility in tomato. <i>Plant Journal</i> , 2020, 102, 931-947.	2.8	55
23	Melatonin promotes metabolism of bisphenol A by enhancing glutathione-dependent detoxification in <i>Solanum lycopersicum</i> L. <i>Journal of Hazardous Materials</i> , 2020, 388, 121727.	6.5	31
24	Light-induced HY5 Functions as a Systemic Signal to Coordinate the Photoprotective Response to Light Fluctuation. <i>Plant Physiology</i> , 2020, 184, 1181-1193.	2.3	20
25	The Chromosome-Scale Genome of Melon Dissects Genetic Architecture of Important Agronomic Traits. <i>IScience</i> , 2020, 23, 101422.	1.9	22
26	The <i>HY5</i> and <i>MYB15</i> transcription factors positively regulate cold tolerance in tomato via the <i>CBF</i> pathway. <i>Plant, Cell and Environment</i> , 2020, 43, 2712-2726.	2.8	56
27	Brassinosteroids act as a positive regulator of NBR1-dependent selective autophagy in response to chilling stress in tomato. <i>Journal of Experimental Botany</i> , 2020, 71, 1092-1106.	2.4	56
28	Transcriptomic and genetic approaches reveal an essential role of the NAC transcription factor SINAP1 in the growth and defense response of tomato. <i>Horticulture Research</i> , 2020, 7, 209.	2.9	37
29	Histone acetylation recruits the SWR1 complex to regulate active DNA demethylation in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 16641-16650.	3.3	73
30	Tomato <i>GLR3.3</i> and <i>GLR3.5</i> mediate cold acclimation-induced chilling tolerance by regulating apoplastic H <sub>2</sub> O <sub>2</sub> production and redox homeostasis. <i>Plant, Cell and Environment</i> , 2019, 42, 3326-3339.	2.8	56
31	COP9 Signalosome CSN4 and CSN5 Subunits Are Involved in Jasmonate-Dependent Defense Against Root-Knot Nematode in Tomato. <i>Frontiers in Plant Science</i> , 2019, 10, 1223.	1.7	16
32	Systemic Root-Shoot Signaling Drives Jasmonate-Based Root Defense against Nematodes. <i>Current Biology</i> , 2019, 29, 3430-3438.e4.	1.8	89
33	Glutaredoxin S25 and its interacting TGACC motif-binding factor TGA2 mediate brassinosteroid-induced chlorothaloniol metabolism in tomato plants. <i>Environmental Pollution</i> , 2019, 255, 113256.	3.7	28
34	Role of ethylene biosynthesis and signaling in elevated CO <sub>2</sub> -induced heat stress response in tomato. <i>Planta</i> , 2019, 250, 563-572.	1.6	57
35	Brassinosteroids Act as a Positive Regulator of Photoprotection in Response to Chilling Stress. <i>Plant Physiology</i> , 2019, 180, 2061-2076.	2.3	90
36	A novel CO <sub>2</sub> -responsive systemic signaling pathway controlling plant mycorrhizal symbiosis. <i>New Phytologist</i> , 2019, 224, 106-116.	3.5	28

#	ARTICLE	IF	CITATIONS
37	Strigolactones positively regulate defense against root-knot nematodes in tomato. <i>Journal of Experimental Botany</i> , 2019, 70, 1325-1337.	2.4	59
38	SlHY5 Integrates Temperature, Light, and Hormone Signaling to Balance Plant Growth and Cold Tolerance. <i>Plant Physiology</i> , 2019, 179, 749-760.	2.3	71
39	BZR1 Mediates Brassinosteroid-Induced Autophagy and Nitrogen Starvation in Tomato. <i>Plant Physiology</i> , 2019, 179, 671-685.	2.3	114
40	A Plant Phytosulfokine Peptide Initiates Auxin-Dependent Immunity through Cytosolic Ca <sup>2+</sup> Signaling in Tomato. <i>Plant Cell</i> , 2018, 30, 652-667.	3.1	120
41	The <i>Zip</i> transcription factor <i>HY5</i> mediates <i>CRY1a</i> -induced anthocyanin biosynthesis in tomato. <i>Plant, Cell and Environment</i> , 2018, 41, 1762-1775.	2.8	138
42	Heat Shock Factor HsfA1a Is Essential for <i>R</i> Gene-Mediated Nematode Resistance and Triggers H <sub>2</sub> O <sub>2</sub> Production <sup>1</sup> . <i>Plant Physiology</i> , 2018, 176, 2456-2471.	2.3	52
43	Induction of systemic resistance in tomato against <i>Botrytis cinerea</i> by N-decanoyl-homoserine lactone via jasmonic acid signaling. <i>Planta</i> , 2018, 247, 1217-1227.	1.6	37
44	Tomato photorespiratory glycolate oxidase-derived H <sub>2</sub> O <sub>2</sub> production contributes to basal defence against <i>Pseudomonas syringae</i> . <i>Plant, Cell and Environment</i> , 2018, 41, 1126-1138.	2.8	28
45	Brassinosteroids act as a positive regulator for resistance against root-knot nematode involving RESPIRATORY BURST OXIDASE HOMOLOG-dependent activation of MAPKs in tomato. <i>Plant, Cell and Environment</i> , 2018, 41, 1113-1125.	2.8	59
46	Tomato <i>CRY1a</i> plays a critical role in the regulation of phytohormone homeostasis, plant development, and carotenoid metabolism in fruits. <i>Plant, Cell and Environment</i> , 2018, 41, 354-366.	2.8	44
47	Light Signaling-Dependent Regulation of Photoinhibition and Photoprotection in Tomato. <i>Plant Physiology</i> , 2018, 176, 1311-1326.	2.3	85
48	Brassinosteroid-mediated apoplastic <i>H<sub>2</sub>O<sub>2</sub></i> glutaredoxin 12/14 cascade regulates antioxidant capacity in response to chilling in tomato. <i>Plant, Cell and Environment</i> , 2018, 41, 1052-1064.	2.8	95
49	Phytomelatonin: Recent advances and future prospects. <i>Journal of Pineal Research</i> , 2018, 65, e12526.	3.4	148
50	A Method of High Throughput Monitoring Crop Physiology Using Chlorophyll Fluorescence and Multispectral Imaging. <i>Frontiers in Plant Science</i> , 2018, 9, 407.	1.7	44
51	Glutaredoxin GRXS16 mediates brassinosteroid-induced apoplastic H <sub>2</sub> O <sub>2</sub> production to promote pesticide metabolism in tomato. <i>Environmental Pollution</i> , 2018, 240, 227-234.	3.7	37
52	BZR1 Transcription Factor Regulates Heat Stress Tolerance Through FERONIA Receptor-Like Kinase-Mediated Reactive Oxygen Species Signaling in Tomato. <i>Plant and Cell Physiology</i> , 2018, 59, 2239-2254.	1.5	91
53	The role of calcium-dependent protein kinase in hydrogen peroxide, nitric oxide and ABA-dependent cold acclimation. <i>Journal of Experimental Botany</i> , 2018, 69, 4127-4139.	2.4	73
54	HsfA1a upregulates melatonin biosynthesis to confer cadmium tolerance in tomato plants. <i>Journal of Pineal Research</i> , 2017, 62, e12387.	3.4	219

#	ARTICLE	IF	CITATIONS
55	Nitric oxide is involved in the oxytetracycline-induced suppression of root growth through inhibiting hydrogen peroxide accumulation in the root meristem. <i>Scientific Reports</i> , 2017, 7, 43096.	1.6	16
56	Crosstalk between Nitric Oxide and MPK1/2 Mediates Cold Acclimation-induced Chilling Tolerance in Tomato. <i>Plant and Cell Physiology</i> , 2017, 58, 1963-1975.	1.5	47
57	24-Epibrassinolide alleviates organic pollutants-retarded root elongation by promoting redox homeostasis and secondary metabolism in <i>Cucumis sativus</i> L. <i>Environmental Pollution</i> , 2017, 229, 922-931.	3.7	57
58	A novel ethylene responsive factor CitERF13 plays a role in photosynthesis regulation. <i>Plant Science</i> , 2017, 256, 112-119.	1.7	14
59	Importance of the mitochondrial alternative oxidase (AOX) pathway in alleviating photoinhibition in cucumber leaves under chilling injury and subsequent recovery when leaves are subjected to high light intensity. <i>Journal of Horticultural Science and Biotechnology</i> , 2017, 92, 31-38.	0.9	14
60	A New Strategy in Observer Modeling for Greenhouse Cucumber Seedling Growth. <i>Frontiers in Plant Science</i> , 2017, 8, 1297.	1.7	1
61	Detecting crop population growth using chlorophyll fluorescence imaging. <i>Applied Optics</i> , 2017, 56, 9762.	0.9	6
62	The critical role of autophagy in plant responses to abiotic stresses. <i>Frontiers of Agricultural Science and Engineering</i> , 2017, 4, 28.	0.9	9
63	Physiological and Transcriptome Responses to Combinations of Elevated CO <sub>2</sub> and Magnesium in <i>Arabidopsis thaliana</i> . <i>PLoS ONE</i> , 2016, 11, e0149301.	1.1	19
64	Genome-Wide Identification and Expression Analysis of Calcium-dependent Protein Kinase in Tomato. <i>Frontiers in Plant Science</i> , 2016, 7, 469.	1.7	62
65	Brassinosteroid Ameliorates Zinc Oxide Nanoparticles-Induced Oxidative Stress by Improving Antioxidant Potential and Redox Homeostasis in Tomato Seedling. <i>Frontiers in Plant Science</i> , 2016, 7, 615.	1.7	84
66	Unraveling Main Limiting Sites of Photosynthesis under Below- and Above-Ground Heat Stress in Cucumber and the Alleviatory Role of Luffa Rootstock. <i>Frontiers in Plant Science</i> , 2016, 7, 746.	1.7	33
67	Melatonin mediates selenium-induced tolerance to cadmium stress in tomato plants. <i>Journal of Pineal Research</i> , 2016, 61, 291-302.	3.4	211
68	Apoplastic H <sub>2</sub> O <sub>2</sub> plays a critical role in axillary bud outgrowth by altering auxin and cytokinin homeostasis in tomato plants. <i>New Phytologist</i> , 2016, 211, 1266-1278.	3.5	49
69	Involvement of an ethylene response factor in chlorophyll degradation during citrus fruit degreening. <i>Plant Journal</i> , 2016, 86, 403-412.	2.8	130
70	Microarray and genetic analysis reveals that csa-miR159b plays a critical role in abscisic acid-mediated heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2016, 39, 1790-1804.	2.8	52
71	Grafting cucumber onto luffa improves drought tolerance by increasing ABA biosynthesis and sensitivity. <i>Scientific Reports</i> , 2016, 6, 20212.	1.6	57
72	Interplay between mitogen-activated protein kinase and nitric oxide in brassinosteroid-induced pesticide metabolism in <i>Solanum lycopersicum</i> . <i>Journal of Hazardous Materials</i> , 2016, 316, 221-231.	6.5	39

#	ARTICLE	IF	CITATIONS
73	<i>DWARF</i> overexpression induces alteration in phytohormone homeostasis, development, architecture and carotenoid accumulation in tomato. <i>Plant Biotechnology Journal</i> , 2016, 14, 1021-1033.	4.1	83
74	Overexpression of a brassinosteroid biosynthetic gene Dwarf enhances photosynthetic capacity through activation of Calvin cycle enzymes in tomato. <i>BMC Plant Biology</i> , 2016, 16, 33.	1.6	57
75	Glutathione-mediated regulation of nitric oxide, S-nitrosothiol and redox homeostasis confers cadmium tolerance by inducing transcription factors and stress response genes in tomato. <i>Chemosphere</i> , 2016, 161, 536-545.	4.2	111
76	Melatonin enhances thermotolerance by promoting cellular protein protection in tomato plants. <i>Journal of Pineal Research</i> , 2016, 61, 457-469.	3.4	216
77	Systemic induction of photosynthesis via illumination of the shoot apex is mediated by phytochrome B. <i>Plant Physiology</i> , 2016, 172, pp.01202.2016.	2.3	73
78	Neglecting legumes has compromised human health and sustainable food production. <i>Nature Plants</i> , 2016, 2, 16112.	4.7	529
79	Role of Hormones in Plant Adaptation to Heat Stress. , 2016, , 1-21.		23
80	Interactions between 2-Cys peroxiredoxins and ascorbate in autophagosome formation during the heat stress response in <i>Solanum lycopersicum</i> . <i>Journal of Experimental Botany</i> , 2016, 67, 1919-1933.	2.4	34
81	Phytochrome A and B Function Antagonistically to Regulate Cold Tolerance via Abscisic Acid-Dependent Jasmonate Signaling. <i>Plant Physiology</i> , 2016, 170, 459-471.	2.3	216
82	Guard cell hydrogen peroxide and nitric oxide mediate elevated $CO_2$ -induced stomatal movement in tomato. <i>New Phytologist</i> , 2015, 208, 342-353.	3.5	95
83	Involvement of nitric oxide in the jasmonate-dependent basal defense against root-knot nematode in tomato plants. <i>Frontiers in Plant Science</i> , 2015, 6, 193.	1.7	57
84	Melatonin mitigates cadmium phytotoxicity through modulation of phytochelatin biosynthesis, vacuolar sequestration, and antioxidant potential in <i>Solanum lycopersicum</i> L. <i>Frontiers in Plant Science</i> , 2015, 6, 601.	1.7	278
85	Tomato HsfA1a plays a critical role in plant drought tolerance by activating <i>ATG</i> genes and inducing autophagy. <i>Autophagy</i> , 2015, 11, 2033-2047.	4.3	166
86	The relationship between the plant-encoded RNA-dependent RNA polymerase 1 and alternative oxidase in tomato basal defense against Tobacco mosaic virus. <i>Planta</i> , 2015, 241, 641-650.	1.6	18
87	Salicylic acid binding of mitochondrial alpha-ketoglutarate dehydrogenase E2 affects mitochondrial oxidative phosphorylation and electron transport chain components and plays a role in basal defense against tobacco mosaic virus in tomato. <i>New Phytologist</i> , 2015, 205, 1296-1307.	3.5	55
88	Characterization of the promoter and extended C-terminal domain of Arabidopsis WRKY33 and functional analysis of tomato WRKY33 homologues in plant stress responses. <i>Journal of Experimental Botany</i> , 2015, 66, 4567-4583.	2.4	86
89	Brassinosteroids play a critical role in the regulation of pesticide metabolism in crop plants. <i>Scientific Reports</i> , 2015, 5, 9018.	1.6	110
90	NPR1-dependent salicylic acid signaling is not involved in elevated $CO_2$ -induced heat stress tolerance in <i>Arabidopsis thaliana</i> . <i>Plant Signaling and Behavior</i> , 2015, 10, e1011944.	1.2	13

#	ARTICLE	IF	CITATIONS
91	RNA-seq analysis reveals the role of red light in resistance against <i>Pseudomonas syringae</i> pv. tomato DC3000 in tomato plants. <i>BMC Genomics</i> , 2015, 16, 120.	1.2	82
92	Phosphorus and magnesium interactively modulate the elongation and directional growth of primary roots in <i>Arabidopsis thaliana</i> (L.) Heynh. <i>Journal of Experimental Botany</i> , 2015, 66, 3841-3854.	2.4	35
93	Interplay between reactive oxygen species and hormones in the control of plant development and stress tolerance. <i>Journal of Experimental Botany</i> , 2015, 66, 2839-2856.	2.4	572
94	Enhanced photosynthetic capacity and antioxidant potential mediate brassinosteroid-induced phenanthrene stress tolerance in tomato. <i>Environmental Pollution</i> , 2015, 201, 58-66.	3.7	37
95	Antagonism between phytohormone signalling underlies the variation in disease susceptibility of tomato plants under elevated CO <sub>2</sub> . <i>Journal of Experimental Botany</i> , 2015, 66, 1951-1963.	2.4	116
96	High atmospheric carbon dioxide-dependent alleviation of salt stress is linked to RESPIRATORY BURST OXIDASE 1 ( <i>RBOH1</i> )-dependent H <sub>2</sub> O <sub>2</sub> production in tomato ( <i>Solanum</i> ) Tj ETQ 2010 0 rg BT/Overlock	2.4	119
97	Application of 24-epibrassinolide decreases the susceptibility to cucumber mosaic virus in zucchini ( <i>Cucurbita pepo</i> L). <i>Scientia Horticulturae</i> , 2015, 195, 116-123.	1.7	14
98	Light-induced systemic resistance in tomato plants against root-knot nematode <i>Meloidogyne incognita</i> . <i>Plant Growth Regulation</i> , 2015, 76, 167-175.	1.8	22
99	Tomato- <i>Pseudomonas syringae</i> interactions under elevated CO <sub>2</sub> concentration: the role of stomata. <i>Journal of Experimental Botany</i> , 2015, 66, 307-316.	2.4	40
100	Role of Brassinosteroid in Plant Adaptation to Abiotic Stresses and its Interplay with Other Hormones. <i>Current Protein and Peptide Science</i> , 2015, 16, 462-473.	0.7	86
101	<i>Arabidopsis</i> LIP5, a Positive Regulator of Multivesicular Body Biogenesis, Is a Critical Target of Pathogen-Responsive MAPK Cascade in Plant Basal Defense. <i>PLoS Pathogens</i> , 2014, 10, e1004243.	2.1	90
102	E3 Ubiquitin Ligase CHIP and NBR1-Mediated Selective Autophagy Protect Additively against Proteotoxicity in Plant Stress Responses. <i>PLoS Genetics</i> , 2014, 10, e1004116.	1.5	127
103	<i>RBOH1</i> -dependent H <sub>2</sub> O <sub>2</sub> production and subsequent activation of MPK1/2 play an important role in acclimation-induced cross-tolerance in tomato. <i>Journal of Experimental Botany</i> , 2014, 65, 595-607.	2.4	129
104	Chloroplastic thioredoxin-f and thioredoxin-m1/4 play important roles in brassinosteroids-induced changes in CO <sub>2</sub> assimilation and cellular redox homeostasis in tomato. <i>Journal of Experimental Botany</i> , 2014, 65, 4335-4347.	2.4	32
105	The perplexing role of autophagy in plant innate immune responses. <i>Molecular Plant Pathology</i> , 2014, 15, 637-645.	2.0	82
106	Role of $H_2O_2$ dynamics in brassinosteroid-induced stomatal closure and opening in <i>Solanum lycopersicum</i> . <i>Plant, Cell and Environment</i> , 2014, 37, 2036-2050.	2.8	139
107	Hydrogen peroxide mediates abscisic acid-induced HSP70 accumulation and heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2014, 37, 2768-2780.	2.8	135
108	The sub/supra-optimal temperature-induced inhibition of photosynthesis and oxidative damage in cucumber leaves are alleviated by grafting onto figleaf gourd/luffa rootstocks. <i>Physiologia Plantarum</i> , 2014, 152, 571-584.	2.6	39

#	ARTICLE	IF	CITATIONS
109	Effects of <i>Fusarium oxysporum</i> on rhizosphere microbial communities of two cucumber genotypes with contrasting <i>Fusarium</i> wilt resistance under hydroponic condition. <i>European Journal of Plant Pathology</i> , 2014, 140, 643-653.	0.8	10
110	H <sub>2</sub> O <sub>2</sub> mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses. <i>Journal of Experimental Botany</i> , 2014, 65, 4371-4383.	2.4	257
111	Role and regulation of autophagy in heat stress responses of tomato plants. <i>Frontiers in Plant Science</i> , 2014, 5, 174.	1.7	162
112	Plant-Soil Feedbacks and Soil Sickness: From Mechanisms to Application in Agriculture. <i>Journal of Chemical Ecology</i> , 2013, 39, 232-242.	0.9	248
113	Brassinosteroids accelerate recovery of photosynthetic apparatus from cold stress by balancing the electron partitioning, carboxylation and redox homeostasis in cucumber. <i>Physiologia Plantarum</i> , 2013, 148, 133-145.	2.6	107
114	Glutathione biosynthesis and regeneration play an important role in the metabolism of chlorothalonil in tomato. <i>Chemosphere</i> , 2013, 90, 2563-2570.	4.2	52
115	Role of brassinosteroids in alleviation of phenanthrene-cadmium co-contamination-induced photosynthetic inhibition and oxidative stress in tomato. <i>Journal of Experimental Botany</i> , 2013, 64, 199-213.	2.4	230
116	Flexible change and cooperation between mitochondrial electron transport and cytosolic glycolysis as the basis for chilling tolerance in tomato plants. <i>Planta</i> , 2013, 237, 589-601.	1.6	21
117	Protein-Protein Interactions in the Regulation of WRKY Transcription Factors. <i>Molecular Plant</i> , 2013, 6, 287-300.	3.9	276
118	Brassinosteroid alleviates polychlorinated biphenyls-induced oxidative stress by enhancing antioxidant enzymes activity in tomato. <i>Chemosphere</i> , 2013, 90, 2645-2653.	4.2	92
119	NBR1-Mediated Selective Autophagy Targets Insoluble Ubiquitinated Protein Aggregates in Plant Stress Responses. <i>PLoS Genetics</i> , 2013, 9, e1003196.	1.5	281
120	Stimulated Leaf Dark Respiration in Tomato in an Elevated Carbon Dioxide Atmosphere. <i>Scientific Reports</i> , 2013, 3, 3433.	1.6	72
121	Brassinosteroids-Induced Systemic Stress Tolerance was Associated with Increased Transcripts of Several Defence-Related Genes in the Phloem in <i>Cucumis sativus</i> . <i>PLoS ONE</i> , 2013, 8, e66582.	1.1	52
122	Cytokinin-Induced Parthenocarpic Fruit Development in Tomato Is Partly Dependent on Enhanced Gibberellin and Auxin Biosynthesis. <i>PLoS ONE</i> , 2013, 8, e70080.	1.1	79
123	The Role of Hydrogen Peroxide and Nitric Oxide in the Induction of Plant-Encoded RNA-Dependent RNA Polymerase 1 in the Basal Defense against Tobacco Mosaic Virus. <i>PLoS ONE</i> , 2013, 8, e76090.	1.1	45
124	Structural and Functional Analysis of VQ Motif-Containing Proteins in Arabidopsis as Interacting Proteins of WRKY Transcription Factors. <i>Plant Physiology</i> , 2012, 159, 810-825.	2.3	216
125	Brassinosteroid improves seed germination and early development of tomato seedling under phenanthrene stress. <i>Plant Growth Regulation</i> , 2012, 68, 87-96.	1.8	28
126	Interaction of Brassinosteroids and Polyamines Enhances Copper Stress Tolerance in <i>Raphanus Sativus</i> . <i>Journal of Experimental Botany</i> , 2012, 63, 5659-5675.	2.4	142



#	ARTICLE	IF	CITATIONS
127	Cellular glutathione redox homeostasis plays an important role in the brassinosteroid-induced increase in CO <sub>2</sub> assimilation in <i>Cucumis sativus</i> . <i>New Phytologist</i> , 2012, 194, 932-943.	3.5	120
128	Brassinosteroids induce plant tolerance against phenanthrene by enhancing degradation and detoxification in <i>Solanum lycopersicum</i> L.. <i>Ecotoxicology and Environmental Safety</i> , 2012, 80, 28-36.	2.9	68
129	The growth, photosynthesis and antioxidant defense responses of five vegetable crops to phenanthrene stress. <i>Ecotoxicology and Environmental Safety</i> , 2012, 80, 132-139.	2.9	82
130	Hydrogen peroxide functions as a secondary messenger for brassinosteroids-induced CO <sub>2</sub> assimilation and carbohydrate metabolism in <i>Cucumis sativus</i> . <i>Journal of Zhejiang University: Science B</i> , 2012, 13, 811-823.	1.3	45
131	Benefits of brassinosteroid crosstalk. <i>Trends in Plant Science</i> , 2012, 17, 594-605.	4.3	271
132	A Combined Approach of High-Throughput Sequencing and Degradome Analysis Reveals Tissue Specific Expression of MicroRNAs and Their Targets in Cucumber. <i>PLoS ONE</i> , 2012, 7, e33040.	1.1	113
133	Brassinosteroid alleviates phenanthrene and pyrene phytotoxicity by increasing detoxification activity and photosynthesis in tomato. <i>Chemosphere</i> , 2012, 86, 546-555.	4.2	110
134	Interactive effects of CO <sub>2</sub> enrichment and brassinosteroid on CO <sub>2</sub> assimilation and photosynthetic electron transport in <i>Cucumis sativus</i> . <i>Environmental and Experimental Botany</i> , 2012, 75, 98-106.	2.0	43
135	Temperature effects on the reactive oxygen species formation and antioxidant defence in roots of two cucurbit species with contrasting root zone temperature optima. <i>Acta Physiologiae Plantarum</i> , 2012, 34, 713-720.	1.0	20
136	The reduction of reactive oxygen species formation by mitochondrial alternative respiration in tomato basal defense against TMV infection. <i>Planta</i> , 2012, 235, 225-238.	1.6	46
137	Chromium Stress Mitigation by Polyamine-Brassinosteroid Application Involves Phytohormonal and Physiological Strategies in <i>Raphanus sativus</i> L.. <i>PLoS ONE</i> , 2012, 7, e33210.	1.1	159
138	Role of nitric oxide in hydrogen peroxide-dependent induction of abiotic stress tolerance by brassinosteroids in cucumber. <i>Plant, Cell and Environment</i> , 2011, 34, 347-358.	2.8	160
139	Induction of systemic stress tolerance by brassinosteroid in <i>Cucumis sativus</i> . <i>New Phytologist</i> , 2011, 191, 706-720.	3.5	124
140	Effects of nitrogen form on growth, CO <sub>2</sub> assimilation, chlorophyll fluorescence, and photosynthetic electron allocation in cucumber and rice plants. <i>Journal of Zhejiang University: Science B</i> , 2011, 12, 126-134.	1.3	49
141	Alleviation of chilling-induced oxidative damage by salicylic acid pretreatment and related gene expression in eggplant seedlings. <i>Plant Growth Regulation</i> , 2011, 65, 101-108.	1.8	63
142	<i>Arabidopsis</i> Sigma Factor Binding Proteins Are Activators of the WRKY33 Transcription Factor in Plant Defense. <i>Plant Cell</i> , 2011, 23, 3824-3841.	3.1	260
143	Systemic Induction and Role of Mitochondrial Alternative Oxidase and Nitric Oxide in a Compatible Tomato-Tobacco mosaic virus Interaction. <i>Molecular Plant-Microbe Interactions</i> , 2010, 23, 39-48.	1.4	85
144	Induction and origin of adventitious roots from chimeras of <i>Brassica juncea</i> and <i>Brassica oleracea</i> . <i>Plant Cell, Tissue and Organ Culture</i> , 2010, 101, 287-294.	1.2	13

#	ARTICLE	IF	CITATIONS
145	Functional Analysis of the Arabidopsis <i>PAL</i> Gene Family in Plant Growth, Development, and Response to Environmental Stress. <i>Plant Physiology</i> , 2010, 153, 1526-1538.	2.3	668
146	ROS mediate brassinosteroids-induced plant stress responses. <i>Plant Signaling and Behavior</i> , 2010, 5, 532-534.	1.2	24
147	Photoprotective roles of anthocyanins in <i>Begonia semperflorens</i> . <i>Plant Science</i> , 2010, 179, 202-208.	1.7	93
148	The different responses of glutathione-dependent detoxification pathway to fungicide chlorothalonil and carbendazim in tomato leaves. <i>Chemosphere</i> , 2010, 79, 958-965.	4.2	49
149	Degradation of chlorpyrifos in laboratory soil and its impact on soil microbial functional diversity. <i>Journal of Environmental Sciences</i> , 2009, 21, 380-386.	3.2	103
150	Brassinosteroids promote photosynthesis and growth by enhancing activation of Rubisco and expression of photosynthetic genes in <i>Cucumis sativus</i> . <i>Planta</i> , 2009, 230, 1185-1196.	1.6	232
151	Grafting of <i>Cucumis sativus</i> onto <i>Cucurbita ficifolia</i> leads to improved plant growth, increased light utilization and reduced accumulation of reactive oxygen species in chilled plants. <i>Journal of Plant Research</i> , 2009, 122, 529-540.	1.2	44
152	Mitochondrial nad2 gene is co-transcribed with CMS-associated orfB gene in cytoplasmic male-sterile stem mustard ( <i>Brassica juncea</i> ). <i>Molecular Biology Reports</i> , 2009, 36, 345-351.	1.0	20
153	Genetic and heterosis analysis for important agronomic traits of Chinese vegetable mustard ( <i>Brassica</i> ) Tj ETQq1 1 0,784314 1gBT /Ov	0.5	0
154	Effects of Phenylcarboxylic Acids on Mitosis, Endoreduplication and Expression of Cell Cycle-Related Genes in Roots of Cucumber ( <i>Cucumis sativus</i> L.). <i>Journal of Chemical Ecology</i> , 2009, 35, 679-688.	0.9	27
155	Selective trans-Cinnamic Acid Uptake Impairs [Ca <sup>2+</sup> ] <sub>cyt</sub> Homeostasis and Growth in <i>Cucumis sativus</i> L.. <i>Journal of Chemical Ecology</i> , 2009, 35, 1471-1477.	0.9	20
156	Effects of cucumber mosaic virus infection on electron transport and antioxidant system in chloroplasts and mitochondria of cucumber and tomato leaves. <i>Physiologia Plantarum</i> , 2009, 135, 246-257.	2.6	82
157	Effects of light quality on CO <sub>2</sub> assimilation, chlorophyll-fluorescence quenching, expression of Calvin cycle genes and carbohydrate accumulation in <i>Cucumis sativus</i> . <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2009, 96, 30-37.	1.7	226
158	Effects of calcium cyanamide on soil microbial communities and <i>Fusarium oxysporum</i> f. sp. <i>cucumerinum</i> . <i>Chemosphere</i> , 2009, 75, 872-877.	4.2	37
159	Detached leaves of tomato differ in their photosynthetic physiological response to moderate high and low temperature stress. <i>Scientia Horticulturae</i> , 2009, 123, 17-22.	1.7	49
160	Impact of Light Variation on Development of Photoprotection, Antioxidants, and Nutritional Value in <i>Lactuca sativa</i> L.. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 5494-5500.	2.4	53
161	Reactive Oxygen Species Are Involved in Brassinosteroid-Induced Stress Tolerance in Cucumber. <i>Plant Physiology</i> , 2009, 150, 801-814.	2.3	640
162	Effects of Root and Foliar Applications of 24-Epibrassinolide on <i>Fusarium</i> Wilt and Antioxidant Metabolism in Cucumber Roots. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2009, 44, 1340-1345.	0.5	20

#	ARTICLE	IF	CITATIONS
163	MADS-box genes are associated with cytoplasmic homeosis in cytoplasmic male-sterile stem mustard as partially mimicked by specifically inhibiting mtETC. <i>Plant Growth Regulation</i> , 2008, 56, 191-201.	1.8	12
164	Decreased energy synthesis is partially compensated by a switch to sucrose synthase pathway of sucrose degradation in restricted root of tomato plants. <i>Plant Physiology and Biochemistry</i> , 2008, 46, 1040-1044.	2.8	16
165	Relationship between cytoplasmic male sterility and <i>SPL</i> -like gene expression in stem mustard. <i>Physiologia Plantarum</i> , 2008, 133, 426-434.	2.6	10
166	Root restriction-induced limitation to photosynthesis in tomato ( <i>Lycopersicon esculentum</i> Mill.) leaves. <i>Scientia Horticulturae</i> , 2008, 117, 197-202.	1.7	30
167	Mitochondrial retrograde regulation tuning fork in nuclear genes expressions of higher plants. <i>Journal of Genetics and Genomics</i> , 2008, 35, 65-71.	1.7	26
168	Chill-Induced Decrease in Capacity of RuBP Carboxylation and Associated H <sub>2</sub> O <sub>2</sub> Accumulation in Cucumber Leaves are Alleviated by Grafting onto Figleaf Gourd. <i>Annals of Botany</i> , 2007, 100, 839-848.	1.4	90
169	Alterations of RNA Editing for the Mitochondrial ATP9 Gene in a New orf220-type Cytoplasmic Male-sterile Line of Stem Mustard ( <i>Brassica juncea</i> var. <i>tumida</i> ). <i>Journal of Integrative Plant Biology</i> , 2007, 49, 672-677.	4.1	10
170	Low O <sub>2</sub> supply is involved in the poor growth in root-restricted plants of tomato ( <i>Lycopersicon</i> ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 46	2.0	40
171	Adaptation of cucurbit species to changes in substrate temperature: Root growth, antioxidants, and peroxidation. <i>Journal of Plant Biology</i> , 2007, 50, 527-532.	0.9	17
172	Diurnal variations in gas exchange, chlorophyll fluorescence quenching and light allocation in soybean leaves: The cause for midday depression in CO <sub>2</sub> assimilation. <i>Scientia Horticulturae</i> , 2006, 110, 214-218.	1.7	35
173	Different effects of chilling on respiration in leaves and roots of cucumber ( <i>Cucumis sativus</i> ). <i>Plant Physiology and Biochemistry</i> , 2006, 44, 837-843.	2.8	29
174	Effect of excess manganese on the antioxidant system in <i>Cucumis sativus</i> L. under two light intensities. <i>Environmental and Experimental Botany</i> , 2006, 58, 197-205.	2.0	73
175	Genotypic Variation of Rubisco Expression, Photosynthetic Electron Flow and Antioxidant Metabolism in the Chloroplasts of Chill-exposed Cucumber Plants. <i>Plant and Cell Physiology</i> , 2006, 47, 192-199.	1.5	87
176	Silicon-mediated alleviation of Mn toxicity in <i>Cucumis sativus</i> in relation to activities of superoxide dismutase and ascorbate peroxidase. <i>Phytochemistry</i> , 2005, 66, 1551-1559.	1.4	216
177	Effects of pollination and N-(2-chloro-4-pyridyl)-N-phenylurea on the expression of acid invertase in ovaries of <i>Lagenaria leucantha</i> . <i>Plant Growth Regulation</i> , 2004, 42, 263-270.	1.8	1
178	Effects of Potato Virus Y <sup>NTN</sup> Infection on Gas Exchange and Photosystem 2 Function in Leaves of <i>Solanum tuberosum</i> L. <i>Photosynthetica</i> , 2004, 42, 417-423.	0.9	39
179	Incidence of Fusarium wilt in <i>Cucumis sativus</i> L. is promoted by cinnamic acid, an autotoxin in root exudates. <i>Plant and Soil</i> , 2004, 263, 143-150.	1.8	128
180	Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber ( <i>Cucumis sativus</i> L.). <i>Plant Science</i> , 2004, 167, 527-533.	1.7	703

#	ARTICLE	IF	CITATIONS
181	Greenhouse and field cucumber genotypes use different mechanisms to protect against dark chilling. <i>Functional Plant Biology</i> , 2004, 31, 1215.	1.1	14
182	Expression of CycD3 is transiently increased by pollination and N-(2-chloro-4-pyridyl)-N <sup>ε</sup> -phenylurea in ovaries of <i>Lagenaria leucantha</i> . <i>Journal of Experimental Botany</i> , 2003, 54, 1245-1251.	2.4	22
183	Chill-Induced Inhibition of Photosynthesis: Genotypic Variation within <i>Cucumis sativus</i> . <i>Plant and Cell Physiology</i> , 2002, 43, 1182-1188.	1.5	50
184	Effects of Simulated Acid Precipitation on Photosynthesis, Chlorophyll Fluorescence, and Antioxidative Enzymes in <i>Cucumis sativus</i> L.. <i>Photosynthetica</i> , 2002, 40, 331-335.	0.9	25
185	Hinoki ( <i>Chamaecyparis obtusa</i> ) bark, a substrate with anti-pathogen properties that suppress some root diseases of tomato. <i>Scientia Horticulturae</i> , 1999, 81, 13-24.	1.7	34