

Ana Sofia Cerdeira

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6053242/ana-sofia-cerdeira-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

17 papers	383 citations	8 h-index	18 g-index
18 ext. papers	517 ext. citations	8.9 avg, IF	3.68 L-index

#	Paper	IF	Citations
17	Placental Vesicles Carry Active Endothelial Nitric Oxide Synthase and Their Activity is Reduced in Preeclampsia. <i>Hypertension</i> , 2017 , 70, 372-381	8.5	85
16	Meta-Analysis and Systematic Review to Assess the Role of Soluble FMS-Like Tyrosine Kinase-1 and Placenta Growth Factor Ratio in Prediction of Preeclampsia: The SaPPPhirE Study. <i>Hypertension</i> , 2018 , 71, 306-316	8.5	83
15	Placental Syncytiotrophoblast-Derived Extracellular Vesicles Carry Active NEP (Neprilysin) and Are Increased in Preeclampsia. <i>Hypertension</i> , 2019 , 73, 1112-1119	8.5	50
14	Randomized Interventional Study on Prediction of Preeclampsia/Eclampsia in Women With Suspected Preeclampsia: INSPIRE. <i>Hypertension</i> , 2019 , 74, 983-990	8.5	43
13	Plasma concentrations of soluble endoglin versus standard evaluation in patients with suspected preeclampsia. <i>PLoS ONE</i> , 2012 , 7, e48259	3.7	41
12	Predictive Performance of PlGF (Placental Growth Factor) for Screening Preeclampsia in Asymptomatic Women: A Systematic Review and Meta-Analysis. <i>Hypertension</i> , 2019 , 74, 1124-1135	8.5	30
11	Maternal circulating syncytiotrophoblast-derived extracellular vesicles contain biologically active 5WtRNA halves. <i>Biochemical and Biophysical Research Communications</i> , 2019 , 518, 107-113	3.4	19
10	The role of angiogenic factors in the management of preeclampsia. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019 , 98, 700-707	3.8	17
9	Circulating soluble fms-like tyrosine kinase-1 is placentally derived in normal pregnancy: First in vivo evidence. <i>Pregnancy Hypertension</i> , 2019 , 16, 145-147	2.6	5
8	Seeing red degeneration in uterine fibroids in pregnancy: proceed with caution. <i>Lancet, The</i> , 2019 , 394, e37	4.0	3
7	Performance of soluble fms-like tyrosine kinase-1-to-placental growth factor ratio of 85 for ruling in preeclampsia within 4 weeks. <i>American Journal of Obstetrics and Gynecology</i> , 2021 , 224, 322-323	6.4	3
6	In vivo evidence of significant placental growth factor release by normal pregnancy placentas. <i>Scientific Reports</i> , 2020 , 10, 132	4.9	2
5	Ruling out preeclampsia in the next 4 weeks using a soluble fms-like tyrosine kinase 1/placental growth factor ratio 88: secondary analysis of the Interventional Study on Prediction of Preeclampsia/Eclampsia in Women With Suspected Preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2021 ,	6.4	1
4	Paracetamol use in pregnancy - neglecting context promotes misinterpretation.. <i>Nature Reviews Endocrinology</i> , 2022 ,	15.2	1
3	Retroperitoneal leiomyomatosis: a benign outcome of power morcellation with potentially serious consequences.. <i>Lancet, The</i> , 2022 , 399, 554	4.0	
2	The value of MRI in management of uterine fibroids in pregnancy. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2021 , 256, 522-523	2.4	
1	The diagnostic value of angiogenic and antiangiogenic factors in differential diagnosis of preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2021 ,	6.4	

