

# Hui Quan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6059155/publications.pdf>

Version: 2024-02-01

7  
papers

122  
citations

1478505  
6  
h-index

1720034  
7  
g-index

7  
all docs

7  
docs citations

7  
times ranked

226  
citing authors

#	ARTICLE	IF	CITATIONS
1	Resveratrol suppresses the reprogramming of macrophages into an endotoxin-tolerant state through the activation of AMP-activated protein kinase. <i>European Journal of Pharmacology</i> , 2021, 899, 173993.	3.5	9
2	Ginsenoside Rb1 increases macrophage phagocytosis through p38 mitogen-activated protein kinase/Akt pathway. <i>Journal of Ginseng Research</i> , 2019, 43, 394-401.	5.7	27
3	Ginsenoside Rg3 promotes Fc gamma receptor-mediated phagocytosis of bacteria by macrophages via an extracellular signal-regulated kinase 1/2 and p38 mitogen-activated protein kinase-dependent mechanism. <i>International Immunopharmacology</i> , 2019, 77, 105945.	3.8	23
4	Stearoyl lysophosphatidylcholine inhibits LPS-induced extracellular release of HMGB1 through the G2A/calcium/CaMKK $\beta$ /AMPK pathway. <i>European Journal of Pharmacology</i> , 2019, 852, 125-133.	3.5	13
5	Stearoyl lysophosphatidylcholine enhances the phagocytic ability of macrophages through the AMP-activated protein kinase/p38 mitogen activated protein kinase pathway. <i>International Immunopharmacology</i> , 2016, 39, 328-334.	3.8	15
6	AICAR Enhances the Phagocytic Ability of Macrophages towards Apoptotic Cells through P38 Mitogen Activated Protein Kinase Activation Independent of AMP-Activated Protein Kinase. <i>PLoS ONE</i> , 2015, 10, e0127885.	2.5	19
7	Stearoyl lysophosphatidylcholine prevents lipopolysaccharide-induced extracellular release of high mobility group box-1 through AMP-activated protein kinase activation. <i>International Immunopharmacology</i> , 2015, 28, 540-545.	3.8	16