

Delphine Javelaud

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6068398/publications.pdf>

Version: 2024-02-01

41
papers

3,576
citations

218677

26
h-index

361022

35
g-index

43
all docs

43
docs citations

43
times ranked

5348
citing authors

#	ARTICLE	IF	CITATIONS
1	Large-scale pan-cancer analysis reveals broad prognostic association between TGF- β ligands, not Hedgehog, and GLI1/2 expression in tumors. <i>Scientific Reports</i> , 2020, 10, 14491.	3.3	10
2	GLI1/GLI2 functional interplay is required to control Hedgehog/GLI targets gene expression. <i>Biochemical Journal</i> , 2020, 477, 3131-3145.	3.7	23
3	Transcriptional repression of the tyrosinase-related protein 2 gene by transforming growth factor- β and the Kruppel-like transcription factor GLI2. <i>Journal of Dermatological Science</i> , 2019, 94, 321-329.	1.9	2
4	How Bad Is the Hedgehog? GLI-Dependent, Hedgehog-Independent Cancers on the Importance of Biomarkers for Proper Patients Selection. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2018, 19, S87-S88.	0.8	2
5	487 Expression of metastasis and invasion related molecules in malignant melanoma. <i>Journal of Investigative Dermatology</i> , 2016, 136, S243.	0.7	1
6	Cell Density Sensing Alters TGF- β Signaling in a Cell-Type-Specific Manner, Independent from Hippo Pathway Activation. <i>Developmental Cell</i> , 2015, 32, 640-651.	7.0	59
7	Abstract LB-028: Cell density sensing alters TGF-beta signaling in a cell type-specific manner, independent from Hippo pathway activation. , 2015, , .		0
8	<scp>GLI</scp>2 cooperates with <scp>ZEB</scp>1 for transcriptional repression of <scp><i>CDH1</i></scp> expression in human melanoma cells. <i>Pigment Cell and Melanoma Research</i> , 2013, 26, 861-873.	3.3	30
9	Overlapping activities of TGF- β and Hedgehog signaling in cancer: Therapeutic targets for cancer treatment. , 2013, 137, 183-199.		51
10	Insights into the Transforming Growth Factor- β Signaling Pathway in Cutaneous Melanoma. <i>Annals of Dermatology</i> , 2013, 25, 135.	0.9	72
11	The Role of TGF- β in Cutaneous Melanoma Biology. , 2013, , 235-254.		0
12	Expression of Microphthalmia-associated Transcription Factor (MITF), Which Is Critical for Melanoma Progression, Is Inhibited by Both Transcription Factor GLI2 and Transforming Growth Factor- β . <i>Journal of Biological Chemistry</i> , 2012, 287, 17996-18004.	3.4	84
13	Halofuginone Inhibits the Establishment and Progression of Melanoma Bone Metastases. <i>Cancer Research</i> , 2012, 72, 6247-6256.	0.9	66
14	Systematic classification of melanoma cells by phenotype-specific gene expression mapping. <i>Pigment Cell and Melanoma Research</i> , 2012, 25, 343-353.	3.3	155
15	Crosstalk between TGF- β and hedgehog signaling in cancer. <i>FEBS Letters</i> , 2012, 586, 2016-2025.	2.8	135
16	GLI2 and MITF transcription factors control exclusive gene expression programs and inversely regulate invasion in human melanoma cells. <i>Pigment Cell and Melanoma Research</i> , 2011, 24, 932-943.	3.3	71
17	Efficient TGF- β /SMAD signaling in human melanoma cells associated with high c-SKI/SnoN expression. <i>Molecular Cancer</i> , 2011, 10, 2.	19.2	46
18	Correction: TGF- β -RI Kinase Inhibitor SD-208 Reduces the Development and Progression of Melanoma Bone Metastases. <i>Cancer Research</i> , 2011, 71, 2023-2023.	0.9	0

#	ARTICLE	IF	CITATIONS
19	TGF- β 2/SMAD/GLI2 Signaling Axis in Cancer Progression and Metastasis. <i>Cancer Research</i> , 2011, 71, 5606-5610.	0.9	182
20	TGF- β 2-RI Kinase Inhibitor SD-208 Reduces the Development and Progression of Melanoma Bone Metastases. <i>Cancer Research</i> , 2011, 71, 175-184.	0.9	203
21	GLI2-Mediated Melanoma Invasion and Metastasis. <i>Journal of the National Cancer Institute</i> , 2010, 102, 1148-1159.	6.3	149
22	Smad7 restricts melanoma invasion by restoring N-cadherin expression and establishing heterotypic cell-cell interactions in vivo. <i>Pigment Cell and Melanoma Research</i> , 2010, 23, 795-808.	3.3	24
23	Abstract LB-240: Smad7 blocks melanoma invasion by suppressing n-cadherin cleavage and preserving heterotypic cell-cell interactions in vivo.., 2010, , .		0
24	Transforming growth factor- β 2 in cutaneous melanoma. <i>Pigment Cell and Melanoma Research</i> , 2008, 21, 123-132.	3.3	125
25	JNK supports survival in melanoma cells by controlling cell cycle arrest and apoptosis. <i>Pigment Cell and Melanoma Research</i> , 2008, 21, 429-438.	3.3	51
26	Response to the letter by Reed et al.. <i>Pigment Cell and Melanoma Research</i> , 2008, 21, 496-497.	3.3	2
27	Transforming Growth Factor- β 2 Suppresses the Ability of Ski to Inhibit Tumor Metastasis by Inducing Its Degradation. <i>Cancer Research</i> , 2008, 68, 3277-3285.	0.9	94
28	Stable Overexpression of Smad7 in Human Melanoma Cells Impairs Bone Metastasis. <i>Cancer Research</i> , 2007, 67, 2317-2324.	0.9	187
29	Stable overexpression of Smad7 in human melanoma cells inhibits bone metastasis. <i>Melanoma Research</i> , 2006, 16, S93.	1.2	0
30	Interplays Between The Smad and Map Kinase Signaling Pathways. , 2006, , 317-334.		2
31	Stable overexpression of Smad7 in human melanoma cells inhibits their tumorigenicity in vitro and in vivo. <i>Oncogene</i> , 2005, 24, 7624-7629.	5.9	100
32	Crosstalk mechanisms between the mitogen-activated protein kinase pathways and Smad signaling downstream of TGF- β 2: implications for carcinogenesis. <i>Oncogene</i> , 2005, 24, 5742-5750.	5.9	373
33	Amelioration of Radiation-induced Fibrosis. <i>Journal of Biological Chemistry</i> , 2004, 279, 15167-15176.	3.4	187
34	NF- κ B activation prevents apoptotic oxidative stress via an increase of both thioredoxin and MnSOD levels in TNF- α -treated Ewing sarcoma cells. <i>FEBS Letters</i> , 2004, 578, 111-115.	2.8	109
35	Mammalian transforming growth factor- β 2s: Smad signaling and physio-pathological roles. <i>International Journal of Biochemistry and Cell Biology</i> , 2004, 36, 1161-1165.	2.8	153
36	TGF- β 2-induced SMAD signaling and gene regulation: consequences for extracellular matrix remodeling and wound healing. <i>Journal of Dermatological Science</i> , 2004, 35, 83-92.	1.9	392

#	ARTICLE	IF	CITATIONS
37	Disruption of Basal JNK Activity Differentially Affects Key Fibroblast Functions Important for Wound Healing. <i>Journal of Biological Chemistry</i> , 2003, 278, 24624-24628.	3.4	103
38	Inactivation of p21 Sensitizes Cells to Apoptosis via an Increase of Both p14ARF and p53 Levels and an Alteration of the Bax/Bcl-2 Ratio. <i>Journal of Biological Chemistry</i> , 2002, 277, 37949-37954.	3.4	107
39	Inhibition of constitutive NF- κ B activity suppresses tumorigenicity of ewing sarcoma EW7 cells. <i>International Journal of Cancer</i> , 2002, 98, 193-198.	5.1	25
40	NF- κ B activation results in rapid inactivation of JNK in TNF α -treated Ewing sarcoma cells: a mechanism for the anti-apoptotic effect of NF- κ B. <i>Oncogene</i> , 2001, 20, 4365-4372.	5.9	123
41	Induction of p21Waf1/Cip1 by TNF α requires NF- κ B activity and antagonizes apoptosis in Ewing tumor cells. <i>Oncogene</i> , 2000, 19, 61-68.	5.9	60