

# Amar Bahadur Singh

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6069560/publications.pdf>

Version: 2024-02-01

46  
papers

1,600  
citations

236612

25  
h-index

301761

39  
g-index

48  
all docs

48  
docs citations

48  
times ranked

2645  
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of p115 as a novel ACSL4 interacting protein and its role in regulating ACSL4 degradation. <i>Journal of Proteomics</i> , 2020, 229, 103926.	1.2	8
2	FXR activation promotes intestinal cholesterol excretion and attenuates hyperlipidemia in SR-BI-deficient mice fed a high-fat and high-cholesterol diet. <i>Physiological Reports</i> , 2020, 8, e14387.	0.7	7
3	Berberine decreases plasma triglyceride levels and upregulates hepatic TRIB1 in LDLR wild type mice and in LDLR deficient mice. <i>Scientific Reports</i> , 2019, 9, 15641.	1.6	11
4	Activation of FXR by obeticholic acid induces hepatic gene expression of SR-BI through a novel mechanism of transcriptional synergy with the nuclear receptor LXR. <i>International Journal of Molecular Medicine</i> , 2019, 43, 1927-1938.	1.8	11
5	Liver-specific knockdown of long-chain acyl-CoA synthetase 4 reveals its key role in VLDL-TG metabolism and phospholipid synthesis in mice fed a high-fat diet. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2019, 316, E880-E894.	1.8	19
6	Identification of a novel function of hepatic long-chain acyl-CoA synthetase-1 (ACSL1) in bile acid synthesis and its regulation by bile acid-activated farnesoid X receptor. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019, 1864, 358-371.	1.2	11
7	Farnesoid X Receptor Activation by Obeticholic Acid Elevates Liver Low-Density Lipoprotein Receptor Expression by mRNA Stabilization and Reduces Plasma Low-Density Lipoprotein Cholesterol in Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 2448-2459.	1.1	19
8	Hepatic HNF1 transcription factors control the induction of PCSK9 mediated by rosuvastatin in normolipidemic hamsters. <i>International Journal of Molecular Medicine</i> , 2017, 39, 749-756.	1.8	27
9	Identification of Hepatic Lysophosphatidylcholine Acyltransferase 3 as a Novel Target Gene Regulated by Peroxisome Proliferator-activated Receptor $\gamma$ . <i>Journal of Biological Chemistry</i> , 2017, 292, 884-897.	1.6	24
10	Regulation of lipid metabolism by obeticholic acid in hyperlipidemic hamsters. <i>Journal of Lipid Research</i> , 2017, 58, 350-363.	2.0	28
11	SREBP2 Activation Induces Hepatic Long-chain Acyl-CoA Synthetase 1 (ACSL1) Expression in Vivo and in Vitro through a Sterol Regulatory Element (SRE) Motif of the ACSL1 C-promoter. <i>Journal of Biological Chemistry</i> , 2016, 291, 5373-5384.	1.6	32
12	A novel peroxisome proliferator response element modulates hepatic low-density lipoprotein receptor gene transcription in response to PPAR $\gamma$ activation. <i>Biochemical Journal</i> , 2015, 472, 275-286.	1.7	12
13	PPAR $\gamma$ activation induces hepatic long-chain acyl-CoA synthetase 4 expression in vivo and in vitro. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2015, 1851, 577-587.	1.2	19
14	Inhibition of PCSK9 Transcription by Berberine Involves Down-regulation of Hepatic HNF1 $\alpha$ Protein Expression through the Ubiquitin-Proteasome Degradation Pathway. <i>Journal of Biological Chemistry</i> , 2015, 290, 4047-4058.	1.6	117
15	High-fructose feeding promotes accelerated degradation of hepatic LDL receptor and hypercholesterolemia in hamsters via elevated circulating PCSK9 levels. <i>Atherosclerosis</i> , 2015, 239, 364-374.	0.4	29
16	Reduction of circulating PCSK9 and LDL-C levels by liver-specific knockdown of HNF1 $\alpha$ in normolipidemic mice. <i>Journal of Lipid Research</i> , 2015, 56, 801-809.	2.0	48
17	Arachidonic acid downregulates acyl-CoA synthetase 4 expression by promoting its ubiquitination and proteasomal degradation. <i>Journal of Lipid Research</i> , 2014, 55, 1657-1667.	2.0	43
18	The Critical Role of mRNA Destabilizing Protein Heterogeneous Nuclear Ribonucleoprotein D in 3 $\alpha$ -OH Untranslated Region-Mediated Decay of Low-Density Lipoprotein Receptor mRNA in Liver Tissue. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014, 34, 8-16.	1.1	23

#	ARTICLE	IF	CITATIONS
19	A novel posttranscriptional mechanism for dietary cholesterol-mediated suppression of liver LDL receptor expression. <i>Journal of Lipid Research</i> , 2014, 55, 1397-1407.	2.0	24
20	CETP inhibitors downregulate hepatic LDL receptor and PCSK9 expression in vitro and in vivo through a SREBP2 dependent mechanism. <i>Atherosclerosis</i> , 2014, 235, 449-462.	0.4	49
21	Activation of retinoid receptor-mediated signaling ameliorates diabetes-induced cardiac dysfunction in Zucker diabetic rats. <i>Journal of Molecular and Cellular Cardiology</i> , 2013, 57, 106-118.	0.9	37
22	Retinoic acid protects cardiomyocytes from high glucose-induced apoptosis through inhibition of NF- $\kappa$ B signaling Pathway. <i>Journal of Cellular Physiology</i> , 2013, 228, 380-392.	2.0	42
23	High-fructose diet downregulates long-chain acyl-CoA synthetase 3 expression in liver of hamsters via impairing LXR/RXR signaling pathway. <i>Journal of Lipid Research</i> , 2013, 54, 1241-1254.	2.0	38
24	Anti-diabetic and anti-oxidative effects of 4-hydroxypipicolinic acid in C57BL/KsJ-db/db mice. <i>Human and Experimental Toxicology</i> , 2012, 31, 57-65.	1.1	20
25	Coagulanolide modulates hepatic glucose metabolism in C57BL/KsJ-db/db mice. <i>Human and Experimental Toxicology</i> , 2012, 31, 1056-1065.	1.1	17
26	Flavone-Based Novel Antidiabetic and Antidyslipidemic Agents. <i>Journal of Medicinal Chemistry</i> , 2012, 55, 4551-4567.	2.9	37
27	High Glucose-induced repression of RAR/RXR in cardiomyocytes is mediated through oxidative stress/JNK signaling. <i>Journal of Cellular Physiology</i> , 2012, 227, 2632-2644.	2.0	44
28	Differential effects of formononetin and cladrin on osteoblast function, peak bone mass achievement and bioavailability in rats. <i>Journal of Nutritional Biochemistry</i> , 2011, 22, 318-327.	1.9	69
29	Tectone, a New Antihyperglycemic Anthraquinone from <i>Tectona grandis</i> Leaves. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	10
30	Antihyperglycaemic effect of an unusual amino acid (4-hydroxyisoleucine) in C57BL/KsJ-db/db mice. <i>Natural Product Research</i> , 2010, 24, 258-265.	1.0	49
31	db/+ Mice as an Alternate Model in Antidiabetic Drug Discovery Research. <i>Archives of Medical Research</i> , 2009, 40, 73-78.	1.5	8
32	Methoxylated isoflavones, cajanin and isoformononetin, have non-estrogenic bone forming effect via differential mitogen activated protein kinase (MAPK) signaling. <i>Journal of Cellular Biochemistry</i> , 2009, 108, 388-399.	1.2	85
33	Novel 2-aryl-naphtho[1,2-d]oxazole derivatives as potential PTP-1B inhibitors showing antihyperglycemic activities. <i>European Journal of Medicinal Chemistry</i> , 2009, 44, 109-116.	2.6	48
34	Synthesis of $\pm$ -amyrin derivatives and their in vivo antihyperglycemic activity. <i>European Journal of Medicinal Chemistry</i> , 2009, 44, 1215-1222.	2.6	31
35	Design and synthesis of 3,5-diarylisoxazole derivatives as novel class of anti-hyperglycemic and lipid lowering agents. <i>Bioorganic and Medicinal Chemistry</i> , 2009, 17, 5285-5292.	1.4	50
36	Synthesis of protein tyrosine phosphatase 1B inhibitors: Model validation and docking studies. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 2320-2323.	1.0	17

#	ARTICLE	IF	CITATIONS
37	5,6-Diarylanthranilo-1,3-dinitriles as a new class of antihyperglycemic agents. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 2158-2161.	1.0	37
38	Synthesis of novel triterpenoid (lupeol) derivatives and their in vivo antihyperglycemic and antidyslipidemic activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 4463-4466.	1.0	63
39	Pyranocoumarins: A new class of anti-hyperglycemic and anti-dyslipidemic agents. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 6447-6451.	1.0	129
40	Novel class of hybrid natural products as antidiabetic agents. <i>Natural Product Research</i> , 2009, 23, 60-69.	1.0	9
41	Antihyperglycaemic activity of $\hat{I}\pm$ -amyrin acetate in rats and db/db mice. <i>Natural Product Research</i> , 2009, 23, 876-882.	1.0	35
42	Synthesis of 3,5-disubstituted isoxazolines as protein tyrosine phosphatase 1B inhibitors. <i>Medicinal Chemistry Research</i> , 2008, 17, 123-136.	1.1	10
43	Preliminary studies on the hypoglycemic effect of <i>Peganum harmala</i> L. Seeds ethanol extract on normal and streptozotocin induced diabetic rats. <i>Indian Journal of Clinical Biochemistry</i> , 2008, 23, 391-393.	0.9	36
44	Synthesis and antihyperglycemic activity of novel N-acyl-2-arylethylamines and N-acyl-3-coumarylamines. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008, 18, 2301-2305.	1.0	25
45	Coagulanolide, a withanolide from <i>Withania coagulans</i> fruits and antihyperglycemic activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008, 18, 6534-6537.	1.0	68
46	Chalcone based aryloxypropanolamines as potential antihyperglycemic agents. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2007, 17, 799-802.	1.0	25