

David Esteban

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/607815/david-esteban-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

151
papers

5,732
citations

35
h-index

72
g-index

155
ext. papers

6,153
ext. citations

4.9
avg, IF

5.59
L-index

#	Paper	IF	Citations
151	Rigidified Derivative of the Non-macrocyclic Ligand HOCTAPA for Stable Lanthanide(III) Complexation.. <i>Inorganic Chemistry</i> , 2022 ,	5.1	3
150	Prediction of Gd(III) complex thermodynamic stability. <i>Coordination Chemistry Reviews</i> , 2022 , 467, 214606	5.2	1
149	Rigid versions of PDTA incorporating a 1,3-diaminocyclobutyl spacer for Mn complexation: stability, water exchange dynamics and relaxivity. <i>Dalton Transactions</i> , 2021 , 50, 16290-16303	4.3	0
148	Understanding the Effect of the Electron Spin Relaxation on the Relaxivities of Mn(II) Complexes with Triazacyclononane Derivatives. <i>Inorganic Chemistry</i> , 2021 , 60, 15055-15068	5.1	1
147	Complexation of Mn(II) by Rigid PycLen Diacetates: Equilibrium, Kinetic, Relaxometric, Density Functional Theory, and Superoxide Dismutase Activity Studies. <i>Inorganic Chemistry</i> , 2021 , 60, 1133-1148	5.1	12
146	Expanding the Ligand Classes Used for Mn(II) Complexation: Oxa-aza Macrocycles Make the Difference. <i>Molecules</i> , 2021 , 26,	4.8	2
145	Complexation of -Functionalized Cyclams with Copper(II) and Zinc(II): Similarities and Changes When Compared to Parent Cyclam Analogues. <i>Inorganic Chemistry</i> , 2021 , 60, 10857-10872	5.1	0
144	Paramagnetic chemical exchange saturation transfer agents and their perspectives for application in magnetic resonance imaging. <i>International Reviews in Physical Chemistry</i> , 2021 , 40, 51-79	7	4
143	Lanthanide(III) Complexes Based on an 18-Membered Macrocycle Containing Acetamide Pendants. Structural Characterization and paraCEST Properties. <i>Inorganic Chemistry</i> , 2021 , 60, 1902-1914	5.1	1
142	Stability, relaxometric and computational studies on Mn complexes with ligands containing a cyclobutane scaffold. <i>Dalton Transactions</i> , 2021 , 50, 1076-1085	4.3	2
141	PycLen-Based Ligands Bearing Pendant Picolinate Arms for Gadolinium Complexation. <i>Inorganic Chemistry</i> , 2021 , 60, 2390-2405	5.1	4
140	Scrutinising the role of intramolecular hydrogen bonding in water exchange dynamics of Gd(III) complexes. <i>Dalton Transactions</i> , 2021 , 50, 5506-5518	4.3	2
139	Macrocyclic PycLen-Based Gd Complex with High Relaxivity and pH Response. <i>Inorganic Chemistry</i> , 2020 , 59, 7306-7317	5.1	3
138	Axial Ligation in Ytterbium(III) DOTAM Complexes Rationalized with Multireference and Ligand-Field ab Initio Calculations. <i>Journal of Physical Chemistry A</i> , 2020 , 124, 1362-1371	2.8	5
137	pH-Dependent Hydration Change in a Gd-Based MRI Contrast Agent with a Phosphonated Ligand. <i>Chemistry - A European Journal</i> , 2020 , 26, 5407-5418	4.8	3
136	Unexpected Trends in the Stability and Dissociation Kinetics of Lanthanide(III) Complexes with Cyclen-Based Ligands across the Lanthanide Series. <i>Inorganic Chemistry</i> , 2020 , 59, 8184-8195	5.1	11
135	Inert macrocyclic Eu ³⁺ complex with affirmative paraCEST features. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 2274-2286	6.8	3

134	Hydrothermal synthesis of six new lanthanides coordination polymers based on 1-H-benzimidazole-5-carboxylic acid: Structure, Hirshfeld analysis, thermal and spectroscopic properties. <i>Inorganica Chimica Acta</i> , 2020 , 510, 119740	2.7	2
133	Ditopic binuclear copper(II) complexes for DNA cleavage. <i>Journal of Inorganic Biochemistry</i> , 2020 , 205, 110995	4.2	0
132	Combined NMR, DFT and X-ray studies highlight structural and hydration changes of [Ln(AAZTA)] ⁺ complexes across the series. <i>Inorganic Chemistry Frontiers</i> , 2020 , 7, 795-803	6.8	10
131	Mn Complexes Containing Sulfonamide Groups with pH-Responsive Relaxivity. <i>Inorganic Chemistry</i> , 2020 , 59, 14306-14317	5.1	5
130	Mn(II) compounds as an alternative to Gd-based MRI probes. <i>Future Medicinal Chemistry</i> , 2019 , 11, 1461-1483	4.8	44
129	Gadolinium Complexes of Highly Rigid, Open-Chain Ligands Containing a Cyclobutane Ring in the Backbone: Decreasing Ligand Denticity Might Enhance Kinetic Inertness. <i>Inorganic Chemistry</i> , 2019 , 58, 13170-13183	5.1	6
128	Electronic versus steric control in palladium complexes of carboranyl phosphine-iminophosphorane ligands. <i>Dalton Transactions</i> , 2019 , 48, 486-503	4.3	1
127	A pentadentate member of the picolinate family for Mn(II) complexation and an amphiphilic derivative. <i>Dalton Transactions</i> , 2019 , 48, 696-710	4.3	6
126	Methylthiazolyl Tacn Ligands for Copper Complexation and Their Bifunctional Chelating Agent Derivatives for Bioconjugation and Copper-64 Radiolabeling: An Example with Bombesin. <i>Inorganic Chemistry</i> , 2019 , 58, 2669-2685	5.1	8
125	Gadolinium(III)-Based Dual H/ F Magnetic Resonance Imaging Probes. <i>Chemistry - A European Journal</i> , 2019 , 25, 4782-4792	4.8	11
124	Accelerating water exchange in Gd-DO3A-derivatives by favouring the dissociative mechanism through hydrogen bonding. <i>Chemical Communications</i> , 2019 , 55, 513-516	5.8	13
123	Water exchange in lanthanide complexes for MRI applications. Lessons learned over the last 25 years. <i>Dalton Transactions</i> , 2019 , 48, 11161-11180	4.3	23
122	Phosphate and polyphosphate anion recognition by a dinuclear copper(II) complex of an unsymmetrical squaramide. <i>Dalton Transactions</i> , 2019 , 48, 10104-10115	4.3	6
121	PIDAZTA: Structurally Constrained Chelators for the Efficient Formation of Stable Gallium-68 Complexes at Physiological pH. <i>Chemistry - A European Journal</i> , 2019 , 25, 10698-10709	4.8	6
120	Lanthanide Complexes with H paraCEST and F Response for Magnetic Resonance Imaging Applications. <i>Inorganic Chemistry</i> , 2019 , 58, 7571-7583	5.1	14
119	endo- versus exo-Cyclic coordination in copper complexes with methylthiazolylcarboxylate tacn derivatives. <i>Dalton Transactions</i> , 2019 , 48, 8740-8755	4.3	2
118	Understanding the Optical and Magnetic Properties of Ytterbium(III) Complexes. <i>Inorganic Chemistry</i> , 2019 , 58, 3732-3743	5.1	17
117	Reinforced Ni(II)-cyclam derivatives as dual H/F MRI probes. <i>Chemical Communications</i> , 2019 , 55, 4115-4118	5.1	14

116	Controlling water exchange rates in potential Mn-based MRI agents derived from NO ₂ A. <i>Dalton Transactions</i> , 2019 , 48, 3962-3972	4.3	10
115	Highly Stable and Inert Complexation of Indium(III) by Reinforced Cyclam Dipicolinate and a Bifunctional Derivative for Bead Encoding in Mass Cytometry. <i>Chemistry - A European Journal</i> , 2019 , 25, 15387-15400	4.8	6
114	The role of ligand to metal charge-transfer states on the luminescence of Europium complexes with 18-membered macrocyclic ligands. <i>Dalton Transactions</i> , 2019 , 48, 4035-4045	4.3	20
113	Characterisation of magnetic resonance imaging (MRI) contrast agents using NMR relaxometry. <i>Molecular Physics</i> , 2019 , 117, 898-909	1.7	34
112	MetalOrganic Self-Assembled Trefoil Knots for CBr Bond Activation. <i>ACS Catalysis</i> , 2019 , 9, 1907-1914	13.1	22
111	Synthesis and Characterization of Positively Charged tris-Imidazolium Calix[6]arene Hosts for Anion Recognition. <i>ChemistrySelect</i> , 2019 , 4, 321-328	1.8	3
110	Taking the next step toward inert Mn ²⁺ complexes of open-chain ligands: the case of the rigid PhDTA ligand. <i>New Journal of Chemistry</i> , 2018 , 42, 8001-8011	3.6	23
109	Stable and Inert Yttrium(III) Complexes with Pyclen-Based Ligands Bearing Pendant Picolinate Arms: Toward New Pharmaceuticals for Radiotherapy. <i>Inorganic Chemistry</i> , 2018 , 57, 2051-2063	5.1	18
108	Coordination Properties of GdDO ₃ A-Based Model Compounds of Bioresponsive MRI Contrast Agents. <i>Inorganic Chemistry</i> , 2018 , 57, 5973-5986	5.1	16
107	Steric Effects on the Binding of Phosphate and Polyphosphate Anions by Zinc(II) and Copper(II) Dinuclear Complexes of m-Xylyl-bis-cyclen. <i>Inorganic Chemistry</i> , 2018 , 57, 6466-6478	5.1	10
106	Modulating the DNA cleavage ability of copper(II) Schiff bases through ternary complex formation. <i>New Journal of Chemistry</i> , 2018 , 42, 15170-15183	3.6	8
105	Long Wavelength Excitation of Europium Luminescence in Extended, Carboline-Based Cryptates. <i>Inorganic Chemistry</i> , 2018 , 57, 7390-7401	5.1	11
104	A Coordination Chemistry Approach to Fine-Tune the Physicochemical Parameters of Lanthanide Complexes Relevant to Medical Applications. <i>Chemistry - A European Journal</i> , 2018 , 24, 3127-3131	4.8	17
103	On the consequences of the stereochemical activity of the Bi(III) 6s lone pair in cyclen-based complexes. The [Bi(DO ₃ A)] case. <i>Dalton Transactions</i> , 2018 , 47, 13830-13842	4.3	11
102	Modeling the OEC with Two New Biomimetic Models: Preparations, Structural Characterization, and Water Photolysis Studies of a BaMn Box Type Complex and a Mn ₄ N ₆ Planar-Diamond Cluster. <i>Catalysts</i> , 2018 , 8, 382	4	1
101	Recognition of AMP, ADP and ATP through Cooperative Binding by Cu(II) and Zn(II) Complexes Containing Urea and/or Phenylboronic-Acid Moieties. <i>Molecules</i> , 2018 , 23,	4.8	13
100	Expanding the Family of Pyclen-Based Ligands Bearing Pendant Picolinate Arms for Lanthanide Complexation. <i>Inorganic Chemistry</i> , 2018 , 57, 6932-6945	5.1	25
99	Ditopic receptors containing urea groups for solvent extraction of Cu(II) salts. <i>Dalton Transactions</i> , 2017 , 46, 3192-3206	4.3	15

98	Spectroscopic Properties of a Family of Mono- to Trinuclear Lanthanide Complexes. <i>European Journal of Inorganic Chemistry</i> , 2017 , 2017, 2122-2129	2.3	7
97	1,4,7-Triazacyclononane-Based Bifunctional Picolinate Ligands for Efficient Copper Complexation. <i>European Journal of Inorganic Chemistry</i> , 2017 , 2017, 2435-2443	2.3	12
96	Spectrally Undiscerned Isomers Might Lead to Erroneous Determination of Water Exchange Rates of paraCEST Eu(III) Agents. <i>Inorganic Chemistry</i> , 2017 , 56, 7737-7745	5.1	15
95	Developing the family of picolinate ligands for Mn complexation. <i>Dalton Transactions</i> , 2017 , 46, 1546-1558	4.3	31
94	Enantiomeric Recognition of d- and l-Lactate by CEST with the Aid of a Paramagnetic Shift Reagent. <i>Journal of the American Chemical Society</i> , 2017 , 139, 17431-17437	16.4	19
93	Toward inert paramagnetic Ni(ii)-based chemical exchange saturation transfer MRI agents. <i>Dalton Transactions</i> , 2017 , 46, 15095-15106	4.3	10
92	Effects of the substituents of pyrazole/thiazine ligands on the magnetic properties of chloro-bridged Cu(II) complexes. <i>New Journal of Chemistry</i> , 2017 , 41, 8818-8827	3.6	8
91	The role of the capping bond effect on pyclen Y/Y chelates: full control of the regiospecific N-functionalization makes the difference. <i>Chemical Communications</i> , 2017 , 53, 9534-9537	5.8	18
90	Tuning the copper(ii) coordination properties of cyclam by subtle chemical modifications. <i>Dalton Transactions</i> , 2017 , 46, 11479-11490	4.3	5
89	Dimer formation of GdDO3A-arylsulfonamide complexes causes loss of pH-dependency of relaxivity. <i>Dalton Transactions</i> , 2017 , 46, 16828-16836	4.3	12
88	Definition of the Labile Capping Bond Effect in Lanthanide Complexes. <i>Chemistry - A European Journal</i> , 2017 , 23, 1110-1117	4.8	17
87	Chapter 2: Gadolinium-based Contrast Agents. <i>New Developments in NMR</i> , 2017 , 121-242	0.9	11
86	Chapter 5: Transition Metal-based T1 Contrast Agents. <i>New Developments in NMR</i> , 2017 , 448-478	0.9	2
85	Optimising the relaxivities of Mn complexes by targeting human serum albumin (HSA). <i>Dalton Transactions</i> , 2017 , 46, 8494-8504	4.3	20
84	Sulphur-rich functionalized calix[4]arenes for selective complexation of Hg over Cu, Zn and Cd. <i>Dalton Transactions</i> , 2016 , 45, 15211-15224	4.3	12
83	Complexation of Ln(3+) Ions with Cyclam Dipicolinates: A Small Bridge that Makes Huge Differences in Structure, Equilibrium, and Kinetic Properties. <i>Inorganic Chemistry</i> , 2016 , 55, 2227-39	5.1	20
82	Magnetic Anisotropies in Rhombic Lanthanide(III) Complexes Do Not Conform to Bleaney's Theory. <i>Inorganic Chemistry</i> , 2016 , 55, 3490-7	5.1	41
81	Approaching the Kinetic Inertness of Macrocyclic Gadolinium(III)-Based MRI Contrast Agents with Highly Rigid Open-Chain Derivatives. <i>Chemistry - A European Journal</i> , 2016 , 22, 896-901	4.8	27

80	The Relationship between NMR Chemical Shifts of Thermally Polarized and Hyperpolarized Y Complexes and Their Solution Structures. <i>Chemistry - A European Journal</i> , 2016 , 22, 16657-16667	4.8	11
79	Transient versus Static Electron Spin Relaxation in Mn(2+) Complexes Relevant as MRI Contrast Agents. <i>Journal of Physical Chemistry A</i> , 2016 , 120, 6467-76	2.8	14
78	Water exchange rates and mechanisms in tetrahedral [Be(H ₂ O) ₄] ²⁺ and [Li(H ₂ O) ₄] ⁺ complexes using DFT methods and cluster-continuum models. <i>International Journal of Quantum Chemistry</i> , 2016 , 116, 1388-1396	2.1	3
77	Stabilizing divalent europium in aqueous solution using size-discrimination and electrostatic effects. <i>Inorganic Chemistry</i> , 2015 , 54, 4940-52	5.1	33
76	Stable Mn(2+), Cu(2+) and Ln(3+) complexes with cyclen-based ligands functionalized with picolinate pendant arms. <i>Dalton Transactions</i> , 2015 , 44, 5017-31	4.3	49
75	Mono-, bi-, and trinuclear bis-hydrated Mn(2+) complexes as potential MRI contrast agents. <i>Inorganic Chemistry</i> , 2015 , 54, 9576-87	5.1	33
74	Gd(3+)-Based Magnetic Resonance Imaging Contrast Agent Responsive to Zn(2+). <i>Inorganic Chemistry</i> , 2015 , 54, 10342-50	5.1	26
73	Importance of outer-sphere and aggregation phenomena in the relaxation properties of phosphonated gadolinium complexes with potential applications as MRI contrast agents. <i>Chemistry - A European Journal</i> , 2015 , 21, 6535-46	4.8	20
72	Exceptionally Inert Lanthanide(III) PARACEST MRI Contrast Agents Based on an 18-Membered Macrocyclic Platform. <i>Chemistry - A European Journal</i> , 2015 , 21, 18662-70	4.8	14
71	Reasons behind the relative abundances of heptacoordinate complexes along the late first-row transition metal series. <i>Inorganic Chemistry</i> , 2014 , 53, 12859-69	5.1	27
70	¹⁷ O and ¹ H relaxometric and DFT study of hyperfine coupling constants in [Mn(H ₂ O) ₆] ²⁺ . <i>RSC Advances</i> , 2014 , 4, 7094	3.7	43
69	Cooperative anion recognition in copper(II) and zinc(II) complexes with a ditopic tripodal ligand containing a urea group. <i>Inorganic Chemistry</i> , 2014 , 53, 2554-68	5.1	27
68	Understanding stability trends along the lanthanide series. <i>Chemistry - A European Journal</i> , 2014 , 20, 3974-81	4.8	57
67	Highly Stable Complexes of Divalent Metal Ions (Mg ²⁺ , Ca ²⁺ , Cu ²⁺ , Zn ²⁺ , Cd ²⁺ , and Pb ²⁺) with a DOTA-Like Ligand Containing a Picolinate Pendant. <i>European Journal of Inorganic Chemistry</i> , 2014 , 2014, 6165-6173	2.3	10
66	Lanthanide(III) complexes with a reinforced cyclam ligand show unprecedented kinetic inertness. <i>Journal of the American Chemical Society</i> , 2014 , 136, 17954-7	16.4	44
65	High relaxivity Mn(2+)-based MRI contrast agents. <i>Chemistry - A European Journal</i> , 2014 , 20, 17300-5	4.8	24
64	Cinderella Elements: Strategies to increase the stability of group 1 complexes by tailoring crown macrocycles. <i>Inorganica Chimica Acta</i> , 2014 , 417, 155-162	2.7	3
63	Self-aggregated dinuclear lanthanide(III) complexes as potential bimodal probes for magnetic resonance and optical imaging. <i>Chemistry - A European Journal</i> , 2013 , 19, 11696-706	4.8	17

62	Hyperfine coupling constants on inner-sphere water molecules of a triazacyclononane-based Mn(II) complex and related systems relevant as MRI contrast agents. <i>Inorganic Chemistry</i> , 2013 , 52, 11173-84	5.1	39
61	Solid state and solution structures of alkaline-earth complexes with lariat ethers containing aniline and benzimidazole pendants. <i>Polyhedron</i> , 2012 , 31, 402-412	2.7	3
60	Highly relaxing gadolinium based MRI contrast agents responsive to Mg ²⁺ sensing. <i>Chemical Communications</i> , 2012 , 48, 4085-7	5.8	26
59	Lanthanide complexes based on a diazapyridinophane platform containing picolinate pendants. <i>Inorganic Chemistry</i> , 2012 , 51, 10893-903	5.1	29
58	Lanthanide(III) complexes with ligands derived from a cyclen framework containing pyridinecarboxylate pendants. The effect of steric hindrance on the hydration number. <i>Inorganic Chemistry</i> , 2012 , 51, 2509-21	5.1	58
57	Hyperfine coupling constants on inner-sphere water molecules of Gd(III)-based MRI contrast agents. <i>ChemPhysChem</i> , 2012 , 13, 3640-50	3.2	72
56	Density functional dependence of molecular geometries in lanthanide(III) complexes relevant to bioanalytical and biomedical applications. <i>Computational and Theoretical Chemistry</i> , 2012 , 999, 93-104	2	45
55	Solution structure of Ln(III) complexes with macrocyclic ligands through theoretical evaluation of ¹ H NMR contact shifts. <i>Inorganic Chemistry</i> , 2012 , 51, 13419-29	5.1	36
54	Monopicolinate cyclen and cyclam derivatives for stable copper(II) complexation. <i>Inorganic Chemistry</i> , 2012 , 51, 6916-27	5.1	67
53	Definition of an intramolecular Eu-to-Eu energy transfer within a discrete [Eu ₂ L] complex in solution. <i>Chemistry - A European Journal</i> , 2012 , 18, 8163-73	4.8	35
52	The effect of ring size variation on the structure and stability of lanthanide(III) complexes with crown ethers containing picolinate pendants. <i>Dalton Transactions</i> , 2011 , 40, 384-92	4.3	26
51	Applications of Density Functional Theory (DFT) to Investigate the Structural, Spectroscopic and Magnetic Properties of Lanthanide(III) Complexes. <i>Current Inorganic Chemistry</i> , 2011 , 1, 91-116		41
50	A merged experimental and theoretical conformational study on alkaline-earth complexes with lariat ethers derived from 4,13-diaza-18-crown-6. <i>Inorganica Chimica Acta</i> , 2011 , 370, 270-278	2.7	6
49	Macrocyclic receptor showing extremely high Sr(II)/Ca(II) and Pb(II)/Ca(II) selectivities with potential application in chelation treatment of metal intoxication. <i>Inorganic Chemistry</i> , 2011 , 50, 3772-84	5.1	52
48	Lanthanide dota-like complexes containing a picolinate pendant: structural entry for the design of Ln(III)-based luminescent probes. <i>Inorganic Chemistry</i> , 2011 , 50, 4125-41	5.1	69
47	Molecular recognition of sialic acid by lanthanide(III) complexes through cooperative two-site binding. <i>Inorganic Chemistry</i> , 2010 , 49, 4212-23	5.1	29
46	Macrocyclic Receptor Showing Improved Pb(II)/Zn(II) and Pb(II)/Ca(II) Selectivities. <i>European Journal of Inorganic Chemistry</i> , 2010 , 2010, 2495-2503	2.3	15
45	Structure and Dynamics of Lanthanide(III) Complexes with an N-Alkylated do3a Ligand (H ₃ do3a = 1,4,7,10-Tetraazacyclododecane-1,4,7-triacetic Acid): A Combined Experimental and DFT Study. <i>European Journal of Inorganic Chemistry</i> , 2010 , 2010, 3586-3595	2.3	28

44	Lead(II) Complexes of Lateral Macrobicyclic Receptors That Incorporate a Crown Moiety and a Pyridine Head Unit. <i>European Journal of Inorganic Chemistry</i> , 2010 , 2010, 5027-5034	2.3	10
43	Towards Selective Recognition of Sialic Acid Through Simultaneous Binding to Its cis-Diol and Carboxylate Functions. <i>European Journal of Organic Chemistry</i> , 2010 , 2010, 3237-3248	3.2	24
42	Binuclear Co(II), Ni(II), Cu(II) and Zn(II) complexes with Schiff-bases derived from crown ether platforms: Rare examples of ether oxygen atoms bridging metal centers. <i>Polyhedron</i> , 2010 , 29, 2269-2277	2.7	6
41	Anion Coordination Effect on the Nuclearity of Co(II), Ni(II), Cu(II), and Zn(II) Complexes with a Benzimidazole Pendant-Armed Crown. <i>European Journal of Inorganic Chemistry</i> , 2009 , 2009, 400-411	2.3	11
40	Macrocyclic receptor exhibiting unprecedented selectivity for light lanthanides. <i>Journal of the American Chemical Society</i> , 2009 , 131, 3331-41	16.4	100
39	Selective chelation of Cd(II) and Pb(II) versus Ca(II) and Zn(II) by using octadentate ligands containing pyridinecarboxylate and pyridyl pendants. <i>Inorganic Chemistry</i> , 2009 , 48, 10976-87	5.1	20
38	Eight-coordinate Zn(II), Cd(II), and Pb(II) complexes based on a 1,7-diaza-12-crown-4 platform endowed with a remarkable selectivity over Ca(II). <i>Inorganic Chemistry</i> , 2009 , 48, 11821-31	5.1	33
37	Stability, water exchange, and anion binding studies on lanthanide(III) complexes with a macrocyclic ligand based on 1,7-diaza-12-crown-4: extremely fast water exchange on the Gd ³⁺ complex. <i>Inorganic Chemistry</i> , 2009 , 48, 8878-89	5.1	51
36	Lanthanide complexes based on a 1,7-diaza-12-crown-4 platform containing picolinate pendants: a new structural entry for the design of magnetic resonance imaging contrast agents. <i>Inorganic Chemistry</i> , 2008 , 47, 7840-51	5.1	76
35	Zn(II), Cd(II) and Pb(II) complexation with pyridinecarboxylate containing ligands. <i>Dalton Transactions</i> , 2008 , 5754-65	4.3	56
34	Protonated macrobicyclic hosts containing pyridine head units for anion recognition. <i>Chemistry - A European Journal</i> , 2008 , 14, 5829-38	4.8	20
33	Conformational study of lanthanide(III) complexes of N-(2-salicylaldehydato)benzyl-1-aza-18-crown-6 by using X-ray and ab initio methods. <i>Polyhedron</i> , 2008 , 27, 1415-1422	2.7	5
32	Seven-coordination versus six-coordination in divalent first-row transition-metal complexes derived from 1,10-diaza-15-crown-5. <i>Inorganic Chemistry</i> , 2007 , 46, 8271-82	5.1	38
31	Effect of Protonation and Interaction with Anions on a Lead(II) Complex with a Lateral Macrobicycle Containing a Phenol Schiff-Base Spacer. <i>European Journal of Inorganic Chemistry</i> , 2007 , 2007, 1635-1643	2.3	15
30	Receptor versus Counterion: Capability of N,N'-Bis(2-aminobenzyl)-diazacrowns for Giving Endo- and/or Exocyclic Coordination of Zn(II). <i>European Journal of Inorganic Chemistry</i> , 2007 , 2007, 1874-1883	2.3	11
29	Metal Ion Complementarity: Effect of Ring-Size Variation on the Conformation and Stability of Lead(II) and Cadmium(II) Complexes with Pendant-Armed Crowns. <i>European Journal of Inorganic Chemistry</i> , 2007 , 2007, 2198-2207	2.3	21
28	Synthesis and crystal structure of manganese(II) complexes with high-denticity ligands derived from azacrowns. <i>Polyhedron</i> , 2007 , 26, 4141-4146	2.7	6
27	What anions do to N-H-containing receptors. <i>Accounts of Chemical Research</i> , 2006 , 39, 343-53	24.3	718

26	Lone-pair activity in lead(II) complexes with unsymmetrical lariat ethers. <i>Inorganic Chemistry</i> , 2006 , 45, 5407-16	5.1	48
25	{4,10-Bis[2-(2-oxidobenzylideneamino-kappa2N,O)benzyl]-1,7-dioxa-4,10-diazacyclododecane-kappa4O1,N4,O3,N10]yt perchlorate acetonitrile solvate. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2006 , 62, m360-2		1
24	Some guidelines for the design of anion receptors. <i>Coordination Chemistry Reviews</i> , 2006 , 250, 1451-1470	3.2	222
23	Urea vs. thiourea in anion recognition. <i>Organic and Biomolecular Chemistry</i> , 2005 , 3, 1495-500	3.9	321
22	Designing binuclear transition metal complexes: a new example of the versatility of N,N'-bis(2-aminobenzyl)-4,13-diaza-18-crown-6. <i>Dalton Transactions</i> , 2005 , 2031-7	4.3	13
21	Electronic structure study of seven-coordinate first-row transition metal complexes derived from 1,10-diaza-15-crown-5: a successful marriage of theory with experiment. <i>Inorganic Chemistry</i> , 2005 , 44, 9704-13	5.1	50
20	Lateral macrobicyclic architectures: toward new lead(II) sequestering agents. <i>Inorganic Chemistry</i> , 2005 , 44, 5428-36	5.1	20
19	Chiral receptors for phosphate ions. <i>Organic and Biomolecular Chemistry</i> , 2005 , 3, 2632-9	3.9	90
18	Lead(II) thiocyanate complexes with bibracchial lariat ethers: an X-ray and DFT study. <i>Inorganic Chemistry</i> , 2005 , 44, 2224-33	5.1	62
17	Why, on interaction of urea-based receptors with fluoride, beautiful colors develop. <i>Journal of Organic Chemistry</i> , 2005 , 70, 5717-20	4.2	456
16	Metal-enhanced H-bond donor tendencies of urea and thiourea toward anions: ditopic receptors for silver(I) salts. <i>Inorganic Chemistry</i> , 2005 , 44, 8690-8	5.1	62
15	Barium(II) thiocyanate templating Schiff-base lateral macrobicycles derived from 1,10-diaza-15-crown-5. <i>Polyhedron</i> , 2005 , 24, 289-294	2.7	8
14	A two-channel chemosensor for the optical detection of carboxylic acids, including cholic acid. <i>Journal of Materials Chemistry</i> , 2005 , 15, 2670		47
13	Anion-induced urea deprotonation. <i>Chemistry - A European Journal</i> , 2005 , 11, 3097-104	4.8	240
12	A Schiff base lateral macrobicycle derived from 4,13-diaza-18-crown-6 in its protonated form. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2005 , 61, o92-4		4
11	Nature of urea-fluoride interaction: incipient and definitive proton transfer. <i>Journal of the American Chemical Society</i> , 2004 , 126, 16507-14	16.4	758
10	Synthesis and structural characterisation of lead(II) isothiocyanate complexes with receptors derived from 1,10-diaza-15-crown-5. <i>Polyhedron</i> , 2003 , 22, 2709-2717	2.7	11
9	[7,13-Bis(2-aminobenzyl)-1,4,10-trioxa-7,13-diazacyclopentadecane]diisothiocyanatobarium(II). <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003 , 59, m16-7		1

8	A barium perchlorate complex with a lateral macrobicyclic derived from 1,10-diaza-15-crown-5 containing a phenol Schiff base spacer. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003 , 59, m93-4		2
7	A barium perchlorate complex with a lateral macrobicyclic derived from 4,13-diaza-18-crown-6 containing a pyridine Schiff base spacer. <i>Acta Crystallographica Section C: Crystal Structure Communications</i> , 2003 , 59, m450-1		1
6	Templating Schiff-base lateral macrobicyclics: an experimental and theoretical structural study of the intermediates. <i>Inorganic Chemistry</i> , 2003 , 42, 4299-307	5.1	30
5	Lead(II) complexes with macrocyclic receptors derived from 4,13-diaza-18-crown-6. <i>Inorganic Chemistry</i> , 2002 , 41, 4337-47	5.1	44
4	Copper complexes with bibracchial lariat ethers: from mono- to binuclear structures. <i>Inorganica Chimica Acta</i> , 2001 , 317, 190-198	2.7	30
3	Cadmium(II) and Lead(II) Complexes with Novel Macrocyclic Receptors Derived from 1,10-Diaza-15-crown-5. <i>European Journal of Inorganic Chemistry</i> , 2000 , 2000, 1445-1456	2.3	35
2	Barium Templating Schiff-Base Lateral Macrobicyclics. <i>Inorganic Chemistry</i> , 1999 , 38, 1937-1944	5.1	27
1	The critical role of ligand topology: strikingly different properties of Gd(III) complexes with regioisomeric AAZTA derivatives. <i>Inorganic Chemistry Frontiers</i> ,	6.8	1