

# Krzysztof Buczkowski

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6080465/publications.pdf>

Version: 2024-02-01

20  
papers

5,995  
citations

932766

10  
h-index

839053

18  
g-index

21  
all docs

21  
docs citations

21  
times ranked

8091  
citing authors

#	ARTICLE	IF	CITATIONS
1	Gender differences in self-care for common colds by primary care patients: a European multicenter survey on the prevalence and patterns of practices (the COCO study). <i>Journal of Gender Studies</i> , 2021, 30, 756-771.	1.3	5
2	Reasons for and Scenarios Associated with Failure to Cease Smoking: Results from a Qualitative Study Among Polish Smokers Who Had Unsuccessfully Attempted to Quit. <i>Patient Preference and Adherence</i> , 2021, Volume 15, 2071-2084.	0.8	9
3	Primary care practitioners' diagnostic action when the patient may have cancer: an exploratory vignette study in 20 European countries. <i>BMJ Open</i> , 2020, 10, e035678.	0.8	9
4	Znieczulenie podpajczynkowe do laparoskopowej cholecystektomii u pacjentki z zespoÅem Marfana. <i>Forum Leczenia Ran</i> , 2020, 1, 167-170.	0.0	0
5	How European primary care practitioners think the timeliness of cancer diagnosis can be improved: a thematic analysis. <i>BMJ Open</i> , 2019, 9, e030169.	0.8	15
6	Adipose tissue – the structure and its functions, with particular emphasis on the characteristics of selected adipokines and their effects on the organism. <i>Medycyna Ogólna i Nauki o Zdrowiu</i> , 2019, 25, 162-169.	0.1	0
7	Identifying important health system factors that influence primary care practitioners' referrals for cancer suspicion: a European cross-sectional survey. <i>BMJ Open</i> , 2018, 8, e022904.	0.8	17
8	Quality control of oral anticoagulation with vitamin K antagonists in primary care patients in Poland: a multi-centre study. <i>Kardiologia Polska</i> , 2018, 76, 764-769.	0.3	7
9	Smoking Status and the Five-Factor Model of Personality: Results of a Cross-Sectional Study Conducted in Poland. <i>International Journal of Environmental Research and Public Health</i> , 2017, 14, 126.	1.2	14
10	Self-Care for Common Colds by Primary Care Patients: A European Multicenter Survey on the Prevalence and Patterns of Practices – The COCO Study. <i>Evidence-based Complementary and Alternative Medicine</i> , 2016, 2016, 1-9.	0.5	10
11	Self-Care Practices for Common Colds by Primary Care Patients: Study Protocol of a European Multicenter Survey – The COCO Study. <i>Evidence-based Complementary and Alternative Medicine</i> , 2015, 2015, 1-6.	0.5	4
12	Association between Genetic Variants on Chromosome 15q25 Locus and Several Nicotine Dependence Traits in Polish Population: A Case-Control Study. <i>BioMed Research International</i> , 2015, 2015, 1-8.	0.9	13
13	Motivations toward smoking cessation, reasons for relapse, and modes of quitting: results from a qualitative study among former and current smokers. <i>Patient Preference and Adherence</i> , 2014, 8, 1353.	0.8	69
14	The attractiveness of family medicine among Polish medical students. <i>European Journal of General Practice</i> , 2014, 20, 121-124.	0.9	5
15	“What kind of general practitioner do I need for smoking cessation?” Results from a qualitative study in Poland. <i>BMC Family Practice</i> , 2013, 14, 159.	2.9	12
16	Influences of polymorphic variants of DRD2 and SLC6A3 genes, and their combinations on smoking in Polish population. <i>BMC Medical Genetics</i> , 2009, 10, 92.	2.1	21
17	Polish medical students' perceptions of the nursing profession: a cross-sectional study. <i>Scandinavian Journal of Caring Sciences</i> , 2009, 23, 438-445.	1.0	7
18	Lack of association between serotonin transporter gene polymorphism 5-HTTLPR and smoking among Polish population: a case-control study. <i>BMC Medical Genetics</i> , 2008, 9, 76.	2.1	18

#	ARTICLE	IF	CITATIONS
19	Patterns of motivations and ways of quitting smoking among Polish smokers: A questionnaire study. BMC Public Health, 2008, 8, 274.	1.2	48
20	Rosuvastatin to Prevent Vascular Events in Men and Women with Elevated C-Reactive Protein. New England Journal of Medicine, 2008, 359, 2195-2207.	13.9	5,712