

# Paul Ziegler

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6082204/publications.pdf>

Version: 2024-02-01

8  
papers

2,613  
citations

1163117

8  
h-index

1588992

8  
g-index

9  
all docs

9  
docs citations

9  
times ranked

5896  
citing authors

#	ARTICLE	IF	CITATIONS
1	Loss of Stat6 affects chromatin condensation in intestinal epithelial cells causing diverse outcome in murine models of inflammation-associated and sporadic colon carcinogenesis. <i>Oncogene</i> , 2019, 38, 1787-1801.	5.9	13
2	IKK $\hat{2}$ acts as a tumor suppressor in cancer-associated fibroblasts during intestinal tumorigenesis. <i>Journal of Experimental Medicine</i> , 2015, 212, 2253-2266.	8.5	116
3	High-fat-diet-mediated dysbiosis promotes intestinal carcinogenesis independently of obesity. <i>Nature</i> , 2014, 514, 508-512.	27.8	366
4	IL-6R/STAT3/miR-34a feedback loop promotes EMT-mediated colorectal cancer invasion and metastasis. <i>Journal of Clinical Investigation</i> , 2014, 124, 1853-1867.	8.2	613
5	Interleukin-11 Is the Dominant IL-6 Family Cytokine during Gastrointestinal Tumorigenesis and Can Be Targeted Therapeutically. <i>Cancer Cell</i> , 2013, 24, 257-271.	16.8	341
6	Intestinal Tumorigenesis Initiated by Dedifferentiation and Acquisition of Stem-Cell-like Properties. <i>Cell</i> , 2013, 152, 25-38.	28.9	889
7	Loss of p53 in Enterocytes Generates an Inflammatory Microenvironment Enabling Invasion and Lymph Node Metastasis of Carcinogen-Induced Colorectal Tumors. <i>Cancer Cell</i> , 2013, 23, 93-106.	16.8	241
8	TNF- $\hat{1}$ -dependent loss of IKK $\hat{2}$ -deficient myeloid progenitors triggers a cytokine loop culminating in granulocytosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 6567-6572.	7.1	34