

Eveline Gart

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6086687/publications.pdf>

Version: 2024-02-01

12
papers

1,125
citations

1039406

9
h-index

1281420

11
g-index

12
all docs

12
docs citations

12
times ranked

2970
citing authors

#	ARTICLE	IF	CITATIONS
1	Diet-Independent Correlations between Bacteria and Dysfunction of Gut, Adipose Tissue, and Liver: A Comprehensive Microbiota Analysis in Feces and Mucosa of the Ileum and Colon in Obese Mice with NAFLD. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1.	1.8	929
2	Combined Treatment with L-Carnitine and Nicotinamide Riboside Improves Hepatic Metabolism and Attenuates Obesity and Liver Steatosis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4359.	1.8	31
3	Propionic acid and not caproic acid, attenuates nonalcoholic steatohepatitis and improves (cerebro) vascular functions in obese Ldlr ^{-/-} .Leiden mice. <i>FASEB Journal</i> , 2020, 34, 9575-9593.	0.2	29
4	Atorvastatin accelerates clearance of lipoprotein remnants generated by activated brown fat to further reduce hypercholesterolemia and atherosclerosis. <i>Atherosclerosis</i> , 2017, 267, 116-126.	0.4	23
5	Butyrate Protects against Diet-Induced NASH and Liver Fibrosis and Suppresses Specific Non-Canonical TGF- β 2 Signaling Pathways in Human Hepatic Stellate Cells. <i>Biomedicines</i> , 2021, 9, 1954.	1.4	23
6	Sex-Specific Differences in Fat Storage, Development of Non-Alcoholic Fatty Liver Disease and Brain Structure in Juvenile HFD-Induced Obese Ldlr ^{-/-} .Leiden Mice. <i>Nutrients</i> , 2019, 11, 1861.	1.7	21
7	Cholesterol Accumulation as a Driver of Hepatic Inflammation Under Translational Dietary Conditions Can Be Attenuated by a Multicomponent Medicine. <i>Frontiers in Endocrinology</i> , 2021, 12, 601160.	1.5	16
8	Krill Oil Treatment Increases Distinct PUFAs and Oxylipins in Adipose Tissue and Liver and Attenuates Obesity-Associated Inflammation via Direct and Indirect Mechanisms. <i>Nutrients</i> , 2021, 13, 2836.	1.7	16
9	Intervention with isoleucine or valine corrects hyperinsulinemia and reduces intrahepatic diacylglycerols, liver steatosis, and inflammation in Ldlr ^{-/-} .Leiden mice with manifest obesity-associated NASH. <i>FASEB Journal</i> , 2022, 36, .	0.2	16
10	Heat-Inactivated Akkermansia muciniphila Improves Gut Permeability but Does Not Prevent Development of Non-Alcoholic Steatohepatitis in Diet-Induced Obese Ldlr ^{-/-} .Leiden Mice. <i>International Journal of Molecular Sciences</i> , 2022, 23, 2325.	1.8	10
11	The Human Milk Oligosaccharide 2-Fucosyllactose Alleviates Liver Steatosis, ER Stress and Insulin Resistance by Reducing Hepatic Diacylglycerols and Improved Gut Permeability in Obese Ldlr ^{-/-} .Leiden Mice. <i>Frontiers in Nutrition</i> , 0, 9, .	1.6	10
12	Propionic acid intervention in obese Ldlr ^{-/-} .Leiden mice attenuates NASH development, but negatively affects cognition. <i>Proceedings of the Nutrition Society</i> , 2020, 79, .	0.4	1