

Ana Charrua

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6088887/ana-charrua-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

38

papers

898

citations

16

h-index

29

g-index

70

ext. papers

1,023

ext. citations

2.5

avg, IF

3.72

L-index

#	Paper	IF	Citations
38	DU Is Induced by Low Levels of Urinary ATP in a Rat Model of Partial Bladder Outlet Obstruction: The Incidence of Both Events Decreases after Deobstruction.. <i>Advances in Urology</i> , 2022 , 2022, 6292457	1.6	0
37	The Role of Urinary VEGF in Observational Studies of BPS/IC Patients: A Systematic Review. <i>Diagnostics</i> , 2022 , 12, 1037	3.8	0
36	Preclinical models of endometriosis and interstitial cystitis/bladder pain syndrome: an Innovative Medicines Initiative-PainCare initiative to improve their value for translational research in pelvic pain. <i>Pain</i> , 2021 , 162, 2349-2365	8	2
35	Biomarkers for Bladder Pain Syndrome/Interstitial Cystitis. <i>Current Bladder Dysfunction Reports</i> , 2021 , 16, 12-18	0.4	1
34	Understanding underactive bladder: a review of the contemporary literature. <i>Porto Biomedical Journal</i> , 2020 , 5, e070	1.1	7
33	Fatty acid amide hydrolase inhibition normalises bladder function and reduces pain through normalising the anandamide/palmitoylethanolamine ratio in the inflamed bladder of rats. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2020 , 393, 263-272	3.4	3
32	Effect of Water Avoidance Stress on serum and urinary NGF levels in rats: diagnostic and therapeutic implications for BPS/IC patients. <i>Scientific Reports</i> , 2019 , 9, 14113	4.9	4
31	TRPV1 Antagonists as Novel Anti-Diabetic Agents: Regulation of Oral Glucose Tolerance and Insulin Secretion Through Reduction of Low-Grade Inflammation?. <i>Medical Sciences (Basel, Switzerland)</i> , 2019 , 7,	3.3	6
30	Underactive bladder in aging rats is associated with a reduced number of serotonin-expressing cells in the urethra and is improved by serotonin application to the urethra. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, 248-254	1.9	6
29	The Impact of Chronic Pelvic Ischemia on LUTS and Urinary Levels of Neuroinflammatory, Inflammatory, and Oxidative Stress Markers in Elderly Men: A Case-control Study. <i>Urology</i> , 2019 , 123, 230-234	1.6	7
28	Intratriganal OnabotulinumtoxinA Improves Bladder Symptoms and Quality of Life in Patients with Bladder Pain Syndrome/Interstitial Cystitis: A Pilot, Single Center, Randomized, Double-Blind, Placebo Controlled Trial. <i>Journal of Urology</i> , 2018 , 199, 998-1003	2.5	28
27	Expression of cleaved SNAP-25 after bladder wall injection of onabotulinumtoxinA or abobotulinumtoxinA: A comparative study in the mice. <i>Neurourology and Urodynamics</i> , 2017 , 36, 86-90	2.3	6
26	The water avoidance stress induces bladder pain due to a prolonged alpha1A adrenoceptor stimulation. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2017 , 390, 839-844	3.4	16
25	MP29-03 CHILDHOOD STRESSFUL EVENTS INDUCE CHRONIC BLADDER PAIN IN ADULTHOOD THROUGH A TRPV1 DEPENDENT MECHANISM. <i>Journal of Urology</i> , 2017 , 197,	2.5	1
24	Endovanilloid control of pain modulation by the rostroventromedial medulla in an animal model of diabetic neuropathy. <i>Neuropharmacology</i> , 2016 , 107, 49-57	5.5	19
23	Bladder pain induced by prolonged peripheral alpha 1A adrenoceptor stimulation involves the enhancement of transient receptor potential vanilloid 1 activity and an increase of urothelial adenosine triphosphate release. <i>Acta Physiologica</i> , 2016 , 218, 265-275	5.6	7
22	Intravesical Capsaicin and Resiniferatoxin for Bladder Disorders 2015 , 119-127		

21	Can the adrenergic system be implicated in the pathophysiology of bladder pain syndrome/interstitial cystitis? A clinical and experimental study. <i>Neurourology and Urodynamics</i> , 2015 , 34, 489-96	2.3	21
20	Co-administration of transient receptor potential vanilloid 4 (TRPV4) and TRPV1 antagonists potentiate the effect of each drug in a rat model of cystitis. <i>BJU International</i> , 2015 , 115, 452-60	5.6	16
19	Sympathetic nervous system and chronic bladder pain: a new tune for an old song. <i>Translational Andrology and Urology</i> , 2015 , 4, 534-42	2.3	7
18	Brain-derived neurotrophic factor, acting at the spinal cord level, participates in bladder hyperactivity and referred pain during chronic bladder inflammation. <i>Neuroscience</i> , 2013 , 234, 88-102	3.9	20
17	Transient receptor potential channels in bladder function. <i>Acta Physiologica</i> , 2013 , 207, 110-22	5.6	34
16	Transient receptor potential vanilloid 1 mediates nerve growth factor-induced bladder hyperactivity and noxious input. <i>BJU International</i> , 2012 , 110, E422-8	5.6	20
15	Rat detrusor overactivity induced by chronic spinalization can be abolished by a transient receptor potential vanilloid 1 (TRPV1) antagonist. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2012 , 166, 35-8	2.4	26
14	TRP Channels in the Genitourinary Tract. <i>Methods in Pharmacology and Toxicology</i> , 2012 , 373-395	1.1	
13	365 TRPV1 and TRPV4 antagonists have synergistic effect for treating bladder overactivity in rats. <i>European Urology Supplements</i> , 2012 , 11, e365	0.9	4
12	Expression of apoptosis-regulating genes in the rat prostate following botulinum toxin type A injection. <i>BMC Urology</i> , 2012 , 12, 1	2.2	11
11	Modulation of urinary bladder innervation: TRPV1 and botulinum toxin A. <i>Handbook of Experimental Pharmacology</i> , 2011 , 345-74	3.2	9
10	GRC-6211, a new oral specific TRPV1 antagonist, decreases bladder overactivity and noxious bladder input in cystitis animal models. <i>Journal of Urology</i> , 2009 , 181, 379-86	2.5	78
9	Functional transient receptor potential vanilloid 1 is expressed in human urothelial cells. <i>Journal of Urology</i> , 2009 , 182, 2944-50	2.5	56
8	Intrathecal delivery of resiniferatoxin (RTX) reduces detrusor overactivity and spinal expression of TRPV1 in spinal cord injured animals. <i>Experimental Neurology</i> , 2008 , 214, 301-8	5.7	26
7	Cystitis is associated with TRPV1b-downregulation in rat dorsal root ganglia. <i>NeuroReport</i> , 2008 , 19, 1469-72	1.7	16
6	Transient receptor potential vanilloid subfamily 1 is essential for the generation of noxious bladder input and bladder overactivity in cystitis. <i>Journal of Urology</i> , 2007 , 177, 1537-41	2.5	95
5	Neurochemical characterization of insulin receptor-expressing primary sensory neurons in wild-type and vanilloid type 1 transient receptor potential receptor knockout mice. <i>Journal of Comparative Neurology</i> , 2007 , 503, 334-47	3.4	32
4	The distribution of sensory fibers immunoreactive for the TRPV1 (capsaicin) receptor in the human prostate. <i>European Urology</i> , 2005 , 48, 162-7	10.2	43

3	Anandamide-evoked activation of vanilloid receptor 1 contributes to the development of bladder hyperreflexia and nociceptive transmission to spinal dorsal horn neurons in cystitis. <i>Journal of Neuroscience</i> , 2004 , 24, 11253-63	6.6	164
2	Intravesical resiniferatoxin decreases spinal c-fos expression and increases bladder volume to reflex micturition in rats with chronic inflamed urinary bladders. <i>BJU International</i> , 2004 , 94, 153-7	5.6	54
1	Insulin induces cobalt uptake in a subpopulation of rat cultured primary sensory neurons. <i>European Journal of Neuroscience</i> , 2003 , 18, 2477-86	3.5	40