

Yasuyoshi Miyata

List of Publications by Citations

Source: <https://exaly.com/author-pdf/609044/yasuyoshi-miyata-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

123 papers	1,854 citations	25 h-index	37 g-index
139 ext. papers	2,164 ext. citations	3.8 avg, IF	4.83 L-index

#	Paper	IF	Citations
123	Lymphangiogenesis and angiogenesis in bladder cancer: prognostic implications and regulation by vascular endothelial growth factors-A, -C, and -D. <i>Clinical Cancer Research</i> , 2006 , 12, 800-6	12.9	98
122	Expression of cyclooxygenase-2 in renal cell carcinoma: correlation with tumor cell proliferation, apoptosis, angiogenesis, expression of matrix metalloproteinase-2, and survival. <i>Clinical Cancer Research</i> , 2003 , 9, 1741-9	12.9	92
121	Presence of phosphorylated hepatocyte growth factor receptor/c-Met is associated with tumor progression and survival in patients with conventional renal cell carcinoma. <i>Clinical Cancer Research</i> , 2006 , 12, 4876-81	12.9	66
120	Expression of matrix metalloproteinase-7 on cancer cells and tissue endothelial cells in renal cell carcinoma: prognostic implications and clinical significance for invasion and metastasis. <i>Clinical Cancer Research</i> , 2006 , 12, 6998-7003	12.9	64
119	Pathological significance and prognostic role of microvessel density, evaluated using CD31, CD34, and CD105 in prostate cancer patients after radical prostatectomy with neoadjuvant therapy. <i>Prostate</i> , 2015 , 75, 84-91	4.2	57
118	Expression of urokinase-type plasminogen activator, urokinase-type plasminogen activator receptor and plasminogen activator inhibitors in patients with renal cell carcinoma: correlation with tumor associated macrophage and prognosis. <i>Journal of Urology</i> , 2005 , 174, 461-5	2.5	56
117	Anti-Cancer Effects of Green Tea Polyphenols Against Prostate Cancer. <i>Molecules</i> , 2019 , 24,	4.8	56
116	Phosphorylated hepatocyte growth factor receptor/c-Met is associated with tumor growth and prognosis in patients with bladder cancer: correlation with matrix metalloproteinase-2 and -7 and E-cadherin. <i>Human Pathology</i> , 2009 , 40, 496-504	3.7	54
115	High expression of HuR in cytoplasm, but not nuclei, is associated with malignant aggressiveness and prognosis in bladder cancer. <i>PLoS ONE</i> , 2013 , 8, e59095	3.7	44
114	A Mini-Review of Reactive Oxygen Species in Urological Cancer: Correlation with NADPH Oxidases, Angiogenesis, and Apoptosis. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	37
113	Expression of metalloproteinase-2, metalloproteinase-9, and tissue inhibitor of metalloproteinase-1 in transitional cell carcinoma of upper urinary tract: correlation with tumor stage and survival. <i>Urology</i> , 2004 , 63, 602-8	1.6	37
112	Thrombospondin-1 in urological cancer: pathological role, clinical significance, and therapeutic prospects. <i>International Journal of Molecular Sciences</i> , 2013 , 14, 12249-72	6.3	36
111	Tumor lymphangiogenesis in transitional cell carcinoma of the upper urinary tract: association with clinicopathological features and prognosis. <i>Journal of Urology</i> , 2006 , 176, 348-53	2.5	35
110	Anticancer Effects of Green Tea and the Underlying Molecular Mechanisms in Bladder Cancer. <i>Medicines (Basel, Switzerland)</i> , 2018 , 5,	4.1	34
109	Heme oxygenase-1 expression is associated with tumor aggressiveness and outcomes in patients with bladder cancer: a correlation with smoking intensity. <i>Translational Research</i> , 2014 , 164, 468-76	11	34
108	Expression of cyclooxygenase-2 and EP4 receptor in transitional cell carcinoma of the upper urinary tract. <i>Journal of Urology</i> , 2005 , 173, 56-60	2.5	34
107	Reconsideration of the clinical and histopathological significance of angiogenesis in prostate cancer: Usefulness and limitations of microvessel density measurement. <i>International Journal of Urology</i> , 2015 , 22, 806-15	2.3	33

106	Serum insulin-like growth factor binding protein-3/prostate-specific antigen ratio is a useful predictive marker in patients with advanced prostate cancer. <i>Prostate</i> , 2003 , 54, 125-32	4.2	33
105	CD105 is a more appropriate marker for evaluating angiogenesis in urothelial cancer of the upper urinary tract than CD31 or CD34. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2013 , 463, 673-9	5.1	31
104	Expression of matrix metalloproteinase-10 in renal cell carcinoma and its prognostic role. <i>European Urology</i> , 2007 , 52, 791-7	10.2	31
103	Green tea polyphenol suppresses tumor invasion and angiogenesis in N-butyl-(4-hydroxybutyl) nitrosamine-induced bladder cancer. <i>Cancer Epidemiology</i> , 2010 , 34, 350-4	2.8	29
102	Expression of nm23-H1 gene product in sarcomatous cancer cells of renal cell carcinoma: correlation with tumor stage and expression of matrix metalloproteinase-2, matrix metalloproteinase-9, sialyl Lewis X, and c-erbB-2. <i>Urology</i> , 2005 , 65, 1029-34	1.6	29
101	Immune complexome analysis reveals the specific and frequent presence of immune complex antigens in lung cancer patients: A pilot study. <i>International Journal of Cancer</i> , 2017 , 140, 370-380	7.5	26
100	Human antigen R is positively associated with malignant aggressiveness via upregulation of cell proliferation, migration, and vascular endothelial growth factors and cyclooxygenase-2 in prostate cancer. <i>Translational Research</i> , 2016 , 175, 116-28	11	25
99	Anti-Cancer and Protective Effects of Royal Jelly for Therapy-Induced Toxicities in Malignancies. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	25
98	Daily salt intake is an independent risk factor for pollakiuria and nocturia. <i>International Journal of Urology</i> , 2017 , 24, 384-389	2.3	23
97	Pathological Characteristics of Periodontal Disease in Patients with Chronic Kidney Disease and Kidney Transplantation. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	23
96	Lymphangiogenesis and angiogenesis in conventional renal cell carcinoma: association with vascular endothelial growth factors A to D immunohistochemistry. <i>Urology</i> , 2008 , 71, 749-54	1.6	23
95	Pheochromocytoma multisystem crisis treated with emergency surgery: a case report and literature review. <i>BMC Research Notes</i> , 2015 , 8, 758	2.3	22
94	Relationship between prostaglandin E2 receptors and clinicopathologic features in human prostate cancer tissue. <i>Urology</i> , 2006 , 68, 1360-5	1.6	21
93	Green Tea Polyphenol Induces Changes in Cancer-Related Factors in an Animal Model of Bladder Cancer. <i>PLoS ONE</i> , 2017 , 12, e0171091	3.7	20
92	High expression of Twist is associated with tumor aggressiveness and poor prognosis in patients with renal cell carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2014 , 7, 3158-65	1.4	20
91	Expression of thrombospondin-derived 4N1K peptide-containing proteins in renal cell carcinoma tissues is associated with a decrease in tumor growth and angiogenesis. <i>Clinical Cancer Research</i> , 2003 , 9, 1734-40	12.9	20
90	Erectile dysfunction in hemodialysis patients with diabetes mellitus: association with age and hemoglobin A1c levels. <i>International Journal of Urology</i> , 2004 , 11, 530-4	2.3	19
89	Pigment epithelium-derived factor inhibits fibroblast-growth-factor-2-induced capillary morphogenesis of endothelial cells through Fyn. <i>Journal of Cell Science</i> , 2005 , 118, 961-70	5.3	19

88	Pathological significance and prognostic implications of heme oxygenase 1 expression in non-muscle-invasive bladder cancer: Correlation with cell proliferation, angiogenesis, lymphangiogenesis and expression of VEGFs and COX-2. <i>Oncology Letters</i> , 2017 , 13, 275-280	2.6	18
87	Molecular Mechanisms of the Anti-Cancer Effects of Isothiocyanates from Cruciferous Vegetables in Bladder Cancer. <i>Molecules</i> , 2020 , 25,	4.8	18
86	The efficacy of mirabegron additional therapy for lower urinary tract symptoms after treatment with α -adrenergic receptor blocker monotherapy: prospective analysis of elderly men. <i>BMC Urology</i> , 2016 , 16, 45	2.2	18
85	Feline sarcoma-related protein expression correlates with malignant aggressiveness and poor prognosis in renal cell carcinoma. <i>Cancer Science</i> , 2013 , 104, 681-6	6.9	18
84	Tumor-associated stromal cells expressing E-prostanoid 2 or 3 receptors in prostate cancer: correlation with tumor aggressiveness and outcome by angiogenesis and lymphangiogenesis. <i>Urology</i> , 2013 , 81, 136-42	1.6	17
83	Met in urological cancers. <i>Cancers</i> , 2014 , 6, 2387-403	6.6	17
82	Effect of salt intake reduction on nocturia in patients with excessive salt intake. <i>Neurourology and Urodynamics</i> , 2019 , 38, 927-933	2.3	16
81	A Review of Oxidative Stress and Urinary Dysfunction Caused by Bladder Outlet Obstruction and Treatments Using Antioxidants. <i>Antioxidants</i> , 2019 , 8,	7.1	16
80	Pathological significance and predictive value for biochemical recurrence of c-Fes expression in prostate cancer. <i>Prostate</i> , 2012 , 72, 201-8	4.2	13
79	Overexpression of hepatocyte growth factor receptor in renal carcinoma cells indirectly stimulates tumor growth in vivo. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 302, 892-7	3.4	12
78	The role of c-Fes in vascular endothelial growth factor-A-mediated signaling by endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 306, 1056-63	3.4	12
77	Expression of insulin-like growth factor binding protein-3 before and after neoadjuvant hormonal therapy in human prostate cancer tissues: correlation with histopathologic effects and biochemical recurrence. <i>Urology</i> , 2004 , 63, 1184-90	1.6	12
76	Periodontal Disease in Patients Receiving Dialysis. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	11
75	A pilot study to assess the safety and usefulness of combined transurethral endoscopic mucosal resection and en-bloc resection for non-muscle invasive bladder cancer. <i>BMC Urology</i> , 2019 , 19, 56	2.2	11
74	High density of tryptase-positive mast cells in patients with renal cell carcinoma on hemodialysis: correlation with expression of stem cell factor and protease activated receptor-2. <i>Human Pathology</i> , 2012 , 43, 888-97	3.7	11
73	Clinical significance and prognostic value of S100A4 and matrix metalloproteinase-14 in patients with organ-confined bladder cancer. <i>Experimental and Therapeutic Medicine</i> , 2010 , 1, 27-31	2.1	11
72	Expression of X-linked inhibitor of apoptosis protein in human prostate cancer specimens with and without neo-adjuvant hormonal therapy. <i>Journal of Cancer Research and Clinical Oncology</i> , 2010 , 136, 787-93	4.9	11
71	Relationship between changes in prostate cancer cell proliferation, apoptotic index, and expression of apoptosis-related proteins by neoadjuvant hormonal therapy and duration of such treatment. <i>Urology</i> , 2005 , 65, 1238-43	1.6	11

70	Pathological function of prostaglandin E2 receptors in transitional cell carcinoma of the upper urinary tract. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2006 , 448, 822-9	5.1	11
69	Current status and perspective of antiangiogenic therapy for cancer: urinary cancer. <i>International Journal of Clinical Oncology</i> , 2006 , 11, 90-107	4.2	11
68	Immune complexome analysis of antigens in circulating immune complexes from patients with acute cellular rejection after living donor liver transplantation. <i>Transplant Immunology</i> , 2018 , 48, 60-64	1.7	10
67	Thrombospondin-1-derived 4N1K peptide expression is negatively associated with malignant aggressiveness and prognosis in urothelial carcinoma of the upper urinary tract. <i>BMC Cancer</i> , 2012 , 12, 372	4.8	10
66	E1AF expression is associated with extra-prostatic growth and matrix metalloproteinase-7 expression in prostate cancer. <i>Apmis</i> , 2009 , 117, 791-6	3.4	10
65	Pathological significance and prognostic roles of densities of CD57+ cells, CD68+ cells, and mast cells, and their ratios in clear cell renal cell carcinoma. <i>Human Pathology</i> , 2018 , 79, 102-108	3.7	10
64	The Pathological Significance and Prognostic Roles of Thrombospondin-1, and -2, and 4N1K-peptide in Bladder Cancer. <i>Anticancer Research</i> , 2019 , 39, 2317-2324	2.3	9
63	Safety and efficacy of combination therapy with low-dose gemcitabine, paclitaxel, and sorafenib in patients with cisplatin-resistant urothelial cancer. <i>Medical Oncology</i> , 2015 , 32, 235	3.7	9
62	Smoking-induced changes in cancer-related factors in patients with upper tract urothelial cancer. <i>Molecular and Clinical Oncology</i> , 2015 , 3, 287-294	1.6	9
61	Efficacy and safety of systemic chemotherapy and intra-arterial chemotherapy with/without radiotherapy for bladder preservation or as neo-adjuvant therapy in patients with muscle-invasive bladder cancer: a single-centre study of 163 patients. <i>European Journal of Surgical Oncology</i> , 2015 , 41, 261-7	3.6	9
60	Oral Intake of Royal Jelly Has Protective Effects Against Tyrosine Kinase Inhibitor-Induced Toxicity in Patients with Renal Cell Carcinoma: A Randomized, Double-Blinded, Placebo-Controlled Trial. <i>Medicines (Basel, Switzerland)</i> , 2018 , 6,	4.1	9
59	Risk for intravesical recurrence of bladder cancer stratified by the results on two consecutive UroVysion fluorescence in situ hybridization tests: a prospective follow-up study in Japan. <i>International Journal of Clinical Oncology</i> , 2020 , 25, 1163-1169	4.2	8
58	Selective, sensitive and comprehensive detection of immune complex antigens by immune complexome analysis with papain-digestion and elution. <i>Journal of Immunological Methods</i> , 2018 , 461, 85-90	2.5	8
57	Predictive Markers for the Recurrence of Nonmuscle Invasive Bladder Cancer Treated with Intravesical Therapy. <i>Disease Markers</i> , 2015 , 2015, 857416	3.2	8
56	Use of low-dose combined therapy with gemcitabine and paclitaxel for advanced urothelial cancer patients with resistance to cisplatin-containing therapy: a retrospective analysis. <i>Cancer Chemotherapy and Pharmacology</i> , 2012 , 70, 451-9	3.5	8
55	Expression of matrix metalloproteinase-10 in non-metastatic prostate cancer: Correlation with an imbalance in cell proliferation and apoptosis. <i>Oncology Letters</i> , 2010 , 1, 417-421	2.6	8
54	Downregulation of Fes inhibits VEGF-A-induced chemotaxis and capillary-like morphogenesis by cultured endothelial cells. <i>Journal of Cellular and Molecular Medicine</i> , 2007 , 11, 495-501	5.6	8
53	Response of recurrent urachal cancer to gemcitabine and cisplatin therapy: a case report and literature review. <i>Anticancer Research</i> , 2011 , 31, 2335-8	2.3	8

52	Long-term Treatment With Sequential Molecular Targeted Therapy for Xp11.2 Translocation Renal Cell Carcinoma: A Case Report and Review of the Literature. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, e503-e506 ^{3,3}	7
51	Efficacy and safety of sunitinib alternate day regimen in patients with metastatic renal cell carcinoma in Japan: Comparison with standard 4/2 schedule. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2018 , 14, 153-158	1.9 7
50	Stage-specific embryonic antigen-4 is a histological marker reflecting the malignant behavior of prostate cancer. <i>Glycoconjugate Journal</i> , 2019 , 36, 409-418	3 7
49	Expression of Class III Beta-tubulin Predicts Prognosis in Patients with Cisplatin-resistant Bladder Cancer Receiving Paclitaxel-based Second-line Chemotherapy. <i>Anticancer Research</i> , 2018 , 38, 1629-1635 ^{2,3}	7
48	Efficacy of Tadalafil Therapy and Changes in Oxidative Stress Levels in Male Patients with Lower Urinary Tract Symptoms and Overactive Bladder. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2020 , 12, 47-53	1.9 7
47	Clinical significance and predictive value of prostaglandin E2 receptors (EP) 1 - 4 in patients with renal cell carcinoma. <i>Anticancer Research</i> , 2011 , 31, 597-605	2.3 7
46	Stromal expression of Fer suppresses tumor progression in renal cell carcinoma and is a predictor of survival. <i>Oncology Letters</i> , 2017 , 13, 834-840	2.6 6
45	Neoadjuvant hormonal therapy for low-risk prostate cancer induces biochemical recurrence after radical prostatectomy via increased lymphangiogenesis-related parameters. <i>Prostate</i> , 2017 , 77, 1408-1415 ^{4,2}	6
44	Downregulation of the c-Fes protein-tyrosine kinase inhibits the proliferation of human renal carcinoma cells. <i>International Journal of Oncology</i> , 2009 , 34, 89-96	1 6
43	Efficacy of mirabegron for overactive bladder with human T cell lymphotropic virus-1 associated myelopathy. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, O65-O70	1.9 6
42	Pathological significance and prognostic significance of FES expression in bladder cancer vary according to tumor grade. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018 , 144, 21-31	4.9 5
41	Predictive value of serum macrophage colony-stimulating factor for development of aortic calcification in haemodialysis patients: a 6 year longitudinal study. <i>Nephrology Dialysis Transplantation</i> , 2005 , 20, 1647-52	4.3 5
40	High Consumption of Green Tea Suppresses Urinary Tract Recurrence of Urothelial Cancer Down-regulation of Human Antigen-R Expression in Never Smokers. <i>In Vivo</i> , 2018 , 32, 721-729	2.3 5
39	TFE3-renal carcinoma in an adult patient: a case with strong expression of phosphorylated hepatocyte growth factor (HGFR)/Met. <i>Pathology Research and Practice</i> , 2009 , 205, 57-61	3.4 4
38	Relationship Between Stage-specific Embryonic Antigen-4 and Anti-cancer Effects of Neoadjuvant Hormonal Therapy in Prostate Cancer. <i>Anticancer Research</i> , 2020 , 40, 5567-5575	2.3 4
37	Oral intake of royal jelly improves anti-cancer effects and suppresses adverse events of molecular targeted therapy by regulating TNF- α and TGF- β in renal cell carcinoma: A preliminary study based on a randomized double-blind clinical trial. <i>Molecular and Clinical Oncology</i> , 2020 , 13, 29	1.6 4
36	Present Status, Limitations and Future Directions of Treatment Strategies Using Fucoidan-Based Therapies in Bladder Cancer. <i>Cancers</i> , 2020 , 12,	6.6 4
35	Pharmacological Effects and Potential Clinical Usefulness of Polyphenols in Benign Prostatic Hyperplasia. <i>Molecules</i> , 2021 , 26,	4.8 4

34	Pathological and Pharmacological Roles of Mitochondrial Reactive Oxygen Species in Malignant Neoplasms: Therapies Involving Chemical Compounds, Natural Products, and Photosensitizers. <i>Molecules</i> , 2020 , 25,	4.8	3
33	Human antigen R as a predictive marker for response to gemcitabine-based chemotherapy in advanced cisplatin-resistant urothelial cancer. <i>Oncology Letters</i> , 2017 , 13, 811-818	2.6	2
32	Prosultiamine for treatment of lower urinary tract dysfunction accompanied by human T-lymphotropic virus type 1-associated myelopathy/tropical spastic paraparesis. <i>International Journal of Urology</i> , 2018 , 25, 54-60	2.3	2
31	Four-weekly Low-dose Gemcitabine and Paclitaxel in Patients With Platinum-resistant Urothelial Cancer and Performance Status 2/3. <i>In Vivo</i> , 2019 , 33, 2217-2224	2.3	2
30	Changes in Lymphangiogenesis and Vascular Endothelial Growth Factor Expression by Neo-Adjuvant Hormonal Therapy in Prostate Cancer Patients. <i>Prostate</i> , 2017 , 77, 255-262	4.2	2
29	Recovery from life-threatening pelvic organ prolapse in an 80-year-old Japanese woman: a case report. <i>Clinical Case Reports (discontinued)</i> , 2014 , 2, 118-21	0.7	2
28	Efficacy of salt reduction for managing overactive bladder symptoms: a prospective study in patients with excessive daily salt intake. <i>Scientific Reports</i> , 2021 , 11, 4046	4.9	2
27	Intravesical mitomycin C (MMC) and MMC+cytosine arabinoside for non-muscle-invasive bladder cancer: a randomised clinical trial. <i>BJU International</i> , 2021 ,	5.6	2
26	Renal dysfunction due to hydronephrosis by SAPHO syndrome: a case report. <i>Clinical Case Reports (discontinued)</i> , 2015 , 3, 686-9	0.7	1
25	Oral administration of E-type prostanoid (EP) 1 receptor antagonist suppresses carcinogenesis and development of prostate cancer via upregulation of apoptosis in an animal model. <i>Scientific Reports</i> , 2021 , 11, 20279	4.9	1
24	Response to Re: Reconsideration of the clinical and histopathological significance of angiogenesis in prostate cancer: Usefulness and limitations of microvessel density measurement. <i>International Journal of Urology</i> , 2016 , 23, 105-6	2.3	1
23	Detection of Novel Urine Markers Using Immune Complexome Analysis in Bladder Cancer Patients: A Preliminary Study. <i>In Vivo</i> , 2021 , 35, 2073-2080	2.3	1
22	Pathological Significance and Prognostic Roles of Thrombospondin-3, 4 and 5 in Bladder Cancer. <i>In Vivo</i> , 2021 , 35, 1693-1701	2.3	1
21	Presurgical bladder wall thickness is a useful marker to predict the postsurgical improvement of symptoms in patients with pelvic organ prolapse-related overactive bladder. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2021 , 13, 347-355	1.9	1
20	Age-associated bladder and urethral coordination impairment and changes in urethral oxidative stress in rats. <i>Life Sciences</i> , 2021 , 279, 119690	6.8	1
19	Effect of maintenance chemotherapy with gemcitabine and paclitaxel on recurrent squamous cell carcinoma of the bladder: a case report. <i>Anticancer Research</i> , 2011 , 31, 4465-8	2.3	1
18	Effect of oral intake of royal jelly on endothelium function in hemodialysis patients: study protocol for multicenter, double-blind, randomized control trial.. <i>Trials</i> , 2021 , 22, 950	2.8	1
17	Pathological roles of c-Met in bladder cancer: Association with cyclooxygenase-2, heme oxygenase-1, vascular endothelial growth factor-A and programmed death ligand 1. <i>Oncology Letters</i> , 2020 , 20, 135-144	2.6	0

16	Excessive accumulation of visceral fat is associated with lower urinary symptoms including overactive bladder in female patients. <i>International Journal of Urology</i> , 2021 , 28, 397-403	2.3	○
15	Ceruloplasmin Levels in Cancer Tissues and Urine Are Significant Biomarkers of Pathological Features and Outcome in Bladder Cancer. <i>Anticancer Research</i> , 2021 , 41, 3815-3823	2.3	○
14	Editorial Comment to Nutrient intake and urinary incontinence in Korean women: A propensity score-matched analysis from the Korea National Health and Nutrition Examination Survey data. <i>International Journal of Urology</i> , 2017 , 24, 797-798	2.3	
13	Editorial comment on: core biopsies of renal tumors: a study on diagnostic accuracy, interobserver, and intraobserver variability. <i>European Urology</i> , 2008 , 53, 1226	10.2	
12	In vitro method for diagnosing and monitoring renal cell carcinoma using matrix metalloprotease-7 as humoral biomarker. <i>Expert Opinion on Medical Diagnostics</i> , 2007 , 1, 433-5		
11	Alternative Treatment with Every-Other-Day Dosing of Sunitinib for Metastatic Renal Cell Carcinoma: Extended Follow-Up.. <i>Urologia Internationalis</i> , 2022 , 1-7	1.9	
10	Intravesical injection of onabotulinumtoxinA in neurogenic overactive bladder patients with human T-cell leukemia virus type 1-associated myelopathy: A single-institution case series. <i>IJU Case Reports</i> , 2021 , 4, 251-254	0.5	
9	Editorial Comment to Green-light laser en bloc resection versus conventional transurethral resection for initial non-muscle-invasive bladder cancer: A randomized controlled trial. <i>International Journal of Urology</i> , 2021 , 28, 860-861	2.3	
8	Stage-specific Embryogenic Antigen-4 Expression in Castration-resistant Prostate Cancer and its Correlation With the Androgen Receptor. <i>Anticancer Research</i> , 2021 , 41, 3327-3335	2.3	
7	Editorial Comment from Dr Miyata and Dr Sakai to Impact of multimodal treatment on prognosis for patients with metastatic upper urinary tract urothelial cancer: Subanalysis of the multi-institutional nationwide case series study of the Japanese Urological Association. <i>International Journal of Urology</i> , 2021 , 28, 333-334	2.3	
6	Editorial Comment to Does cognitive behavioral therapy using a self-check sheet improve night-time frequency in patients with nocturia? Results of a multicenter randomized controlled trial. <i>International Journal of Urology</i> , 2021 , 28, 449	2.3	
5	Pathological significance and prognostic role of LATS2 in prostate cancer. <i>Prostate</i> , 2021 , 81, 1252-1260	4.2	
4	Pathological Roles of Prostaglandin E2-specific E-type Prostanoid Receptors in Hormone-sensitive and Castration-resistant Prostate Cancer. <i>Anticancer Research</i> , 2021 , 41, 4333-4341	2.3	
3	Dialysis Access and Preemptive Kidney Transplantation. <i>Kidney and Dialysis</i> , 2021 , 1, 79-87		
2	Penile erection occurs when penile vascular resistance is decreased and corporal smooth muscle is relaxed. <i>European Urology</i> , 2007 , 52, 259	10.2	
1	A Clinical Trial Evaluating the Usefulness of Tailored Antimicrobial Prophylaxis Using Rectal-culture Screening Media Prior to Transrectal Prostate Biopsy: A Multicenter, Randomized Controlled Trial. <i>Acta Medica Okayama</i> , 2021 , 75, 663-667	0.5	