

Zhu-Qiu Song

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6110537/publications.pdf>

Version: 2024-02-01

22
papers

131
citations

1478505
6
h-index

1281871
11
g-index

25
all docs

25
docs citations

25
times ranked

148
citing authors

#	ARTICLE	IF	CITATIONS
1	Huchimingia, a new genus segregated from Millettia (Leguminosae, Millettieae) based on morphological and molecular evidence. <i>Phytotaxa</i> , 2022, 532, 37-56.	0.3	0
2	Transfer of Millettia pachycarpa and M. entadooides to Derris (Fabaceae), supported by morphological and molecular data. <i>Phytotaxa</i> , 2022, 531, 230-248.	0.3	2
3	Global warming increases latitudinal divergence in flowering dates of a perennial herb in humid regions across eastern Asia. <i>Agricultural and Forest Meteorology</i> , 2021, 296, 108209.	4.8	7
4	<p align="left">Spatholobus discolor; a new synonym of Craspedolobium unijugum (Fabaceae)</p>. <i>Phytotaxa</i> , 2021, 489, 263-274.	0.3	0
5	<p>Millettia lantsangensis is conspecific with Cruddasia insignis (Fabaceae)</p>. <i>Phytotaxa</i> , 2021, 497, 29-38.	0.3	0
6	<p align="left">Millettia puerariooides Prain (Fabaceae:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 5 China</p>. <i>Phytotaxa</i> , 2021, 507, 183-190.	0.3	0
7	Surprising roles of climate in regulating flowering phenology in a subtropical ecosystem. <i>Ecography</i> , 2021, 44, 1379-1390.	4.5	8
8	New combinations and re-lectotypification in the genus Imbralyx (Fabaceae: Millettieae). <i>Phytotaxa</i> , 2021, 525, 223-231.	0.3	1
9	Effects of winter chilling and photoperiod on leaf-out and flowering in a subtropical evergreen broadleaved forest in China. <i>Forest Ecology and Management</i> , 2020, 458, 117766.	3.2	31
10	Flowering phenology of a widespread perennial herb shows contrasting responses to global warming between humid and non-humid regions. <i>Functional Ecology</i> , 2020, 34, 1870-1881.	3.6	22
11	<i>Bulbophyllum yarlungzangboense</i> (Orchidaceae; Epidendroideae; Malaxideae), a new species from Tibet, China. <i>Phytotaxa</i> , 2019, 404, 79.	0.3	5
12	Exclusion of <i>Glochidion coccineum</i> (Phyllanthaceae) from the flora of China. <i>Phytotaxa</i> , 2019, 408, 215-221.	0.3	0
13	<i>Millettia weizhii</i> (Fabaceae: Millettieae), a new species with four-winged pods from a limestone area in Yunnan, China. <i>Nordic Journal of Botany</i> , 2019, 37, .	0.5	0
14	Untangling the taxonomy of the <i>Cladrastis</i> clade (Leguminosae: Papilionoideae) by integrating phylogenetics and ecological evidence. <i>Taxon</i> , 2019, 68, 1189-1203.	0.7	11
15	<i>Millettia sapindifolia</i> is a synonym of <i>Derris yunnanensis</i> (Leguminosae: Millettieae). <i>Nordic Journal of Botany</i> , 2017, 35, 404-410.	0.5	2
16	The identity of <i>Fordia microphylla</i> and lectotypification of <i>Millettia pulchra</i> (Fabaceae: Millettieae). <i>Phytotaxa</i> , 2017, 332, 51.	0.3	0
17	<i>Mycetia griffithii</i> , a new name for <i>Mycetia angustifolia</i> (Hook.f.) Razafim. & B.Bremer (Rubiaceae). <i>Phytotaxa</i> , 2016, 252, 231.	0.3	1
18	<i>Foonchewia coriacea</i>, a new combination to replace <i>F. guangdongensis</i> (Rubiaceae). <i>Nordic Journal of Botany</i> , 2016, 34, 685-689.	0.5	1

#	ARTICLE	IF	CITATIONS
19	<i>Mycetia fangi</i> (Rubiaceae), a New Species from South China, with Notes on <i>M. cauliflora</i>. Systematic Botany, 2016, 41, 229-237.	0.5	4
20	(2296) Proposal to conserve the name <i>Maackia hupehensis</i> against <i>M. chinensis</i> (<i>Leguminosae</i>). Taxon, 2014, 63, 686-687.	0.7	0
21	Leaves and fruits of Bauhinia (Leguminosae, Caesalpinioideae, Cercideae) from the Oligocene Ningming Formation of Guangxi, South China and their biogeographic implications. BMC Evolutionary Biology, 2014, 14, 88.	3.2	30
22	The identity of Ailanthus guangxiensis (Simaroubaceae) and lectotypification of A. integrifolia Lamarck. Phytotaxa, 2014, 173, 177.	0.3	5