
Eyüp Avcı

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6123704/publications.pdf Version: 2024-02-01



Fvã1/1 DAVCÄ-

#	Article	IF	CITATIONS
1	The monocyte counts to HDL cholesterol ratio in obese and lean patients with polycystic ovary syndrome. Reproductive Biology and Endocrinology, 2018, 16, 34.	3.3	50
2	The Ratio of Fibrinogen to Albumin as a Predictor of Contrast-Induced Nephropathy After Carotid Angiography. Angiology, 2019, 70, 458-464.	1.8	27
3	The predictive value of the prognostic nutritional index for postoperative acute kidney injury in patients undergoing on-pump coronary bypass surgery. Journal of Cardiothoracic Surgery, 2019, 14, 74.	1.1	25
4	Placental fractalkine immunoreactivity in preeclampsia and its correlation with histopathological changes in the placenta and adverse pregnancy outcomes*. Journal of Maternal-Fetal and Neonatal Medicine, 2020, 33, 806-815.	1.5	14
5	Increased Serum CRP-Albumin Ratio Is Independently Associated With Severity of Carotid Artery Stenosis. Angiology, 2020, 71, 740-746.	1.8	13
6	Relationship between high-density lipoprotein cholesterol and the red cell distribution width in patients with coronary artery disease. Lipids in Health and Disease, 2018, 17, 53.	3.0	12
7	Combined value of left ventricular ejection fraction and the Model for End-Stage Liver Disease (MELD) score for predicting mortality in patients with acute coronary syndrome who were undergoing percutaneous coronary intervention. BMC Cardiovascular Disorders, 2018, 18, 44.	1.7	11
8	Prognostic value of rising mean platelet volume during hospitalization in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. BMC Cardiovascular Disorders, 2018, 18, 226.	1.7	11
9	Association of the blood urea nitrogen-to-left ventricular ejection fraction ratio with contrast-induced nephropathy in patients with acute coronary syndrome who underwent percutaneous coronary intervention. International Urology and Nephrology, 2019, 51, 475-481.	1.4	5
10	Using Soluble ST2 to Predict Adverse Postoperative Outcomes in Patients with Impaired Left Ventricular Function Undergoing Coronary Bypass Surgery. Medicina (Lithuania), 2019, 55, 572.	2.0	3
11	Correlation Between the Severity of Coronary Artery Ectasia and Monocyte/Lymphocyte, Platelet/Lymphocyte, and HDL/LDL Ratios. Journal of the College of Physicians and SurgeonsPakistan: JCPSP, 2020, 30, 235-239.	0.4	3
12	The Ratio of Fibrinogen to Albumin as a Predictor of Contrast-Induced Nephropathy After Carotid Angiography: Reply. Angiology, 2019, 70, 467-468.	1.8	2
13	Epicardial Adipose Tissue Predicts Severe Mitral Annular Calcification in Patients Aged ≥60 Years. Medical Science Monitor, 2020, 26, e921553.	1.1	2
14	Bleeding risk in patients with acute coronary syndrome in a Turkish population: Results from the Turkish Acute Coronary Syndrome Registry (TACSER) study. European Journal of Clinical Investigation, 2020, 50, e13271.	3.4	1
15	The Association Between the Ratio of Fibrinogen to Albumin With Contrast-Induced Nephropathy After Carotid Angiography: Reply. Angiology, 2020, 71, 98-98.	1.8	0