## Eyüp Avcı

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6123704/publications.pdf Version: 2024-02-01



ΕνΑΊμο Ανοά

#	Article	IF	CITATIONS
1	Placental fractalkine immunoreactivity in preeclampsia and its correlation with histopathological changes in the placenta and adverse pregnancy outcomes*. Journal of Maternal-Fetal and Neonatal Medicine, 2020, 33, 806-815.	1.5	14
2	The Association Between the Ratio of Fibrinogen to Albumin With Contrast-Induced Nephropathy After Carotid Angiography: Reply. Angiology, 2020, 71, 98-98.	1.8	0
3	Increased Serum CRP-Albumin Ratio Is Independently Associated With Severity of Carotid Artery Stenosis. Angiology, 2020, 71, 740-746.	1.8	13
4	Bleeding risk in patients with acute coronary syndrome in a Turkish population: Results from the Turkish Acute Coronary Syndrome Registry (TACSER) study. European Journal of Clinical Investigation, 2020, 50, e13271.	3.4	1
5	Correlation Between the Severity of Coronary Artery Ectasia and Monocyte/Lymphocyte, Platelet/Lymphocyte, and HDL/LDL Ratios. Journal of the College of Physicians and SurgeonsPakistan: JCPSP, 2020, 30, 235-239.	0.4	3
6	Epicardial Adipose Tissue Predicts Severe Mitral Annular Calcification in Patients Aged ≥60 Years. Medical Science Monitor, 2020, 26, e921553.	1.1	2
7	Using Soluble ST2 to Predict Adverse Postoperative Outcomes in Patients with Impaired Left Ventricular Function Undergoing Coronary Bypass Surgery. Medicina (Lithuania), 2019, 55, 572.	2.0	3
8	The predictive value of the prognostic nutritional index for postoperative acute kidney injury in patients undergoing on-pump coronary bypass surgery. Journal of Cardiothoracic Surgery, 2019, 14, 74.	1.1	25
9	The Ratio of Fibrinogen to Albumin as a Predictor of Contrast-Induced Nephropathy After Carotid Angiography: Reply. Angiology, 2019, 70, 467-468.	1.8	2
10	Association of the blood urea nitrogen-to-left ventricular ejection fraction ratio with contrast-induced nephropathy in patients with acute coronary syndrome who underwent percutaneous coronary intervention. International Urology and Nephrology, 2019, 51, 475-481.	1.4	5
11	The Ratio of Fibrinogen to Albumin as a Predictor of Contrast-Induced Nephropathy After Carotid Angiography. Angiology, 2019, 70, 458-464.	1.8	27
12	Combined value of left ventricular ejection fraction and the Model for End-Stage Liver Disease (MELD) score for predicting mortality in patients with acute coronary syndrome who were undergoing percutaneous coronary intervention. BMC Cardiovascular Disorders, 2018, 18, 44.	1.7	11
13	The monocyte counts to HDL cholesterol ratio in obese and lean patients with polycystic ovary syndrome. Reproductive Biology and Endocrinology, 2018, 16, 34.	3.3	50
14	Relationship between high-density lipoprotein cholesterol and the red cell distribution width in patients with coronary artery disease. Lipids in Health and Disease, 2018, 17, 53.	3.0	12
15	Prognostic value of rising mean platelet volume during hospitalization in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. BMC Cardiovascular Disorders, 2018, 18, 226.	1.7	11