Eric O Long

List of Publications by Citations

Source: https://exaly.com/author-pdf/6124947/eric-o-long-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

167	19,073	75	137
papers	citations	h-index	g-index
184	20,744 ext. citations	14.4	6.77
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
167	Repeated genes in eukaryotes. <i>Annual Review of Biochemistry</i> , 1980 , 49, 727-64	29.1	1195
166	Regulation of immune responses through inhibitory receptors. <i>Annual Review of Immunology</i> , 1999 , 17, 875-904	34.7	827
165	Controlling natural killer cell responses: integration of signals for activation and inhibition. <i>Annual Review of Immunology</i> , 2013 , 31, 227-58	34.7	763
164	A human histocompatibility leukocyte antigen (HLA)-G-specific receptor expressed on all natural killer cells. <i>Journal of Experimental Medicine</i> , 1999 , 189, 1093-100	16.6	596
163	Recruitment of tyrosine phosphatase HCP by the killer cell inhibitor receptor. <i>Immunity</i> , 1996 , 4, 77-85	32.3	560
162	Synergy among receptors on resting NK cells for the activation of natural cytotoxicity and cytokine secretion. <i>Blood</i> , 2006 , 107, 159-66	2.2	546
161	Regulation of human NK-cell cytokine and chemokine production by target cell recognition. <i>Blood</i> , 2010 , 115, 2167-76	2.2	532
160	Molecular clones of the p58 NK cell receptor reveal immunoglobulin-related molecules with diversity in both the extra- and intracellular domains. <i>Immunity</i> , 1995 , 2, 439-49	32.3	525
159	Essential role of LAT in T cell development. <i>Immunity</i> , 1999 , 10, 323-32	32.3	462
158	Activation, coactivation, and costimulation of resting human natural killer cells. <i>Immunological Reviews</i> , 2006 , 214, 73-91	11.3	435
157	Cytolytic granule polarization and degranulation controlled by different receptors in resting NK cells. <i>Journal of Experimental Medicine</i> , 2005 , 202, 1001-12	16.6	356
156	Killer cell inhibitory receptors specific for HLA-C and HLA-B identified by direct binding and by functional transfer. <i>Immunity</i> , 1995 , 3, 801-9	32.3	290
155	Cold urticaria, immunodeficiency, and autoimmunity related to PLCG2 deletions. <i>New England Journal of Medicine</i> , 2012 , 366, 330-8	59.2	288
154	Peptide specificity in the recognition of MHC class I by natural killer cell clones. <i>Science</i> , 1995 , 267, 1016	5 -9 3.3	271
153	A myelin basic protein peptide is recognized by cytotoxic T cells in the context of four HLA-DR types associated with multiple sclerosis. <i>Journal of Experimental Medicine</i> , 1991 , 173, 19-24	16.6	267
152	Negative signaling by inhibitory receptors: the NK cell paradigm. <i>Immunological Reviews</i> , 2008 , 224, 70-	84 1.3	266
151	Defective cytotoxic lymphocyte degranulation in syntaxin-11 deficient familial hemophagocytic lymphohistiocytosis 4 (FHL4) patients. <i>Blood</i> , 2007 , 110, 1906-15	2.2	248

150	Activation of NK cells by an endocytosed receptor for soluble HLA-G. <i>PLoS Biology</i> , 2006 , 4, e9	9.7	245
149	Antigen presentation mediated by recycling of surface HLA-DR molecules. <i>Nature</i> , 1995 , 375, 603-6	50.4	238
148	Crystal structure of the human natural killer cell inhibitory receptor KIR2DL1-HLA-Cw4 complex. <i>Nature Immunology</i> , 2001 , 2, 452-60	19.1	220
147	LFA-1 contributes an early signal for NK cell cytotoxicity. <i>Journal of Immunology</i> , 2004 , 173, 3653-9	5.3	213
146	Vav1 dephosphorylation by the tyrosine phosphatase SHP-1 as a mechanism for inhibition of cellular cytotoxicity. <i>Molecular and Cellular Biology</i> , 2003 , 23, 6291-9	4.8	210
145	Cutting edge: induction of IFN-gamma production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. <i>Journal of Immunology</i> , 2001 , 167, 1877-81	5.3	208
144	Minimal requirement for induction of natural cytotoxicity and intersection of activation signals by inhibitory receptors. <i>Blood</i> , 2009 , 114, 2657-66	2.2	194
143	Cell surface HLA-DR-invariant chain complexes are targeted to endosomes by rapid internalization. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 8581-5	11.5	189
142	Processing pathways for presentation of cytosolic antigen to MHC class II-restricted T cells. <i>Nature</i> , 1992 , 357, 702-4	50.4	187
141	Killer cell inhibitory receptors: diversity, specificity, and function. <i>Immunological Reviews</i> , 1997 , 155, 13) F -/4/10	.0 .
		55 1114 3	184
140	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54	16.6	179
140	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic		179
	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54 Mapping of the class II region of the human major histocompatibility complex by pulsed-field gel	16.6	179
139	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54 Mapping of the class II region of the human major histocompatibility complex by pulsed-field gel electrophoresis. <i>Nature</i> , 1986 , 323, 453-5 KIR2DL4 (CD158d), an NK cell-activating receptor with inhibitory potential. <i>Journal of Immunology</i> ,	16.6 50.4	179 179 175
139	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54 Mapping of the class II region of the human major histocompatibility complex by pulsed-field gel electrophoresis. <i>Nature</i> , 1986 , 323, 453-5 KIR2DL4 (CD158d), an NK cell-activating receptor with inhibitory potential. <i>Journal of Immunology</i> , 2002 , 168, 6208-14 Understanding how combinations of HLA and KIR genes influence disease. <i>Journal of Experimental</i>	16.6 50.4 5·3	179 179 175
139 138	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54 Mapping of the class II region of the human major histocompatibility complex by pulsed-field gel electrophoresis. <i>Nature</i> , 1986 , 323, 453-5 KIR2DL4 (CD158d), an NK cell-activating receptor with inhibitory potential. <i>Journal of Immunology</i> , 2002 , 168, 6208-14 Understanding how combinations of HLA and KIR genes influence disease. <i>Journal of Experimental Medicine</i> , 2005 , 201, 1025-9 The direct binding of a p58 killer cell inhibitory receptor to human histocompatibility leukocyte	16.6 50.4 5·3	179 179 175 171
139 138 137	An endogenous processing pathway in vaccinia virus-infected cells for presentation of cytoplasmic antigens to class II-restricted T cells. <i>Journal of Experimental Medicine</i> , 1990 , 172, 947-54 Mapping of the class II region of the human major histocompatibility complex by pulsed-field gel electrophoresis. <i>Nature</i> , 1986 , 323, 453-5 KIR2DL4 (CD158d), an NK cell-activating receptor with inhibitory potential. <i>Journal of Immunology</i> , 2002 , 168, 6208-14 Understanding how combinations of HLA and KIR genes influence disease. <i>Journal of Experimental Medicine</i> , 2005 , 201, 1025-9 The direct binding of a p58 killer cell inhibitory receptor to human histocompatibility leukocyte antigen (HLA)-Cw4 exhibits peptide selectivity. <i>Journal of Experimental Medicine</i> , 1997 , 185, 1523-8	16.6 50.4 5.3 16.6	179 179 175 171

132	Molecular basis for positive and negative signaling by the natural killer cell receptor 2B4 (CD244). <i>Blood</i> , 2005 , 105, 4722-9	2.2	164
131	Infection of natural killer cells by human herpesvirus 6. <i>Nature</i> , 1993 , 362, 458-62	50.4	164
130	A novel phosphotyrosine motif with a critical amino acid at position -2 for the SH2 domain-mediated activation of the tyrosine phosphatase SHP-1. <i>Journal of Biological Chemistry</i> , 1997 , 272, 13066-72	5.4	161
129	Line of attack: NK cell specificity and integration of signals. <i>Current Opinion in Immunology</i> , 2008 , 20, 344-52	7.8	161
128	Expression of ribosomal DNA insertions in Drosophila melanogaster. <i>Cell</i> , 1979 , 18, 1185-96	56.2	160
127	Adhesion to target cells is disrupted by the killer cell inhibitory receptor. Current Biology, 2000, 10, 777-	-803	152
126	Integrin-dependent organization and bidirectional vesicular traffic at cytotoxic immune synapses. <i>Immunity</i> , 2009 , 31, 99-109	32.3	149
125	Cytotoxic immunological synapses. <i>Immunological Reviews</i> , 2010 , 235, 24-34	11.3	147
124	Structure of the inhibitory receptor for human natural killer cells resembles haematopoietic receptors. <i>Nature</i> , 1997 , 389, 96-100	50.4	147
123	Cell surface expression of class II histocompatibility antigens occurs in the absence of the invariant chain. <i>Journal of Experimental Medicine</i> , 1986 , 164, 1490-504	16.6	133
122	Allelic polymorphism and complexity of the genes for HLA-DR beta-chainsdirect analysis by DNA-DNA hybridization. <i>Nature</i> , 1982 , 300, 372-4	50.4	131
121	Ribosomal DNA in Drosophila melanogaster. I. Isolation and characterization of cloned fragments. Journal of Molecular Biology, 1978 , 126, 749-68	6.5	131
120	Inhibition of natural killer cell activation signals by killer cell immunoglobulin-like receptors (CD158). <i>Immunological Reviews</i> , 2001 , 181, 223-33	11.3	121
119	Two forms of the Ia antigen-associated invariant chain result from alternative initiations at two in-phase AUGs. <i>Cell</i> , 1986 , 47, 619-25	56.2	121
118	A p70 killer cell inhibitory receptor specific for several HLA-B allotypes discriminates among peptides bound to HLA-B*2705. <i>Journal of Experimental Medicine</i> , 1996 , 184, 1585-90	16.6	116
117	KIR2DL4 (CD158d): An activation receptor for HLA-G. Frontiers in Immunology, 2012 , 3, 258	8.4	114
116	Identification of HLA-DR1 beta chain residues critical for binding staphylococcal enterotoxins A and E. <i>Journal of Experimental Medicine</i> , 1992 , 175, 415-24	16.6	114
115	Efficient cDNA expression vectors for stable and transient expression of HLA-DR in transfected fibroblast and lymphoid cells. <i>Human Immunology</i> , 1991 , 31, 229-35	2.3	114

(1981-2012)

114	remodeling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 20596-601	11.5	113
113	Natural killer cell inhibitory receptors block actin cytoskeleton-dependent recruitment of 2B4 (CD244) to lipid rafts. <i>Journal of Experimental Medicine</i> , 2003 , 197, 77-85	16.6	109
112	Vav1 phosphorylation is induced by beta2 integrin engagement on natural killer cells upstream of actin cytoskeleton and lipid raft reorganization. <i>Journal of Experimental Medicine</i> , 2003 , 198, 469-74	16.6	105
111	ICAM-1: getting a grip on leukocyte adhesion. <i>Journal of Immunology</i> , 2011 , 186, 5021-3	5.3	102
110	Specific lysis of allogeneic cells after activation of CD3- lymphocytes in mixed lymphocyte culture. Journal of Experimental Medicine, 1988, 168, 2403-8	16.6	102
109	Interleukin-2 activity can be fine tuned with engineered receptor signaling clamps. <i>Immunity</i> , 2015 , 42, 826-38	32.3	100
108	The alpha 1 domain of the HLA-DR molecule is essential for high-affinity binding of the toxic shock syndrome toxin-1. <i>Nature</i> , 1990 , 346, 474-6	50.4	97
107	Isolation of distinct cDNA clones encoding HLA-DR beta chains by use of an expression assay. Proceedings of the National Academy of Sciences of the United States of America, 1982, 79, 7465-9	11.5	97
106	Synergistic signals for natural cytotoxicity are required to overcome inhibition by c-Cbl ubiquitin ligase. <i>Immunity</i> , 2010 , 32, 175-86	32.3	96
105	Nucleotide sequence of the initiation site for ribosomal RNA transcription in Drosophila melanogaster: comparison of genes with and without insertions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1981 , 78, 1513-7	11.5	92
104	Isolation of cDNA clones encoding HLA-DR alpha chains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1982 , 79, 6979-83	11.5	92
103	NK cell inhibitory receptors prevent tyrosine phosphorylation of the activation receptor 2B4 (CD244). <i>Journal of Immunology</i> , 2000 , 165, 3545-8	5.3	90
102	Negative signaling pathways of the killer cell inhibitory receptor and Fc gamma RIIb1 require distinct phosphatases. <i>Journal of Experimental Medicine</i> , 1997 , 186, 473-8	16.6	88
101	Inhibitory receptor signaling via tyrosine phosphorylation of the adaptor Crk. <i>Immunity</i> , 2008 , 29, 578-88	32.3	88
100	Functional analysis of human NK cells by flow cytometry. <i>Methods in Molecular Biology</i> , 2010 , 612, 335-52	2 .4	85
99	Signal transduction during activation and inhibition of natural killer cells. <i>Current Protocols in Immunology</i> , 2010 , Chapter 11, Unit 11.9B	4	84
98	Intracellular traffic and antigen processing. <i>Trends in Immunology</i> , 1989 , 10, 232-4		82
97	Ribosomal insertion-like elements in Drosophila melanogaster are interspersed with mobile sequences. <i>Cell</i> , 1981 , 25, 399-408	56.2	82

96	Natural killer cell receptors. <i>Current Opinion in Immunology</i> , 1997 , 9, 344-50	7.8	80
95	Coexpression of CD58 or CD48 with intercellular adhesion molecule 1 on target cells enhances adhesion of resting NK cells. <i>Journal of Immunology</i> , 2003 , 170, 294-9	5.3	79
94	Direct binding of a soluble natural killer cell inhibitory receptor to a soluble human leukocyte antigen-Cw4 class I major histocompatibility complex molecule. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 7178-83	11.5	78
93	HLA class I recognition by killer cell Ig-like receptors. <i>Seminars in Immunology</i> , 2000 , 12, 101-8	10.7	75
92	Recognition of virus-infected cells by natural killer cell clones is controlled by polymorphic target cell elements. <i>Journal of Experimental Medicine</i> , 1993 , 178, 961-9	16.6	75
91	Inhibitory MHC class I receptors on NK and T cells: a standard nomenclature. <i>Trends in Immunology</i> , 1996 , 17, 100		70
90	Pathways of viral antigen processing and presentation to CTL: defined by the mode of virus entry?. <i>Trends in Immunology</i> , 1989 , 10, 45-8		70
89	The adaptor protein Crk controls activation and inhibition of natural killer cells. <i>Immunity</i> , 2012 , 36, 600)-31 .3	69
88	Restriction analysis of spacers in ribosomal DNA of Drosophila melanogaster. <i>Nucleic Acids Research</i> , 1979 , 7, 205-15	20.1	66
87	Peptide loading onto recycling HLA-DR molecules occurs in early endosomes. <i>European Journal of Immunology</i> , 1998 , 28, 799-804	6.1	65
86	Isoforms of the invariant chain regulate transport of MHC class II molecules to antigen processing compartments. <i>Journal of Cell Biology</i> , 1996 , 133, 281-91	7.3	65
85	Regulation through inhibitory receptors: Lessons from natural killer cells. <i>Trends in Cell Biology</i> , 1997 , 7, 473-9	18.3	64
84	Tumor cell recognition by natural killer cells. Seminars in Cancer Biology, 2002, 12, 57-61	12.7	64
83	New nomenclature for MHC receptors. <i>Nature Immunology</i> , 2001 , 2, 661	19.1	63
82	Tethering of intercellular adhesion molecule on target cells is required for LFA-1-dependent NK cell adhesion and granule polarization. <i>Journal of Immunology</i> , 2010 , 185, 2918-26	5.3	62
81	Alternative pathways in the processing of ribosomal RNA precursor in Drosophila melanogaster. <i>Journal of Molecular Biology</i> , 1980 , 138, 873-8	6.5	61
80	Ia-negative B-cell variants reveal a coordinate regulation in the transcription of the HLA class II gene family. <i>Immunogenetics</i> , 1984 , 19, 349-53	3.2	57
79	Binding of a soluble p70 killer cell inhibitory receptor to HLA-B*5101: requirement for all three p70 immunoglobulin domains. <i>European Journal of Immunology</i> , 1997 , 27, 568-71	6.1	56

(2011-2018)

78	NK cells inhibit growth in red blood cells via antibody-dependent cellular cytotoxicity. <i>ELife</i> , 2018 , 7,	8.9	53
77	T cell response to myelin basic protein in the context of the multiple sclerosis-associated HLA-DR15 haplotype: peptide binding, immunodominance and effector functions of T cells. <i>Journal of Neuroimmunology</i> , 1997 , 77, 195-203	3.5	52
76	Ready for prime time: NK cell priming by dendritic cells. <i>Immunity</i> , 2007 , 26, 385-7	32.3	52
75	Spontaneous clustering and tyrosine phosphorylation of NK cell inhibitory receptor induced by ligand binding. <i>Journal of Immunology</i> , 2003 , 170, 6107-14	5.3	52
74	Presentation of a cytosolic antigen by major histocompatibility complex class II molecules requires a long-lived form of the antigen. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 14692-7	11.5	52
73	Complementary phosphorylation sites in the adaptor protein SLP-76 promote synergistic activation of natural killer cells. <i>Science Signaling</i> , 2012 , 5, ra49	8.8	51
72	The SH2 domain-containing inositol 5@phosphatase (SHIP) recruits the p85 subunit of phosphoinositide 3-kinase during FcgammaRIIb1-mediated inhibition of B cell receptor signaling. <i>Journal of Biological Chemistry</i> , 1999 , 274, 7489-94	5.4	49
71	DNA-PKcs controls an endosomal signaling pathway for a proinflammatory response by natural killer cells. <i>Science Signaling</i> , 2010 , 3, ra14	8.8	48
70	HLA-DR-restricted presentation of purified myelin basic protein is independent of intracellular processing. <i>European Journal of Immunology</i> , 1997 , 27, 941-51	6.1	46
69	Cutting edge: NK cell licensing modulates adhesion to target cells. <i>Journal of Immunology</i> , 2013 , 191, 3981-5	5.3	45
68	A signaling network stimulated by 2 integrin promotes the polarization of lytic granules in cytotoxic cells. <i>Science Signaling</i> , 2014 , 7, ra96	8.8	44
67	Stress signals activate natural killer cells. <i>Journal of Experimental Medicine</i> , 2002 , 196, 1399-402	16.6	43
66	Two modes of lytic granule fusion during degranulation by natural killer cells. <i>Immunology and Cell Biology</i> , 2011 , 89, 728-38	5	41
65	Adaptive NK cells in people exposed to correlate with protection from malaria. <i>Journal of Experimental Medicine</i> , 2019 , 216, 1280-1290	16.6	40
64	Antigen presentation to HLA class II-restricted measles virus-specific T-cell clones can occur in the absence of the invariant chain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988 , 85, 1209-12	11.5	39
63	Expression of a killer cell receptor-like gene in plastic regions of the central nervous system. <i>Journal of Neuroimmunology</i> , 2005 , 161, 177-82	3.5	38
62	Isolation of cDNA clones for the p33 invariant chain associated with HLA-DR antigens. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1983 , 80, 5714-8	11.5	37
61	I integrin induces TCRESyk-phospholipase C-[phosphorylation and paxillin-dependent granule polarization in human NK cells. <i>Journal of Immunology</i> , 2011 , 186, 2998-3005	5.3	35

60	HLA-DR polymorphism affects the interaction with CD4. <i>Journal of Experimental Medicine</i> , 1995 , 182, 733-41	16.6	34
59	Expression of the ribosomal DNA insertions in bobbed mutants of Drosophila melanogaster. <i>Molecular Genetics and Genomics</i> , 1981 , 182, 377-84		34
58	Recruitment of activation receptors at inhibitory NK cell immune synapses. <i>PLoS ONE</i> , 2008 , 3, e3278	3.7	34
57	T cell recognition of an HLA-A2-restricted epitope derived from a cleaved signal sequence. <i>Journal of Experimental Medicine</i> , 1994 , 180, 1989-94	16.6	33
56	Structural relationship of the SB beta-chain gene to HLA-D-region genes and murine I-region genes. <i>Nature</i> , 1984 , 310, 233-5	50.4	31
55	Human NK cell receptor KIR2DS4 detects a conserved bacterial epitope presented by HLA-C. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 12964-12973	3 ^{11.5}	29
54	Palmitoylation of MICA, a ligand for NKG2D, mediates its recruitment to membrane microdomains and promotes its shedding. <i>European Journal of Immunology</i> , 2011 , 41, 3667-76	6.1	29
53	High-affinity oligoclonal TCRs define effective adoptive T cell therapy targeting mutant KRAS-G12D. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 12826-12835	11.5	28
52	Structural requirements for pairing of alpha and beta chains in HLA-DR and HLA-DP molecules. <i>Journal of Experimental Medicine</i> , 1990 , 171, 615-28	16.6	28
51	Distinct role of rab27a in granule movement at the plasma membrane and in the cytosol of NK cells. <i>PLoS ONE</i> , 2010 , 5, e12870	3.7	28
50	Canonical and Cross-reactive Binding of NK Cell Inhibitory Receptors to HLA-C Allotypes Is Dictated by Peptides Bound to HLA-C. <i>Frontiers in Immunology</i> , 2017 , 8, 193	8.4	26
49	Lytic granule polarization, rather than degranulation, is the preferred target of inhibitory receptors in NK cells. <i>Journal of Immunology</i> , 2010 , 185, 4698-704	5.3	26
48	Sequences in both class II major histocompatibility complex alpha and beta chains contribute to the binding of the superantigen toxic shock syndrome toxin 1. <i>Journal of Experimental Medicine</i> , 1992 , 175, 1301-5	16.6	26
47	Recombination within the HLA-D region. Correlation of molecular genotyping with functional data. Journal of Experimental Medicine, 1984 , 160, 222-38	16.6	25
46	Molecular analysis of the genes for human class II antigens of the major histocompatibility complex. <i>Human Immunology</i> , 1983 , 8, 113-21	2.3	23
45	CD28 Homolog Is a Strong Activator of Natural Killer Cells for Lysis of B7H7 Tumor Cells. <i>Cancer Immunology Research</i> , 2019 , 7, 939-951	12.5	22
44	Viral evasion of NK-cell activation. <i>Trends in Immunology</i> , 2005 , 26, 403-5	14.4	21
43	KIR enrichment at the effector-target cell interface is more sensitive than signaling to the strength of ligand binding. <i>European Journal of Immunology</i> , 2003 , 33, 1084-93	6.1	21

(2013-2000)

42	Cobalt-mediated dimerization of the human natural killer cell inhibitory receptor. <i>Journal of Biological Chemistry</i> , 2000 , 275, 23700-6	5.4	19
41	Natural killer cells and mast cells from gp49B null mutant mice are functional. <i>Molecular and Cellular Biology</i> , 2000 , 20, 7178-82	4.8	19
40	Recognition of intracellular measles virus antigens by HLA class II restricted measles virus-specific cytotoxic T lymphocytes. <i>Annals of the New York Academy of Sciences</i> , 1988 , 540, 352-3	6.5	19
39	-endocytosis of intact IL-15R-L-15 complex from presenting cells into NK cells favors signaling for proliferation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 522-531	11.5	19
38	NK Cell Proliferation Induced by IL-15 Transpresentation Is Negatively Regulated by Inhibitory Receptors. <i>Journal of Immunology</i> , 2015 , 195, 4810-21	5.3	18
37	Distinct binding sites on HLA-DR for invariant chain and staphylococcal enterotoxins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992 , 89, 9657-61	11.5	18
36	Zinc-Induced Polymerization of Killer-Cell Ig-like Receptor into Filaments Promotes Its Inhibitory Function at Cytotoxic Immunological Synapses. <i>Molecular Cell</i> , 2016 , 62, 21-33	17.6	17
35	KIR2DL3 and KIR2DL1 show similar impact on licensing of human NK cells. <i>European Journal of Immunology</i> , 2016 , 46, 185-91	6.1	17
34	A disulfide-linked natural killer cell receptor dimer has higher affinity for HLA-C than wild-type monomer. <i>European Journal of Immunology</i> , 2000 , 30, 2692-7	6.1	16
33	Genome-Wide CRISPR Screen Reveals Cancer Cell Resistance to NK Cells Induced by NK-Derived IFN-[[Frontiers in Immunology, 2019 , 10, 2879	8.4	16
32	Cutting edge: NKG2D-dependent cytotoxicity is controlled by ligand distribution in the target cell membrane. <i>Journal of Immunology</i> , 2011 , 186, 5538-42	5.3	12
31	Binding of soluble KIR-Fc fusion proteins to HLA class I. <i>Methods in Molecular Biology</i> , 2000 , 121, 239-50	1.4	12
30	Activated CD3- CD16+ natural killer cells express a subset of the lymphokine genes induced in activated alpha beta + and gamma delta + T cells. <i>Scandinavian Journal of Immunology</i> , 1991 , 33, 247-52	3.4	12
29	Presentation of influenza hemagglutinin peptide in the presence of limited allostimulation by HLA-DR1 transfected human fibroblasts. <i>Human Immunology</i> , 1988 , 21, 173-81	2.3	12
28	Ionomycin Treatment Renders NK Cells Hyporesponsive. <i>PLoS ONE</i> , 2016 , 11, e0150998	3.7	11
27	Inhibitory receptor gp49B regulates eosinophil infiltration during allergic inflammation. <i>Journal of Leukocyte Biology</i> , 2007 , 82, 1531-41	6.5	10
26	Transcription of unrearranged T cell receptor delta genes in CD3- major histocompatibility complex-unrestricted cytotoxic cells. <i>European Journal of Immunology</i> , 1989 , 19, 1973-6	6.1	10
25	Found: a cellular activating ligand for NKp44. <i>Blood</i> , 2013 , 122, 2921-2	2.2	9

24	Plasmodium falciparum-specific IgM B cells dominate in children, expand with malaria, and produce functional IgM. <i>Journal of Experimental Medicine</i> , 2021 , 218,	16.6	9
23	Regulation of immune responses by inhibitory receptors. <i>Advances in Experimental Medicine and Biology</i> , 1998 , 452, 19-28	3.6	8
22	A single amino acid change in inhibitory killer cell Ig-like receptor results in constitutive receptor self-association and phosphorylation. <i>Journal of Immunology</i> , 2015 , 194, 817-26	5.3	7
21	Antagonizing inhibition gets NK cells going. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 10333-4	11.5	7
20	Use of vaccinia virus for functional gene transfer in natural killer cells. <i>Methods in Molecular Biology</i> , 2000 , 121, 265-72	1.4	7
19	Cutting Edge: Quantitative Determination of CD40L Threshold for IL-12 and IL-23 Production from Dendritic Cells. <i>Journal of Immunology</i> , 2018 , 201, 2879-2884	5.3	7
18	Use of transfected Drosophila S2 cells to study NK cell activation. <i>Methods in Molecular Biology</i> , 2010 , 612, 67-88	1.4	6
17	PD-1 Expression on NK Cells in Malaria-Exposed Individuals Is Associated with Diminished Natural Cytotoxicity and Enhanced Antibody-Dependent Cellular Cytotoxicity. <i>Infection and Immunity</i> , 2020 , 88,	3.7	6
16	Comment on "Killer Ig-like receptor 2DL4 does not mediate NK Cell IFN-Iresponses to soluble HLA-G preparations". <i>Journal of Immunology</i> , 2014 , 192, 4003	5.3	5
15	An unusual form of alternative splicing in the HLA-DNA gene. <i>Immunogenetics</i> , 1991 , 33, 124-31	3.2	5
14	TNFR-associated factor 6 and TGF-Eactivated kinase 1 control signals for a senescence response by an endosomal NK cell receptor. <i>Journal of Immunology</i> , 2014 , 192, 714-21	5.3	4
13	Homogenous expression of killer cell immunoglobulin-like receptors (KIR) on polyclonal natural killer cells detected by a monoclonal antibody to KIR2D. <i>Tissue Antigens</i> , 2000 , 56, 240-7		4
12	Complement factor P: promoting the antibacterial activity of natural killer cells. <i>Cellular and Molecular Immunology</i> , 2017 , 14, 797-799	15.4	3
11	Processing requirements for presentation of antigens to T lymphocytes. <i>Current Opinion in Immunology</i> , 1988 , 1, 98-102	7.8	3
10	Author response: NK cells inhibit Plasmodium falciparum growth in red blood cells via antibody-dependent cellular cytotoxicity 2018 ,		3
9	Inhibition-Resistant CARs for NK Cell Cancer Immunotherapy. <i>Trends in Immunology</i> , 2019 , 40, 1078-108	B114.4	3
8	Patients With Natural Killer (NK) Cell Chronic Active Epstein-Barr Virus Have Immature NK Cells and Hyperactivation of PI3K/Akt/mTOR and STAT1 Pathways. <i>Journal of Infectious Diseases</i> , 2020 , 222, 1170	o-7 ₁₇₉	2
7	Structure of a human natural killer cell inhibitory receptor. <i>Transplantation Proceedings</i> , 1999 , 31, 1871	-21.1	2

LIST OF PUBLICATIONS

6	Bone Marrow-Derived Dendritic Cell Cultures from RAG Mice Include IFN-Producing NK Cells. <i>ImmunoHorizons</i> , 2020 , 4, 415-419	2.7	1
5	HLA-C-restricted presentation of a conserved bacterial epitope to an innate NK cell receptor		1
4	NK Cells Equipped With a Chimeric Antigen Receptor That Overcomes Inhibition by HLA Class I for Adoptive Transfer of CAR-NK Cells <i>Frontiers in Immunology</i> , 2022 , 13, 840844	8.4	0
3	How do killer cell Ig-like receptors inhibit natural killer cells? 2001 , 235-241		
2	Tyrosine phosphorylation of an adapter protein induced by NK cell inhibitory receptor contributes to inhibition of cytotoxicity. <i>FASEB Journal</i> , 2008 , 22, 1064.11	0.9	
1	CRISPR Screen to Identify Factors that Render Tumor Cells Sensitive or Resistant to Killing by NK Cells <i>Methods in Molecular Biology</i> , 2022 , 2463, 269-288	1.4	