

# Eghon Guzman

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6127/publications.pdf>

Version: 2024-02-01

13  
papers

57  
citations

1937457

4  
h-index

1719901

7  
g-index

13  
all docs

13  
docs citations

13  
times ranked

54  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of FOLFOX on minimal residual disease in Stage III colon cancer and risk of relapse. <i>Ecancermedalscience</i> , 2019, 13, 935.	0.6	13
2	Subtypes of minimal residual disease, association with Gleason score, risk and time to biochemical failure in pT2 prostate cancer treated with radical prostatectomy. <i>Ecancermedalscience</i> , 2019, 13, 934.	0.6	12
3	Minimal Residual Disease Defines the Risk and Time to Biochemical Failure in Patients with Pt2 and Pt3a Prostate Cancer Treated With Radical Prostatectomy: An Observational Prospective Study. <i>Urology Journal</i> , 2020, 17, 262-270.	0.3	6
4	The association of the neutrophilâ€“lymphocyte ratio with the presence of minimal residual disease and outcome in patients with Stage II colon cancer treated with surgery alone. <i>Colorectal Disease</i> , 2021, 23, 805-813.	0.7	5
5	Subtypes of minimal residual disease and outcome for stage II colon cancer treated by surgery alone. <i>Ecancermedalscience</i> , 2020, 14, 1119.	0.6	5
6	Transient acquired factor XII deficiency associated with moderately severe Covid-19 pneumonia. <i>Hematology, Transfusion and Cell Therapy</i> , 2021, 43, 515-517.	0.1	4
7	The expression of matrix-metalloproteinase-2 in bone marrow micro-metastasis is associated with the presence of circulating prostate cells and a worse prognosis in men treated with radical prostatectomy for prostate cancer. <i>Turkish Journal of Urology</i> , 2020, 46, 186-195.	1.3	4
8	Immune Dysfunction as Measured by the Systemic Immune-Inflammation Index is Associated with the Sub-Type of Minimal Residual Disease and Outcome in Stage II Colon Cancer Treated with Surgery alone. <i>Asian Pacific Journal of Cancer Prevention</i> , 2021, 22, 2391-2397.	0.5	2
9	Subtypes of minimal residual disease and outcome for stage II colon cancer treated by surgery alone. <i>Ecancermedalscience</i> , 2020, 14, 1119.	0.6	2
10	The Epstein criteria predict for organ-confined prostate cancer but not for minimal residual disease and outcome after radical prostatectomy. <i>Turkish Journal of Urology</i> , 2020, 46, 360-366.	1.3	2
11	Management of mild congenital deficiency of Factor XI with a Factor XI inhibitor in pregnancy: A clinical case. <i>Hematology, Transfusion and Cell Therapy</i> , 2021, , .	0.1	1
12	The CAPRA-S score versus subtypes of minimal residual disease to predict biochemical failure after radical prostatectomy. <i>Ecancermedalscience</i> , 2020, 14, 1063.	0.6	1
13	The Epstein criteria predict for organ-confined prostate cancer but not for minimal residual disease and outcome after radical prostatectomy. <i>Turkish Journal of Urology</i> , 2020, 46, 360-366.	1.3	0