

Fabrizio Martelli

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6133477/publications.pdf>

Version: 2024-02-01

50
papers

697
citations

567281

15
h-index

552781

26
g-index

50
all docs

50
docs citations

50
times ranked

880
citing authors

#	ARTICLE	IF	CITATIONS
1	Characterization of the TGF- β 1 signaling abnormalities in the Gata1 ^{low} mouse model of myelofibrosis. <i>Blood</i> , 2013, 121, 3345-3363.	1.4	86
2	Dexamethasone targeted directly to macrophages induces macrophage niches that promote erythroid expansion. <i>Haematologica</i> , 2015, 100, 178-187.	3.5	59
3	Concise Review: Stem Cell-Derived Erythrocytes as Upcoming Players in Blood Transfusion. <i>Stem Cells</i> , 2012, 30, 1587-1596.	3.2	56
4	Altered SDF-1/CXCR4 axis in patients with primary myelofibrosis and in the Gata1 ^{low} mouse model of the disease. <i>Experimental Hematology</i> , 2008, 36, 158-171.	0.4	50
5	The hypomorphic Gata1 ^{low} mutation alters the proliferation/differentiation potential of the common megakaryocytic-erythroid progenitor. <i>Blood</i> , 2007, 109, 1460-1471.	1.4	48
6	Preclinical rationale for TGF- β 2 inhibition as a therapeutic target for the treatment of myelofibrosis. <i>Experimental Hematology</i> , 2016, 44, 1138-1155.e4.	0.4	38
7	Variegation of the phenotype induced by the Gata1 ^{low} mutation in mice of different genetic backgrounds. <i>Blood</i> , 2005, 106, 4102-4113.	1.4	32
8	Mitochondrial alterations induced by serum amine oxidase and spermine on human multidrug resistant tumor cells. <i>Amino Acids</i> , 2004, 26, 273-82.	2.7	31
9	P-Selectin Sustains Extramedullary Hematopoiesis in the Gata1 ^{low} Model of Myelofibrosis. <i>Stem Cells</i> , 2016, 34, 67-82.	3.2	31
10	TGF- β 1 protein trap AVID200 beneficially affects hematopoiesis and bone marrow fibrosis in myelofibrosis. <i>JCI Insight</i> , 2021, 6, .	5.0	31
11	GATA1 insufficiencies in primary myelofibrosis and other hematopoietic disorders: consequences for therapy. <i>Expert Review of Hematology</i> , 2018, 11, 169-184.	2.2	28
12	Gata1 expression driven by the alternative HS2 enhancer in the spleen rescues the hematopoietic failure induced by the hypomorphic Gata1 ^{low} mutation. <i>Blood</i> , 2009, 114, 2107-2120.	1.4	26
13	The thrombopoietin/MPL axis is activated in the Gata1 ^{low} mouse model of myelofibrosis and is associated with a defective RPS14 signature. <i>Blood Cancer Journal</i> , 2017, 7, e572-e572.	6.2	23
14	Thrombopoietin Inhibits Murine Mast Cell Differentiation. <i>Stem Cells</i> , 2008, 26, 912-919.	3.2	20
15	CXCR4-independent rescue of the myeloproliferative defect of the gata1 ^{low} myelofibrosis mouse model by Aplidin®. <i>Journal of Cellular Physiology</i> , 2010, 225, 490-499.	4.1	16
16	Abnormal P-selectin localization during megakaryocyte development determines thrombosis in the gata1 ^{low} model of myelofibrosis. <i>Platelets</i> , 2014, 25, 539-547.	2.3	14
17	A novel interaction between megakaryocytes and activated fibrocytes increases TGF- β 2 bioavailability in the Gata1 ^{low} mouse model of myelofibrosis. <i>American Journal of Blood Research</i> , 2015, 5, 34-61.	0.6	14
18	The Calreticulin control of human stress erythropoiesis is impaired by JAK2V617F in polycythemia vera. <i>Experimental Hematology</i> , 2017, 50, 53-76.	0.4	12

#	ARTICLE	IF	CITATIONS
19	Dynamic regulation of Gata1 expression during the maturation of conventional dendritic cells. <i>Experimental Hematology</i> , 2010, 38, 489-503.e1.	0.4	11
20	Differential localization of P-selectin and von Willebrand factor during megakaryocyte maturation. <i>Biotechnic and Histochemistry</i> , 2010, 85, 157-170.	1.3	10
21	Removal of the Spleen in Mice Alters the Cytokine Expression Profile of the Marrow Microenvironment and Increases Bone Formation. <i>Annals of the New York Academy of Sciences</i> , 2009, 1176, 77-86.	3.8	9
22	The CXCR1/CXCR2 Inhibitor Reparixin Alters the Development of Myelofibrosis in the Gata1 ^{low} Mice. <i>Frontiers in Oncology</i> , 2022, 12, 853484.	2.8	7
23	Evidence for organ-specific stem cell microenvironments. <i>Journal of Cellular Physiology</i> , 2010, 223, 460-470.	4.1	6
24	Phosphoproteomic Landscaping Identifies Non-canonical cKIT Signaling in Polycythemia Vera Erythroid Progenitors. <i>Frontiers in Oncology</i> , 2019, 9, 1245.	2.8	6
25	Aplidin Improves Megakaryocytopoiesis and Halts Neo-Angiogenesis in the Gata1 ^{low} Murine Model of Myelofibrosis. <i>Blood</i> , 2008, 112, 2787-2787.	1.4	6
26	Resident Self-Tissue of Proinflammatory Cytokines Rather Than Their Systemic Levels Correlates with Development of Myelofibrosis in Gata1 ^{low} Mice. <i>Biomolecules</i> , 2022, 12, 234.	4.0	6
27	Novel targets to cure primary myelofibrosis from studies on Gata1 ^{low} mice. <i>IUBMB Life</i> , 2020, 72, 131-141.	3.4	5
28	Role of β 1 integrin in thrombocytopoiesis. <i>Faculty Reviews</i> , 2021, 10, 68.	3.9	4
29	Increased Differentiation of Dermal Mast Cells in Mice Lacking the Mpl Gene. <i>Stem Cells and Development</i> , 2009, 18, 1081-1092.	2.1	3
30	Preclinical Rationale for the Use of Crizanlizumab (SEG101) in Myelofibrosis. <i>Blood</i> , 2020, 136, 26-27.	1.4	3
31	The Marine Tunicate-Derived Cyclic Depsipeptide Aplidin Restores Functional Hematopoiesis in the Marrow of the Gata1 ^{low} Mouse Model of Myelofibrosis. <i>Blood</i> , 2009, 114, 3914-3914.	1.4	1
32	hGATA1 Under the Control of a β 4LCR/ β 2-Globin Promoter Rescues the Erythroid but Not the Megakaryocytic Phenotype Induced by the Gata1 ^{low} Mutation in Mice. <i>Frontiers in Genetics</i> , 2021, 12, 720552.	2.3	1
33	SB431542, An Inhibitor of TGF- β 1 Activin Receptor-Like Kinases, Improves the Natural History of Myelofibrosis In Gata1 ^{low} Mice. <i>Blood</i> , 2010, 116, 462-462.	1.4	1
34	Glucocorticoid Regulation of Erythropoiesis in Humans: A Study of Patients with Cushing's Disease. <i>Blood</i> , 2015, 126, 2135-2135.	1.4	1
35	Altered Megakaryocytes Are Associated with Development of Pulmonary Fibrosis in Mice Carrying the Hypomorphic Gata1 ^{low} Mutation. <i>Blood</i> , 2019, 134, 2336-2336.	1.4	1
36	The CXCL1 Inhibitor Reparixin Rescues Myelofibrosis in the Gata1 ^{low} Model of the Disease. <i>Blood</i> , 2021, 138, 3579-3579.	1.4	1

#	ARTICLE	IF	CITATIONS
37	CALR resets the stress-response of erythroid cells and this function is impaired by CALR and JAK2 mutations alike in MPN. <i>Experimental Hematology</i> , 2016, 44, S70.	0.4	0
38	Activation of non-canonical cKIT signalling in erythroid progenitor cells from polycythemia vera. <i>Experimental Hematology</i> , 2017, 53, S77-S78.	0.4	0
39	Murine Mast Cells Express Mpl, the Thrombopoietin Receptor, and Thrombopoietin Is a Potent Regulator of Mast Cell Differentiation.. <i>Blood</i> , 2006, 108, 1335-1335.	1.4	0
40	Abnormal P-Selectin Localization During Megakaryocyte Development Determines Thrombosis in the Gata1low Model of Myelofibrosis.. <i>Blood</i> , 2009, 114, 1907-1907.	1.4	0
41	Transforming Growth Factor \hat{I}^2 (TGF- \hat{I}^2)- and P-Selectin-Dependent Fibroblast Peripoleosis of Megakaryocytes Contributes to Development of Myelofibrosis in Gata1low mice. <i>Blood</i> , 2011, 118, 1741-1741.	1.4	0
42	Increased TGF- \hat{I}^2 expression Cooperates with the Gata1low Mutation in Determining the Gata1low Phenotype in CD1 Mice. <i>Blood</i> , 2012, 120, 3462-3462.	1.4	0
43	The Glucocorticoid Receptor Plays an Important Role in Controlling Lipid Metabolism During Erythroid Maturation. <i>Blood</i> , 2012, 120, 3195-3195.	1.4	0
44	The JAK2 V617F Mutation Disrupts the Regulatory Activity Exerted By Calreticulin on the Glucocorticoid Receptor in Erythroid Cells. <i>Blood</i> , 2015, 126, 5216-5216.	1.4	0
45	Phosphoproteomic Landscaping Unveils Constitutive cKIT Activation in Human Erythroblasts from Polycythemia Vera (PV) Patients. <i>Blood</i> , 2016, 128, 399-399.	1.4	0
46	The Carboxy-Terminal Domain of Calreticulin (CALR) Exports the Glucocorticoid Receptor (GR) from the Nucleus to the Cytoplasm of Human Erythroid Cells Resetting Their Stress Response. <i>Blood</i> , 2016, 128, 545-545.	1.4	0
47	Ribosomal Deficiency Due to Activation of the Thrombopoietin Axis May be Involved in the Reduced Levels of GATA1 Expressed By Megakaryocytes (MKs) from the Gata1low Model of Myelofibrosis. <i>Blood</i> , 2016, 128, 4275-4275.	1.4	0
48	The Hypomorphic Gata1low Mutation Induces Fibrosis in Multiple Organs. <i>Blood</i> , 2018, 132, 3059-3059.	1.4	0
49	Human GATA1 Driven By the Human \hat{I}^2 -Globin Promoter Rescues the Erythroid but Not the Megakaryocytic Phenotype Induced in Mice By the Gata1low Mutation. <i>Blood</i> , 2018, 132, 1042-1042.	1.4	0
50	The Glucocorticoid Receptor-Dependent Stress Response in Human Erythropoiesis Is BCL11A-Dependent. <i>Blood</i> , 2021, 138, 939-939.	1.4	0