Mahmoud Shoor

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6136729/publications.pdf

Version: 2024-02-01

1478505 1372567 11 106 10 6 citations h-index g-index papers 11 11 11 141 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	The use of native turf mixtures to approach sustainable lawn in urban landscapes. Urban Forestry and Urban Greening, 2013, 12, 532-536.	5.3	24
2	Assessment of phytochemical and agro-morphological variability among different wild accessions of Mentha longifolia L. cultivated in field condition. Industrial Crops and Products, 2019, 140, 111698.	5 . 2	23
3	The performance of Brevicoryne brassicae on ornamental cabbages grown in CO2-enriched atmospheres. Journal of Asia-Pacific Entomology, 2012, 15, 249-253.	0.9	14
4	Interaction Effect of CO2 Enrichment and Nutritional Conditions on Physiological Characteristics, Essential Oil and Yield of Lemon Balm (Melissa officinalis L.). Notulae Scientia Biologicae, 2012, 4, 121-130.	0.4	11
5	Effects of drought stress on cold hardiness of non-acclimated viola (Viola  ×  wittrockiana  lona Gold) Tj	ЕЈ. Q q1 1 ().784314 rg
6	Title is missing!. Notulae Scientia Biologicae, 2011, 3, 100-104.	0.4	8
7	Association of Genetic Structure and Diversity in Iranian Wild Germplasms of <i>Mentha longifolia </i> L. Based on Phenotypical, Biochemical, and Molecular Markers. Chemistry and Biodiversity, 2021, 18, e2001044.	2.1	5
8	Flowering times and some growth indicators of strawberry were affected by physical properties of the growing media. Scientia Horticulturae, 2020, 272, 109601.	3.6	4
9	The effects of three levels of irrigation water on the improvement of cold tolerance of acclimated viola. Acta Physiologiae Plantarum, 2020, 42, 1.	2.1	4
10	Influence of Water Stress on Agro-Morphological Traits and Essential Oil Content Among Iranian Genotypes of Mentha longifolia. Proceedings of the National Academy of Sciences India Section B - Biological Sciences, 2019, 89, 1219-1230.	1.0	3
11	Aphid suitability for the predatory hoverfly <i>Episyrphus balteatus</i> atmospheric CO ₂ and sinigrin. Entomological Science, 2018, 21, 12-21.	0.6	1