Kristine Krakauer

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6138506/kristine-krakauer-publications-by-year.pdf

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

23	302	10	17
papers	citations	h-index	g-index
24	394 ext. citations	5.7	3.01
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
23	Are attenuated positive symptoms and cortisol levels associated?. <i>Schizophrenia Research</i> , 2021 , 228, 621-623	3.6	
22	Generalized neurocognitive impairment in individuals at ultra-high risk for psychosis: The possible key role of slowed processing speed. <i>Brain and Behavior</i> , 2021 , 11, e01962	3.4	3
21	Global fractional anisotropy predicts transition to psychosis after 12 months in individuals at ultra-high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 144, 448-463	6.5	2
20	Changes in negative symptoms are linked to white matter changes in superior longitudinal fasciculus in individuals at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2021 , 237, 192-201	3.6	0
19	Associations between facial affect recognition and neurocognition in subjects at ultra-high risk for psychosis: A case-control study. <i>Psychiatry Research</i> , 2020 , 290, 112969	9.9	2
18	Cerebral Glutamate and Gamma-Aminobutyric Acid Levels in Individuals at Ultra-high Risk for Psychosis and the Association With Clinical Symptoms and Cognition. <i>Biological Psychiatry:</i> Cognitive Neuroscience and Neuroimaging, 2020, 5, 569-579	3.4	7
17	Effectiveness of cognitive remediation in the ultra-high risk state for psychosis. <i>World Psychiatry</i> , 2020 , 19, 401-402	14.4	7
16	Cognitive remediation plus standard treatment versus standard treatment alone for individuals at ultra-high risk of developing psychosis: Results of the FOCUS randomised clinical trial. <i>Schizophrenia Research</i> , 2020 , 224, 151-158	3.6	10
15	No Effects of Cognitive Remediation on Cerebral White Matter in Individuals at Ultra-High Risk for Psychosis-A Randomized Clinical Trial. <i>Frontiers in Psychiatry</i> , 2020 , 11, 873	5	4
14	Baseline measures of cerebral glutamate and GABA levels in individuals at ultrahigh risk for psychosis: Implications for clinical outcome after 12 months. <i>European Psychiatry</i> , 2020 , 63, e83	6	3
13	Interview and questionnaire assessment of cognitive impairment in subjects at ultra-high risk for psychosis: Associations with cognitive test performance, psychosocial functioning, and positive symptoms. <i>Psychiatry Research</i> , 2020 , 294, 113498	9.9	3
12	Widespread higher fractional anisotropy associates to better cognitive functions in individuals at ultra-high risk for psychosis. <i>Human Brain Mapping</i> , 2019 , 40, 5185-5201	5.9	13
11	Multiple measures of HPA axis function in ultra high risk and first-episode schizophrenia patients. <i>Psychoneuroendocrinology</i> , 2018 , 92, 72-80	5	17
10	Premorbid adjustment in individuals at ultra-high risk for developing psychosis: a case-control study. <i>Microbial Biotechnology</i> , 2018 , 12, 839-847	3.3	11
9	Examining speed of processing of facial emotion recognition in individuals at ultra-high risk for psychosis: Associations with symptoms and cognition. <i>Schizophrenia Research</i> , 2018 , 195, 562-563	3.6	4
8	White matter maturation during 12 months in individuals at ultra-high-risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 65-78	6.5	18
7	Psychopathology and social functioning of 42 subjects from a Danish ultra high-risk cohort. <i>Microbial Biotechnology</i> , 2018 , 12, 1181-1187	3.3	6

LIST OF PUBLICATIONS

6	Patterns of white matter microstructure in individuals at ultra-high-risk for psychosis: associations to level of functioning and clinical symptoms. <i>Psychological Medicine</i> , 2017 , 47, 2689-2707	6.9	25
5	Negative symptoms mediate the relationship between neurocognition and function in individuals at ultrahigh risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 135, 250-258	6.5	17
4	EFFECTIVENESS OF DIALECTICAL BEHAVIOR THERAPY VERSUS COLLABORATIVE ASSESSMENT AND MANAGEMENT OF SUICIDALITY TREATMENT FOR REDUCTION OF SELF-HARM IN ADULTS WITH BORDERLINE PERSONALITY TRAITS AND DISORDER-A RANDOMIZED OBSERVER-BLINDED	8.4	64
3	CLINICAL TRIAL. Depression and Anxiety, 2016 , 33, 520-30 Systemic oxidative DNA and RNA damage are not increased during early phases of psychosis: A case control study. <i>Psychiatry Research</i> , 2016 , 241, 201-6	9.9	17
2	Social cognition in patients at ultra-high risk for psychosis: What is the relation to social skills and functioning?. <i>Schizophrenia Research: Cognition</i> , 2016 , 5, 21-27	2.8	37
1	The FOCUS trial: cognitive remediation plus standard treatment versus standard treatment for patients at ultra-high risk for psychosis: study protocol for a randomised controlled trial. <i>Trials</i> , 2015 , 16, 25	2.8	32