

Ki Hyeong Lee

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6142744/publications.pdf>

Version: 2024-02-01

121
papers

21,621
citations

76196

40
h-index

26548

107
g-index

121
all docs

121
docs citations

121
times ranked

14871
citing authors

#	ARTICLE	IF	CITATIONS
1	First-Line Nivolumab Plus Ipilimumab in Advanced NSCLC: 4-Year Outcomes From the Randomized, Open-Label, Phase 3 CheckMate 227 Part 1 Trial. <i>Journal of Thoracic Oncology</i> , 2022, 17, 289-308.	0.5	173
2	Five-Year Survival Outcomes From the PACIFIC Trial: Durvalumab After Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer. <i>Journal of Clinical Oncology</i> , 2022, 40, 1301-1311.	0.8	445
3	Veliparib Plus Carboplatin and Paclitaxel Versus Investigator's Choice of Standard Chemotherapy in Patients With Advanced Non-Squamous Small Cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2022, 23, 214-225.	1.1	14
4	A Phase 1/2 Study of Lazertinib 240 mg in Patients With Advanced EGFR T790M-Positive NSCLC After Previous EGFR Tyrosine Kinase Inhibitors. <i>Journal of Thoracic Oncology</i> , 2022, 17, 558-567.	0.5	43
5	Abstract P1-16-01: Pemetrexed plus vinorelbine versus vinorelbine monotherapy in patients with metastatic breast cancer: A randomized, open-label, multicenter, phase II trial (KCSG-BR15-17). <i>Cancer Research</i> , 2022, 82, P1-16-01-P1-16-01.	0.4	0
6	Diagnostic Value of Ascitic Tumor Markers for Gastric Cancer-associated Malignant Ascites. <i>The Korean Journal of Helicobacter and Upper Gastrointestinal Research</i> , 2022, 22, 38-49.	0.1	3
7	Tiragolumab plus atezolizumab versus placebo plus atezolizumab as a first-line treatment for PD-L1-selected non-small-cell lung cancer (CITYSCAPE): primary and follow-up analyses of a randomised, double-blind, phase 2 study. <i>Lancet Oncology</i> , 2022, 23, 781-792.	5.1	150
8	Association of depth of target lesion response to brigatinib with outcomes in patients with ALK inhibitor-naïve ALK+ NSCLC in ALTA-1L. <i>Journal of Clinical Oncology</i> , 2022, 40, 9072-9072.	0.8	0
9	Two-year update from KEYNOTE-799: Pembrolizumab plus concurrent chemoradiation therapy (cCRT) for unresectable, locally advanced, stage III NSCLC. <i>Journal of Clinical Oncology</i> , 2022, 40, 8508-8508.	0.8	16
10	A randomized phase II study comparing erlotinib with or without bevacizumab in patients with advanced non-small cell lung cancer (NSCLC) with EGFR mutation. <i>Journal of Clinical Oncology</i> , 2022, 40, 9107-9107.	0.8	0
11	Phase I Study of the Efficacy and Safety of Ramucirumab in Combination with Osimertinib in Advanced T790M-positive EGFR-mutant Non-Small Cell Lung Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 992-1002.	3.2	36
12	The patient's perspective on treatment with dacomitinib: patient-reported outcomes from the Phase III trial ARCHER 1050. <i>Future Oncology</i> , 2021, 17, 783-794.	1.1	0
13	TOX-expressing terminally exhausted tumor-infiltrating CD8+ T cells are reinvigorated by co-blockade of PD-1 and TIGIT in bladder cancer. <i>Cancer Letters</i> , 2021, 499, 137-147.	3.2	42
14	Updated Overall Survival in a Randomized Study Comparing Dacomitinib with Gefitinib as First-Line Treatment in Patients with Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations. <i>Drugs</i> , 2021, 81, 257-266.	4.9	57
15	Safety and efficacy of first-line dacomitinib in Asian patients with EGFR mutation-positive non-small cell lung cancer: Results from a randomized, open-label, phase 3 trial (ARCHER 1050). <i>Lung Cancer</i> , 2021, 154, 176-185.	0.9	18
16	KEYNOTE-799: Phase 2 trial of pembrolizumab plus platinum chemotherapy and radiotherapy for unresectable, locally advanced, stage 3 NSCLC. <i>Journal of Clinical Oncology</i> , 2021, 39, 8512-8512.	0.8	13
17	Comparison of Bleeding, Hematoma, Pain, and Discomfort After Bone Marrow Examination With or Without Sandbag Compression. <i>Asian Nursing Research</i> , 2021, 15, 150-156.	0.7	0
18	Patient-Reported Outcomes with Durvalumab With or Without Tremelimumab Versus Standard Chemotherapy as First-Line Treatment of Metastatic Non-Small-Cell Lung Cancer (MYSTIC). <i>Clinical Lung Cancer</i> , 2021, 22, 301-312.e8.	1.1	10

#	ARTICLE	IF	CITATIONS
19	Amivantamab in EGFR Exon 20 Insertion-“Mutated Non-“Small-Cell Lung Cancer Progressing on Platinum Chemotherapy: Initial Results From the CHRYSALIS Phase I Study. <i>Journal of Clinical Oncology</i> , 2021, 39, 3391-3402.	0.8	320
20	Cardiac Safety Assessment of Lazertinib: Findings From Patients With EGFR Mutation-Positive Advanced NSCLC and Preclinical Studies. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100224.	0.6	6
21	Phase 1b Open-Label Trial of Afatinib Plus Xentuzumab (BI 836845) in Patients With EGFR Mutation-Positive NSCLC After Progression on EGFR Tyrosine Kinase Inhibitors. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100206.	0.6	3
22	E-Cadherin and Angiopoietin-2 as Potential Biomarkers for Colorectal Cancer With Peritoneal Carcinomatosis. <i>Anticancer Research</i> , 2021, 41, 4497-4504.	0.5	5
23	Brigatinib Versus Crizotinib in ALK Inhibitor-“Naive Advanced ALK-Positive NSCLC: Final Results of Phase 3 ALTA-1L Trial. <i>Journal of Thoracic Oncology</i> , 2021, 16, 2091-2108.	0.5	156
24	Histologic Changes in Non-Small Cell Lung Cancer Under Various Treatments: A Comparison of Histology and Mutation Status in Serial Samples. <i>Cancer Research and Treatment</i> , 2021, , .	1.3	5
25	Pembrolizumab Plus Concurrent Chemoradiation Therapy in Patients With Unresectable, Locally Advanced, Stage III Non-“Small Cell Lung Cancer. <i>JAMA Oncology</i> , 2021, 7, 1351.	3.4	113
26	Three-Year Overall Survival with Durvalumab after Chemoradiotherapy in Stage III NSCLC-“Update from PACIFIC. <i>Journal of Thoracic Oncology</i> , 2020, 15, 288-293.	0.5	328
27	Overall Survival with Osimertinib in Untreated, <i>EGFR</i>-Mutated Advanced NSCLC. <i>New England Journal of Medicine</i> , 2020, 382, 41-50.	13.9	1,725
28	Brigatinib Versus Crizotinib in Advanced ALK Inhibitor-“Naive ALK-Positive Non-“Small Cell Lung Cancer: Second Interim Analysis of the Phase III ALTA-1L Trial. <i>Journal of Clinical Oncology</i> , 2020, 38, 3592-3603.	0.8	224
29	18F-fluorodeoxyglucose positron emission tomography/computed tomography findings in descending necrotizing mediastinitis and cervical vertebral osteomyelitis in a cancer patient. <i>Medicine (United Tj ETQq1 1 0.7843 14 rgBI /Overl</i>	0.7	14
30	Differences in diagnosis, treatment, and survival rate of acute myeloid leukemia with or without disabilities: A national cohort study in the Republic of Korea. <i>Cancer Medicine</i> , 2020, 9, 5335-5344.	1.3	7
31	Bintrafusp Alfa, a Bifunctional Fusion Protein Targeting TGF- β 2 and PD-L1, in Second-Line Treatment of Patients With NSCLC: Results From an Expansion Cohort of a Phase 1 Trial. <i>Journal of Thoracic Oncology</i> , 2020, 15, 1210-1222.	0.5	119
32	Durvalumab With or Without Tremelimumab vs Standard Chemotherapy in First-line Treatment of Metastatic Non-“Small Cell Lung Cancer. <i>JAMA Oncology</i> , 2020, 6, 661.	3.4	446
33	Phase II study of pembrolizumab (pembro) plus platinum doublet chemotherapy and radiotherapy as first-line therapy for unresectable, locally advanced stage III NSCLC: KEYNOTE-799.. <i>Journal of Clinical Oncology</i> , 2020, 38, 9008-9008.	0.8	15
34	Nivolumab + ipilimumab versus platinum-doublet chemotherapy as first-line treatment for advanced non-small cell lung cancer: Three-year update from CheckMate 227 Part 1.. <i>Journal of Clinical Oncology</i> , 2020, 38, 9500-9500.	0.8	42
35	Primary analysis of a randomized, double-blind, phase II study of the anti-TIGIT antibody tiragolumab (tira) plus atezolizumab (atezo) versus placebo plus atezo as first-line (1L) treatment in patients with PD-L1-selected NSCLC (CITYSCAPE).. <i>Journal of Clinical Oncology</i> , 2020, 38, 9503-9503.	0.8	179
36	Amivantamab (JNJ-61186372), an anti-EGFR-MET bispecific antibody, in patients with EGFR exon 20 insertion (exon20ins)-mutated non-small cell lung cancer (NSCLC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 9512-9512.	0.8	54

#	ARTICLE	IF	CITATIONS
37	Two-year follow-up of bintrafusp alfa, a bifunctional fusion protein targeting TGF- β 2 and PD-L1, for second-line (2L) treatment of non-small cell lung cancer (NSCLC).. Journal of Clinical Oncology, 2020, 38, 9558-9558.	0.8	7
38	ctDNA resistance landscape of lazertinib, a third-generation EGFR tyrosine kinase inhibitor (TKI).. Journal of Clinical Oncology, 2020, 38, 9601-9601.	0.8	1
39	The Effect of Disability on the Diagnosis and Treatment of Multiple Myeloma in Korea: A National Cohort Study. Cancer Research and Treatment, 2020, 52, 1-9.	1.3	9
40	Real World Experience of Nivolumab in Non-Small Cell Lung Cancer in Korea. Cancer Research and Treatment, 2020, 52, 1112-1119.	1.3	10
41	A phase III, open-label, randomized study of atezolizumab in combination with carboplatin + paclitaxel + bevacizumab compared with pemetrexed + cisplatin or carboplatin with stage IV non-squamous non-small cell lung cancer (NSCLC) with activating EGFR mutation or ALK translocation (ATLAS Trial).. Journal of Clinical Oncology, 2020, 38, TPS9636-TPS9636.	0.8	2
42	363â€¦Vactosertib and durvalumab as second or later line treatment for PD-L1 positive non-small cell lung cancer: interim result. , 2020, , .		2
43	Effects of dose modifications on the safety and efficacy of dacomitinib for <i>EGFR</i> mutation-positive non-small-cell lung cancer. Future Oncology, 2019, 15, 2795-2805.	1.1	27
44	Pseudoprogression presenting as intestinal perforation in nonâ€small cell lung cancer treated with antiâ€PDâ€1: A case report. Molecular and Clinical Oncology, 2019, 11, 132-134.	0.4	9
45	Safety, tolerability, and anti-tumor activity of olmutinib in non-small cell lung cancer with T790M mutation: A single arm, open label, phase 1/2 trial. Lung Cancer, 2019, 135, 66-72.	0.9	22
46	Lazertinib in patients with EGFR mutation-positive advanced non-small-cell lung cancer: results from the dose escalation and dose expansion parts of a first-in-human, open-label, multicentre, phase 1â€2 study. Lancet Oncology, The, 2019, 20, 1681-1690.	5.1	92
47	Patient-reported outcomes with durvalumab after chemoradiotherapy in stage III, unresectable non-small-cell lung cancer (PACIFIC): a randomised, controlled, phase 3 study. Lancet Oncology, The, 2019, 20, 1670-1680.	5.1	125
48	Ramucirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2019, 20, 1655-1669.	5.1	418
49	A Case of Acquired Amegakaryocytic Thrombocytopenia with Anti-c-mpl Autoantibody: Comparison with Idiopathic Thrombocytopenic Purpura. Acta Haematologica, 2019, 142, 239-243.	0.7	3
50	Quality of life outcomes including neuropathyâ€associated scale from a phase II, multicenter, randomized trial of eribulin plus gemcitabine versus paclitaxel plus gemcitabine as firstâ€line chemotherapy for HER2â€negative metastatic breast cancer: Korean Cancer Study Group Trial (KCSG) Tj ETQqO 0 0 rfgBT /Overlock 10 T	3.7	5
51	Nivolumab plus ipilimumab versus chemotherapy as first-line treatment in advanced nonâ€small-cell lung cancer with high tumour mutational burden: patient-reported outcomes results from the randomised, open-label, phase III CheckMate 227 trial. European Journal of Cancer, 2019, 116, 137-147.	1.3	167
52	First-line afatinib for advanced EGFRm+ NSCLC: Analysis of long-term responders in the LUX-Lung 3, 6, and 7 trials. Lung Cancer, 2019, 133, 10-19.	0.9	25
53	Molecular alterations and poziotinib efficacy, a panâ€HER inhibitor, in human epidermal growth factor receptor 2 (HER2)â€positive breast cancers: Combined exploratory biomarker analysis from a phase II clinical trial of poziotinib for refractory HER2â€positive breast cancer patients. International Journal of Cancer, 2019, 145, 1669-1678.	2.3	14
54	First-line afatinib vs gefitinib for patients with EGFR mutation-positive NSCLC (LUX-Lung 7): impact of afatinib dose adjustment and analysis of mode of initial progression for patients who continued treatment beyond progression. Journal of Cancer Research and Clinical Oncology, 2019, 145, 1569-1579.	1.2	31

#	ARTICLE	IF	CITATIONS
55	Osimertinib versus Standard of Care EGFR TKI as First-Line Treatment in Patients with EGFRm Advanced NSCLC: FLAURA Asian Subset. <i>Journal of Thoracic Oncology</i> , 2019, 14, 99-106.	0.5	82
56	JNJ-61186372 (JNJ-372), an EGFR-cMet bispecific antibody, in EGFR-driven advanced non-small cell lung cancer (NSCLC).. <i>Journal of Clinical Oncology</i> , 2019, 37, 9009-9009.	0.8	74
57	Blood tumor mutational burden (bTMB) and tumor PD-L1 as predictive biomarkers of survival in MYSTIC: First-line durvalumab (D) ± tremelimumab (T) versus chemotherapy (CT) in metastatic (m) NSCLC.. <i>Journal of Clinical Oncology</i> , 2019, 37, 9016-9016.	0.8	20
58	Patient-reported outcomes (PROs) with first-line durvalumab (D) ± tremelimumab (T) versus chemotherapy (CT) in metastatic NSCLC: Results from MYSTIC.. <i>Journal of Clinical Oncology</i> , 2019, 37, 9048-9048.	0.8	2
59	Randomized Phase III Trial of Irinotecan Plus Cisplatin versus Etoposide Plus Cisplatin in Chemotherapy-Naïve Korean Patients with Extensive-Disease Small Cell Lung Cancer. <i>Cancer Research and Treatment</i> , 2019, 51, 119-127.	1.3	23
60	A Randomized, Open-Label, Phase II Study Comparing Pemetrexed Plus Cisplatin Followed by Maintenance Pemetrexed versus Pemetrexed Alone in Patients with Epidermal Growth Factor Receptor (EGFR)-Mutant Non-small Cell Lung Cancer after Failure of First-Line EGFR Tyrosine Kinase Inhibitor: KCSG-LU12-13. <i>Cancer Research and Treatment</i> , 2019, 51, 718-726.	1.3	10
61	Exosomal miR-181b-5p Downregulation in Ascites Serves as a Potential Diagnostic Biomarker for Gastric Cancer-associated Malignant Ascites. <i>Journal of Gastric Cancer</i> , 2019, 19, 301.	0.9	24
62	Pazopanib maintenance after first-line etoposide and platinum chemotherapy in patients with extensive disease small-cell lung cancer: a multicentre, randomised, placebo-controlled Phase II study (KCSG-LU12-07). <i>British Journal of Cancer</i> , 2018, 118, 648-653.	2.9	31
63	Afatinib as First-line Treatment of Older Patients With EGFR Mutation-Positive Non-Small-Cell Lung Cancer: Subgroup Analyses of the LUX-Lung 3, LUX-Lung 6, and LUX-Lung 7 Trials. <i>Clinical Lung Cancer</i> , 2018, 19, e465-e479.	1.1	56
64	Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. <i>New England Journal of Medicine</i> , 2018, 378, 113-125.	13.9	3,530
65	Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations. <i>Journal of Clinical Oncology</i> , 2018, 36, 2244-2250.	0.8	361
66	CNS Response to Osimertinib Versus Standard Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Patients With Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. <i>Journal of Clinical Oncology</i> , 2018, 36, 3290-3297.	0.8	515
67	Diagnostic benefits of the combined use of liquid-based cytology, cell block, and carcinoembryonic antigen immunocytochemistry in malignant pleural effusion. <i>Journal of Thoracic Disease</i> , 2018, 10, 4931-4939.	0.6	29
68	EGFR Mutation Status in Lung Adenocarcinoma-Associated Malignant Pleural Effusion and Efficacy of EGFR Tyrosine Kinase Inhibitors. <i>Cancer Research and Treatment</i> , 2018, 50, 908-916.	1.3	32
69	Long-term Survival after Repeated Local Therapy and Salvage Chemotherapy for Recurrent Metastases from Gastric Cancer: a Case Report and Literature Review. <i>Journal of Gastric Cancer</i> , 2018, 18, 305.	0.9	0
70	Brigatinib versus Crizotinib in ALK-Positive Non-Small-Cell Lung Cancer. <i>New England Journal of Medicine</i> , 2018, 379, 2027-2039.	13.9	691
71	Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung Cancer. <i>New England Journal of Medicine</i> , 2018, 379, 2040-2051.	13.9	2,676
72	Nivolumab in advanced non-small-cell lung cancer patients who failed prior platinum-based chemotherapy. <i>Lung Cancer</i> , 2018, 122, 234-242.	0.9	22

#	ARTICLE	IF	CITATIONS
73	A phase II trial of the pan-HER inhibitor poziotinib, in patients with HER2-positive metastatic breast cancer who had received at least two prior HER2-directed regimens: results of the NOV120101-203 trial. <i>International Journal of Cancer</i> , 2018, 143, 3240-3247.	2.3	46
74	Feasibility and accessibility of electronic patient-reported outcome measures using a smartphone during routine chemotherapy: a pilot study. <i>Supportive Care in Cancer</i> , 2018, 26, 3721-3728.	1.0	21
75	Do health literacy and self-care behaviours affect quality of life in older persons with lung cancer receiving chemotherapy?. <i>International Journal of Nursing Practice</i> , 2018, 24, e12691.	0.8	11
76	Association of ERBB Mutations With Clinical Outcomes of Afatinib- or Erlotinib-Treated Patients With Lung Squamous Cell Carcinoma. <i>JAMA Oncology</i> , 2018, 4, 1189.	3.4	53
77	Dacomitinib (daco) versus gefitinib (gef) for first-line treatment of advanced NSCLC (ARCHER 1050): Final overall survival (OS) analysis.. <i>Journal of Clinical Oncology</i> , 2018, 36, 9004-9004.	0.8	9
78	Results from a second-line (2L) NSCLC cohort treated with M7824 (MSB0011359C), a bifunctional fusion protein targeting TGF- β 2 and PD-L1.. <i>Journal of Clinical Oncology</i> , 2018, 36, 9017-9017.	0.8	10
79	YH25448, a 3rd generation EGFR-TKI, in patients with EGFR-TKI-resistant NSCLC: Phase I/II study results.. <i>Journal of Clinical Oncology</i> , 2018, 36, 9033-9033.	0.8	2
80	Efficacy and safety results of ramucirumab in combination with osimertinib in advanced T790M-positive EGFR-mutant NSCLC.. <i>Journal of Clinical Oncology</i> , 2018, 36, 9053-9053.	0.8	4
81	Role of HER2 copy number amplification and PI3K pathway as a biomarker for patients with HER2+ MBC treating with Poziotinib, pan-HER TKI.. <i>Journal of Clinical Oncology</i> , 2018, 36, e13009-e13009.	0.8	1
82	Prevalence of and factors associated with anxiety and depression in Korean patients with newly diagnosed advanced gastrointestinal cancer. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 585-594.	0.7	28
83	Molecular Screening of Small Biopsy Samples Using Next-Generation Sequencing in Korean Patients with Advanced Non-small Cell Lung Cancer: Korean Lung Cancer Consortium (KLCC-13-01). <i>Journal of Pathology and Translational Medicine</i> , 2018, 52, 148-156.	0.4	15
84	Quality of life(QoL) outcomes including neuropathy associated scale (FACT-T) from a phase II, multicenter, randomized trial of eribulin plus gemcitabine(EG) versus paclitaxel plus gemcitabine(PG) as first-line chemotherapy for human epidermal growth factor receptor 2 (HER2)- negative metastatic breast cancer(MBC): Korean Cancer Study Group trial (KCSG BR13-11).. <i>Journal of Clinical Oncology</i> , 2018, 36, 10117-10117.	0.8	0
85	Clinical trial of nintedanib in patients with recurrent or metastatic salivary gland cancer of the head and neck: A multicenter phase 2 study (Korean Cancer Study Group HN14-01). <i>Cancer</i> , 2017, 123, 1958-1964.	2.0	44
86	Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial. <i>Lancet Oncology</i> , 2017, 18, 1454-1466.	5.1	877
87	Phase II, multicentre, randomised trial of eribulin plus gemcitabine versus paclitaxel plus gemcitabine as first-line chemotherapy in patients with HER2-negative metastatic breast cancer. <i>European Journal of Cancer</i> , 2017, 86, 385-393.	1.3	23
88	Phase III Trial of Ipilimumab Combined With Paclitaxel and Carboplatin in Advanced Squamous Non-Small-Cell Lung Cancer. <i>Journal of Clinical Oncology</i> , 2017, 35, 3449-3457.	0.8	311
89	Open-Label, Multicenter, Phase II Study of Ceritinib in Patients With Non-Small-Cell Lung Cancer Harboring ROS1 Rearrangement. <i>Journal of Clinical Oncology</i> , 2017, 35, 2613-2618.	0.8	260
90	A randomized, phase II study of gefitinib alone versus nimotuzumab plus gefitinib after platinum-based chemotherapy in advanced non-small cell lung cancer (KCSG LU12-01). <i>Oncotarget</i> , 2017, 8, 15943-15951.	0.8	11

#	ARTICLE	IF	CITATIONS
91	Dacomitinib versus gefitinib for the first-line treatment of advanced EGFR mutation positive non-small cell lung cancer (ARCHER 1050): A randomized, open-label phase III trial.. Journal of Clinical Oncology, 2017, 35, LBA9007-LBA9007.	0.8	3
92	PEARLY: A randomized, multicenter, open-label, phase III trial comparing anthracyclines followed by taxane versus anthracyclines followed by taxane plus carboplatin as (neo)adjuvant therapy in patients with early triple-negative breast cancer.. Journal of Clinical Oncology, 2017, 35, TPS587-TPS587.	0.8	6
93	Dacomitinib versus gefitinib for the first-line treatment of advanced EGFR mutation positive non-small cell lung cancer (ARCHER 1050): A randomized, open-label phase III trial.. Journal of Clinical Oncology, 2017, 35, LBA9007-LBA9007.	0.8	21
94	A Phase II Study of Poziotinib in Patients with Epidermal Growth Factor Receptor (<i>EGFR</i>)-Mutant Lung Adenocarcinoma Who Have Acquired Resistance to EGFR Tyrosine Kinase Inhibitors. Cancer Research and Treatment, 2017, 49, 10-19.	1.3	35
95	Randomized Phase II Study of Afatinib Plus Simvastatin Versus Afatinib Alone in Previously Treated Patients with Advanced Nonadenocarcinomatous Non-small Cell Lung Cancer. Cancer Research and Treatment, 2017, 49, 1001-1011.	1.3	43
96	Phase II study of nivolumab in patients with advanced non-small cell lung cancer (NSCLC) in Korea.. Journal of Clinical Oncology, 2017, 35, 92-92.	0.8	0
97	A phase II, multicenter, randomized trial of eribulin plus gemcitabine (EG) vs. paclitaxel plus gemcitabine (PG) in patients with HER2-negative metastatic breast cancer (MBC) as first-line chemotherapy (KCSG BR13-11, NCT02263495).. Journal of Clinical Oncology, 2017, 35, 1082-1082.	0.8	1
98	Afatinib versus gefitinib as first-line treatment of patients with EGFR mutation-positive non-small-cell lung cancer (LUX-Lung 7): a phase 2B, open-label, randomised controlled trial. Lancet Oncology, The, 2016, 17, 577-589.	5.1	950
99	A randomized, multicenter, phase II/III study to determine the optimal dose and to evaluate the efficacy and safety of pegteograstim (GCPGC) on chemotherapy-induced neutropenia compared to pegfilgrastim in breast cancer patients: KCSG PC10-09. Supportive Care in Cancer, 2016, 24, 1709-1717.	1.0	19
100	First-in-human study of HM95573, a novel oral RAF inhibitor, in patients with solid tumors.. Journal of Clinical Oncology, 2016, 34, 2570-2570.	0.8	4
101	Phase II trial of nintedanib in patients with recurrent or metastatic salivary gland cancer: A multicenter phase II study.. Journal of Clinical Oncology, 2016, 34, 6090-6090.	0.8	3
102	A randomized, open label, phase II study comparing pemetrexed plus cisplatin followed by pemetrexed versus pemetrexed alone in EGFR mutant NSCLC patients who have failed first-line EGFR TKI: KCSG-LU12-13.. Journal of Clinical Oncology, 2016, 34, 9043-9043.	0.8	1
103	First-line afatinib (A) vs gefitinib (G) for patients (pts) with EGFR mutation positive (EGFRm+) NSCLC (LUX-Lung 7): Patient-reported outcomes (PROs) and impact of dose modifications on efficacy and adverse events (AEs).. Journal of Clinical Oncology, 2016, 34, 9046-9046.	0.8	11
104	BI 1482694 (HM61713), an EGFR mutant-specific inhibitor, in T790M+ NSCLC: Efficacy and safety at the RP2D.. Journal of Clinical Oncology, 2016, 34, 9055-9055.	0.8	27
105	Open-label, multicenter, randomized phase III trial of pemetrexed/carboplatin doublet vs pemetrexed singlet in chemotherapy-naïve elderly patients aged 70 or more with advanced non-squamous non-small cell lung cancer and good performance status.. Journal of Clinical Oncology, 2016, 34, 9081-9081.	0.8	1
106	A Prospective Multicenter Study Evaluating Secondary Adrenal Suppression After Antiemetic Dexamethasone Therapy in Cancer Patients Receiving Chemotherapy: A Korean South West Oncology Group Study. Oncologist, 2015, 20, 1432-1439.	1.9	21
107	Impact of the availability of active cytotoxic agents on the survival of patients with advanced gastric cancer. Oncology Letters, 2015, 10, 2481-2486.	0.8	1
108	Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials. Lancet Oncology, The, 2015, 16, 141-151.	5.1	1,369

#	ARTICLE	IF	CITATIONS
109	Afatinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label randomised controlled phase 3 trial. <i>Lancet Oncology</i> , The, 2015, 16, 897-907.	5.1	389
110	Diagnostic performance of CD66c in lung adenocarcinoma-associated malignant pleural effusion: comparison with CEA, CA 19-9, and CYFRA 21-1. <i>Pathology</i> , 2015, 47, 123-129.	0.3	18
111	Therapeutic effect of anti CEACAM6 monoclonal antibody against lung adenocarcinoma by enhancing anoikis sensitivity. <i>Biomaterials</i> , 2015, 67, 32-41.	5.7	29
112	Updated safety and efficacy results from phase I/II study of HM61713 in patients (pts) with EGFR mutation positive non-small cell lung cancer (NSCLC) who failed previous EGFR-tyrosine kinase inhibitor (TKI).. <i>Journal of Clinical Oncology</i> , 2015, 33, 8084-8084.	0.8	28
113	A phase II, single-arm, efficacy and safety study of poziotinib (NOV120101) in Korean patients with advanced or metastatic lung adenocarcinoma who have acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors.. <i>Journal of Clinical Oncology</i> , 2015, 33, 8085-8085.	0.8	0
114	Prevalence and associated factors of anxiety and depression in patients with metastatic gastrointestinal cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, e20718-e20718.	0.8	0
115	Prospective multicenter study evaluating adrenal suppression after dexamethasone therapy as an antiemetic in cancer patients: a KSWOG (Korean South West Oncology Group) study.. <i>Journal of Clinical Oncology</i> , 2015, 33, 9605-9605.	0.8	0
116	E7080 (lenvatinib) in addition to best supportive care (BSC) versus BSC alone in third-line or greater nonsquamous, non-small cell lung cancer (NSCLC).. <i>Journal of Clinical Oncology</i> , 2014, 32, 8043-8043.	0.8	9
117	Multicenter phase II study of docetaxel and oxaliplatin combination in patients with locally advanced or metastatic biliary tract cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, e15150-e15150.	0.8	0
118	Phase III Study of Afatinib or Cisplatin Plus Pemetrexed in Patients With Metastatic Lung Adenocarcinoma With <i>EGFR</i> Mutations. <i>Journal of Clinical Oncology</i> , 2013, 31, 3327-3334.	0.8	2,854
119	LUX-Lung 3: A randomized, open-label, phase III study of afatinib versus pemetrexed and cisplatin as first-line treatment for patients with advanced adenocarcinoma of the lung harboring EGFR-activating mutations.. <i>Journal of Clinical Oncology</i> , 2012, 30, LBA7500-LBA7500.	0.8	74
120	LUX-Lung 3: A randomized, open-label, phase III study of afatinib versus pemetrexed and cisplatin as first-line treatment for patients with advanced adenocarcinoma of the lung harboring EGFR-activating mutations.. <i>Journal of Clinical Oncology</i> , 2012, 30, LBA7500-LBA7500.	0.8	60
121	A Case of Superwarfarin Intoxication without a Definitive History of Brodifacoum Exposure. <i>The Korean Journal of Hematology</i> , 2009, 44, 53.	0.7	2