

# Matthias Oelke

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6143082/matthias-oelke-publications-by-year.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

58  
papers

3,285  
citations

25  
h-index

57  
g-index

69  
ext. papers

3,916  
ext. citations

4.1  
avg, IF

5.14  
L-index

#	Paper	IF	Citations
58	Sonografische Diagnostik des Harntrakts und Beckens <b>2022</b> , 121-145		
57	What uro-gynecologists should know about sacral neuromodulation (SNM) for the treatment of refractory overactive bladder. <i>Archives of Gynecology and Obstetrics</i> , <b>2019</b> , 299, 1243-1252	2.5	2
56	The International Continence Society (ICS) report on the terminology for adult male lower urinary tract and pelvic floor symptoms and dysfunction. <i>Neurourology and Urodynamics</i> , <b>2019</b> , 38, 433-477	2.3	195
55	Intradetrusor onabotulinumtoxinA injections are significantly more efficacious than oral oxybutynin for treatment of neurogenic detrusor overactivity: results of a randomized, controlled, 24-week trial. <i>Einstein (Sao Paulo, Brazil)</i> , <b>2018</b> , 16, eAO4207	1.2	7
54	Efficacy and safety of tadalafil 5 mg once daily in the treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia in men aged $\geq 5$ years: integrated analyses of pooled data from multinational, randomized, placebo-controlled clinical studies. <i>BJU International</i> , <b>2017</b> , 119, 793-803	5.6	19
53	A practical approach to the management of nocturia. <i>International Journal of Clinical Practice</i> , <b>2017</b> , 71, e13027	2.9	54
52	Efficacy and safety of daily mirabegron 50 mg in male patients with overactive bladder: a critical analysis of five phase III studies. <i>Therapeutic Advances in Urology</i> , <b>2017</b> , 9, 137-154	3.2	29
51	Annual direct and indirect costs attributable to nocturia in Germany, Sweden, and the UK. <i>European Journal of Health Economics</i> , <b>2017</b> , 18, 761-771	3.6	12
50	Ultrasound detrusor wall thickness measurement in combination with bladder capacity can safely detect detrusor underactivity in adult men. <i>World Journal of Urology</i> , <b>2017</b> , 35, 153-159	4	22
49	Incidence of urinary retention during treatment with single tablet combinations of solifenacin+tamsulosin OCASIFor up to 1 year in adult men with both storage and voiding LUTS: A subanalysis of the NEPTUNE/NEPTUNE II randomized controlled studies. <i>PLoS ONE</i> , <b>2017</b> , 12, e0170726	3.7	12
48	Nocturia is often inadequately assessed, diagnosed and treated by physicians: results of an observational, real-life practice database containing 8659 European and US-American patients. <i>International Journal of Clinical Practice</i> , <b>2016</b> , 70, 940-949	2.9	19
47	Unravelling detrusor underactivity: Development of a bladder outlet resistance-Bladder contractility nomogram for adult male patients with lower urinary tract symptoms. <i>Neurourology and Urodynamics</i> , <b>2016</b> , 35, 980-986	2.3	32
46	Use of Structural Equation Modeling to Demonstrate the Differential Impact of Storage and Voiding Lower Urinary Tract Symptoms on Symptom Bother and Quality of Life during Treatment for Lower Urinary Tract Symptoms Associated with Benign Prostatic Hyperplasia. <i>Journal of Urology</i> , <b>2016</b> , 196, 824-30	2.5	10
45	Latest Evidence on the Use of Phosphodiesterase Type 5 Inhibitors for the Treatment of Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. <i>European Urology</i> , <b>2016</b> , 70, 124-133	10.2	85
44	Exploratory Decision-Tree Modeling of Data from the Randomized REACTT Trial of Tadalafil Versus Placebo to Predict Recovery of Erectile Function After Bilateral Nerve-Sparing Radical Prostatectomy. <i>European Urology</i> , <b>2016</b> , 70, 529-37	10.2	15
43	Bladder Instillation Therapy With Hyaluronic Acid and Chondroitin Sulfate Improves Symptoms of Postradiation Cystitis: Prospective Pilot Study. <i>Clinical Genitourinary Cancer</i> , <b>2016</b> , 14, 444-449	3.3	18
42	What is the role of combination drug therapy in the treatment of overactive bladder? ICI-RS 2014. <i>Neurourology and Urodynamics</i> , <b>2016</b> , 35, 288-92	2.3	6

41	Bladder wall thickness in women with symptoms of overactive bladder and detrusor overactivity: Results from the randomised, placebo-controlled shrink study. <i>Neurourology and Urodynamics</i> , <b>2016</b> , 35, 819-25	2.3	13
40	Recommendations for future development of contractility and obstruction nomograms for women. ICI-RS 2014. <i>Neurourology and Urodynamics</i> , <b>2016</b> , 35, 307-11	2.3	10
39	Effects of Tadalafil Once-Daily or On-Demand vs Placebo on Return to Baseline Erectile Function After Bilateral Nerve-Sparing Radical Prostatectomy--Results from a Randomized Controlled Trial (REACTT). <i>Journal of Sexual Medicine</i> , <b>2016</b> , 13, 679-83	1.1	12
38	Efficacy of Continuous Dosing of Tadalafil Once Daily vs Tadalafil On Demand in Clinical Subgroups of Men With Erectile Dysfunction: A Descriptive Comparison Using the Integrated Tadalafil Databases. <i>Journal of Sexual Medicine</i> , <b>2016</b> , 13, 860-75	1.1	17
37	Responder and health-related quality of life analyses in men with lower urinary tract symptoms treated with a fixed-dose combination of solifenacin and tamsulosin oral-controlled absorption system: results from the NEPTUNE study. <i>BJU International</i> , <b>2016</b> , 117, 165-72	5.6	11
36	Expression and distribution of phosphodiesterase isoenzymes in the human male urethra. <i>Urology</i> , <b>2015</b> , 85, 964.e1-6	1.6	5
35	The underactive bladder: a new clinical concept?. <i>European Urology</i> , <b>2015</b> , 68, 351-3	10.2	97
34	Acute Urinary Retention Rates in the General Male Population and in Adult Men With Lower Urinary Tract Symptoms Participating in Pharmacotherapy Trials: A Literature Review. <i>Urology</i> , <b>2015</b> , 86, 654-65	1.6	21
33	Sodium hyaluronate and chondroitin sulfate replenishment therapy can improve nocturia in men with post-radiation cystitis: results of a prospective pilot study. <i>BMC Urology</i> , <b>2015</b> , 15, 65	2.2	16
32	Appropriateness of oral drugs for long-term treatment of lower urinary tract symptoms in older persons: results of a systematic literature review and international consensus validation process (LUTS-FORTA 2014). <i>Age and Ageing</i> , <b>2015</b> , 44, 745-55	3	71
31	Long-term safety and efficacy of single-tablet combinations of solifenacin and tamsulosin oral controlled absorption system in men with storage and voiding lower urinary tract symptoms: results from the NEPTUNE Study and NEPTUNE II open-label extension. <i>European Urology</i> , <b>2015</b> , 67, 262-70	10.2	47
30	Impact of cardiovascular risk factors and related comorbid conditions and medical therapy reported at baseline on the treatment response to tadalafil 5 mg once-daily in men with lower urinary tract symptoms associated with benign prostatic hyperplasia: an integrated analysis of four randomised, double-blind, placebo-controlled clinical studies. <i>Urology</i> , <b>2015</b> , 85, 1486-507	2.9	11
29	Factors associated with dose escalation of fesoterodine for treatment of overactive bladder in people >65 years of age: A post hoc analysis of data from the SOFIA study. <i>Neurourology and Urodynamics</i> , <b>2015</b> , 34, 438-43	2.3	9
28	EAU Guidelines on the Assessment of Non-neurogenic Male Lower Urinary Tract Symptoms including Benign Prostatic Obstruction. <i>European Urology</i> , <b>2015</b> , 67, 1099-1109	10.2	541
27	Time to onset of clinically meaningful improvement with tadalafil 5 mg once daily for lower urinary tract symptoms secondary to benign prostatic hyperplasia: analysis of data pooled from 4 pivotal, double-blind, placebo controlled studies. <i>Journal of Urology</i> , <b>2015</b> , 193, 1581-9	2.5	26
26	Effects of tadalafil on nighttime voiding (nocturia) in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: a post hoc analysis of pooled data from four randomized, placebo-controlled clinical studies. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1127-32	4	26
25	Cardiovascular and ocular safety of $\alpha$ -adrenoceptor antagonists in the treatment of male lower urinary tract symptoms. <i>Expert Opinion on Drug Safety</i> , <b>2014</b> , 13, 1187-97	4.1	28
24	Nocturia improvement in the combination of Avodart( $\alpha$ ) and tamsulosin (CombAT) study. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1133-40	4	13

23	Impact of dutasteride on nocturia in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia (LUTS/BPH): a pooled analysis of three phase III studies. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1141-7	4	16
22	Fixed-dose combination PRO 160/120 of sabal and urtica extracts improves nocturia in men with LUTS suggestive of BPH: re-evaluation of four controlled clinical studies. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1149-54	4	7
21	Effects of tadalafil once daily on maximum urinary flow rate in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia. <i>Journal of Urology</i> , <b>2014</b> , 191, 1045-50	2.5	26
20	Phosphodiesterase isoenzymes in the human urethra: a molecular biology and functional study. <i>European Journal of Pharmacology</i> , <b>2014</b> , 741, 330-5	5.3	6
19	Detrusor contraction power parameters (BCI and W max) rise with increasing bladder outlet obstruction grade in men with lower urinary tract symptoms: results from a urodynamic database analysis. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1177-83	4	29
18	Treatment satisfaction with tadalafil or tamsulosin vs placebo in men with lower urinary tract symptoms (LUTS) suggestive of benign prostatic hyperplasia (BPH): results from a randomised, placebo-controlled study. <i>BJU International</i> , <b>2014</b> , 114, 568-75	5.6	25
17	Open Simple Prostatectomy <b>2014</b> , 164-174		
16	Nocturia and its impact on health-related quality of life and health care seeking behaviour in German community-dwelling men aged 50 years or older. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1155-62	4	23
15	Review on ultrasound measurement of bladder or detrusor wall thickness in women: techniques, diagnostic utility, and use in clinical trials. <i>World Journal of Urology</i> , <b>2013</b> , 31, 1093-104	4	26
14	Re: Editorial comment on monotherapy with tadalafil or tamsulosin similarly improved lower urinary tract symptoms suggestive of benign prostatic hyperplasia in an international, randomised, parallel, placebo-controlled clinical trial (Eur Urol 2012; 61: 917-925): S. A. Kaplan J Urol 2013; 189: 1448-1449. <i>Journal of Urology</i> , <b>2013</b> , 190, 2311-2	2.5	
13	EAU guidelines on the treatment and follow-up of non-neurogenic male lower urinary tract symptoms including benign prostatic obstruction. <i>European Urology</i> , <b>2013</b> , 64, 118-40	10.2	808
12	The use of a single daily dose of tadalafil to treat signs and symptoms of benign prostatic hyperplasia and erectile dysfunction. <i>Research and Reports in Urology</i> , <b>2013</b> , 5, 99-111	1.3	15
11	Re: Dutasteride improves outcomes of benign prostatic hyperplasia when evaluated for prostate cancer risk reduction: secondary analysis of the reduction by dutasteride of prostate cancer events trial. <i>European Urology</i> , <b>2012</b> , 61, 624-5	10.2	
10	Monotherapy with tadalafil or tamsulosin similarly improved lower urinary tract symptoms suggestive of benign prostatic hyperplasia in an international, randomised, parallel, placebo-controlled clinical trial. <i>European Urology</i> , <b>2012</b> , 61, 917-25	10.2	200
9	Diagnosis and medical treatment of lower urinary tract symptoms in adult men: applying specialist guidelines in clinical practice. <i>BJU International</i> , <b>2012</b> , 110, 710-8	5.6	27
8	Can we identify men who will have complications from benign prostatic obstruction (BPO)? ICI-RS 2011. <i>Neurourology and Urodynamics</i> , <b>2012</b> , 31, 322-6	2.3	32
7	Phosphodiesterase (PDE) inhibitors in the treatment of lower urinary tract dysfunction. <i>British Journal of Clinical Pharmacology</i> , <b>2011</b> , 72, 197-204	3.8	43
6	Age-stratified normal values for prostate volume, PSA, maximum urinary flow rate, IPSS, and other LUTS/BPH indicators in the German male community-dwelling population aged 50 years or older. <i>World Journal of Urology</i> , <b>2011</b> , 29, 171-8	4	61

5	Efficacy of propiverine ER with or without $\beta$ blockers related to maximum urinary flow rate in adult men with OAB: results of a 12-week, multicenter, non-interventional study. <i>World Journal of Urology</i> , <b>2011</b> , 29, 217-23	4	17
4	Manual versus automatic bladder wall thickness measurements: a method comparison study. <i>World Journal of Urology</i> , <b>2009</b> , 27, 747-53	4	13
3	Age and bladder outlet obstruction are independently associated with detrusor overactivity in patients with benign prostatic hyperplasia. <i>European Urology</i> , <b>2008</b> , 54, 419-26	10.2	129
2	Increase in detrusor wall thickness indicates bladder outlet obstruction (BOO) in men. <i>World Journal of Urology</i> , <b>2002</b> , 19, 443-52	4	117
1	Petroleum jelly is an ideal contact medium for pain reduction and successful treatment with extracorporeal shock wave lithotripsy. <i>Journal of Urology</i> , <b>1999</b> , 162, 18-22	2.5	15