

Keiji Miyazawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6145236/publications.pdf>

Version: 2024-02-01

95
papers

8,075
citations

57719

44
h-index

46771

89
g-index

98
all docs

98
docs citations

98
times ranked

9589
citing authors

#	ARTICLE	IF	CITATIONS
1	MAB21L4 regulates the TGF- β ² -induced expression of target genes in epidermal keratinocytes. <i>Journal of Biochemistry</i> , 2022, 171, 399-410.	0.9	3
2	Ets family proteins regulate the EMT transcription factors Snail and ZEB in cancer cells. <i>FEBS Open Bio</i> , 2022, 12, 1353-1364.	1.0	5
3	Neurotensin receptor 1 signaling promotes pancreatic cancer progression. <i>Molecular Oncology</i> , 2021, 15, 151-166.	2.1	17
4	ZEB1 and oncogenic Ras constitute a regulatory switch for stimulus-dependent E-cadherin downregulation. <i>Cancer Science</i> , 2021, 112, 205-216.	1.7	6
5	Group V Secretory Phospholipase A ₂ Regulates Endocytosis of Acetylated LDL by Transcriptional Activation of PK1 in RAW264.7 Macrophage Cell Line. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, , .	0.9	2
6	TGF- β ² -induced cell motility requires downregulation of ARHGAPs to sustain Rac1 activity. <i>Journal of Biological Chemistry</i> , 2021, 296, 100545.	1.6	5
7	EHF suppresses cancer progression by inhibiting ETS1-mediated ZEB expression. <i>Oncogenesis</i> , 2021, 10, 26.	2.1	22
8	Structural basis for inhibitory effects of Smad7 on TGF- β ² family signaling. <i>Journal of Structural Biology</i> , 2020, 212, 107661.	1.3	14
9	Dissociation of the AhR/ARNT complex by TGF- β ² /Smad signaling represses CYP1A1 gene expression and inhibits benze[a]pyrene-mediated cytotoxicity. <i>Journal of Biological Chemistry</i> , 2020, 295, 9033-9051.	1.6	21
10	Deubiquitylase USP25 prevents degradation of BCR-ABL protein and ensures proliferation of Ph-positive leukemia cells. <i>Oncogene</i> , 2020, 39, 3867-3878.	2.6	25
11	Lipidome-based rapid diagnosis with machine learning for detection of TGF- β ² signalling activated area in head and neck cancer. <i>British Journal of Cancer</i> , 2020, 122, 995-1004.	2.9	9
12	Addiction of mesenchymal phenotypes on the FGF/FGFR axis in oral squamous cell carcinoma cells. <i>PLoS ONE</i> , 2019, 14, e0217451.	1.1	12
13	A comparative analysis of Smad-responsive motifs identifies multiple regulatory inputs for TGF- β ² transcriptional activation. <i>Journal of Biological Chemistry</i> , 2019, 294, 15466-15479.	1.6	18
14	Long Noncoding RNA <i>ELIT-1</i> Acts as a Smad3 Cofactor to Facilitate TGF- β ² /Smad Signaling and Promote Epithelial-Mesenchymal Transition. <i>Cancer Research</i> , 2019, 79, 2821-2838.	0.4	84
15	Transgenic Analyses in <i>Drosophila</i> Reveal That mCORL1 Is Functionally Distinct from mCORL2 and dCORL. <i>G3: Genes, Genomes, Genetics</i> , 2019, 9, 3781-3789.	0.8	4
16	Smad3–STAT3 crosstalk in pathophysiological contexts. <i>Acta Biochimica Et Biophysica Sinica</i> , 2018, 50, 82-90.	0.9	57
17	Reciprocal expression of Slug and Snail in human oral cancer cells. <i>PLoS ONE</i> , 2018, 13, e0199442.	1.1	52
18	Expression of ZEBs in gliomas is associated with invasive properties and histopathological grade. <i>Oncology Letters</i> , 2018, 16, 1758-1764.	0.8	9

#	ARTICLE	IF	CITATIONS
19	Repression of Smad3 by Stat3 and c-Ski/SnoN induces gefitinib resistance in lung adenocarcinoma. <i>Biochemical and Biophysical Research Communications</i> , 2017, 484, 269-277.	1.0	18
20	Ets1 and ESE1 reciprocally regulate expression of ZEB1/ZEB2, dependent on ERK1/2 activity, in breast cancer cells. <i>Cancer Science</i> , 2017, 108, 952-960.	1.7	42
21	Regulation of TGF- β Family Signaling by Inhibitory Smads. <i>Cold Spring Harbor Perspectives in Biology</i> , 2017, 9, a022095.	2.3	327
22	Snail suppresses cellular senescence and promotes fibroblast-mediated cancer cell invasion. <i>FEBS Open Bio</i> , 2017, 7, 1586-1597.	1.0	17
23	Dual Roles for Epithelial Splicing Regulatory Proteins 1 (ESRP1) and 2 (ESRP2) in Cancer Progression. <i>Advances in Experimental Medicine and Biology</i> , 2016, 925, 33-40.	0.8	26
24	The potential role of regulator of G-protein signaling 16 in cell motility mediated by EF 1 family proteins. <i>FEBS Letters</i> , 2016, 590, 270-278.	1.3	10
25	EF1 associates with DNMT1 and maintains DNA methylation of the E-cadherin promoter in breast cancer cells. <i>Cancer Medicine</i> , 2015, 4, 125-135.	1.3	83
26	Maid is a negative regulator of transforming growth factor- β -induced cell migration. <i>Journal of Biochemistry</i> , 2015, 158, 435-444.	0.9	7
27	Arkadia enhances BMP signalling through ubiquitylation and degradation of Smad6. <i>Journal of Biochemistry</i> , 2015, 158, 61-71.	0.9	15
28	Epithelial Splicing Regulatory Proteins 1 (ESRP1) and 2 (ESRP2) Suppress Cancer Cell Motility via Different Mechanisms. <i>Journal of Biological Chemistry</i> , 2014, 289, 27386-27399.	1.6	133
29	A Smad3 and TTF-1/NKX2-1 complex regulates Smad4-independent gene expression. <i>Cell Research</i> , 2014, 24, 994-1008.	5.7	45
30	Phosphoinositide 5-phosphatases: how do they affect tumourigenesis?. <i>Journal of Biochemistry</i> , 2013, 153, 1-3.	0.9	4
31	Identification of integrin α 3 as a molecular marker of cells undergoing epithelial-mesenchymal transition and of cancer cells with aggressive phenotypes. <i>Cancer Science</i> , 2013, 104, 1189-1197.	1.7	45
32	Oligodendrocyte Transcription Factor 1 (Olig1) Is a Smad Cofactor Involved in Cell Motility Induced by Transforming Growth Factor- β . <i>Journal of Biological Chemistry</i> , 2013, 288, 18911-18922.	1.6	19
33	Basolateral BMP Signaling in Polarized Epithelial Cells. <i>PLoS ONE</i> , 2013, 8, e62659.	1.1	22
34	A negative regulator or just an unconcerned passerby: phosphoinositide 3-kinase signalling in IL-12 production. <i>Journal of Biochemistry</i> , 2012, 152, 497-499.	0.9	3
35	Transcriptional and Post-transcriptional Regulation in TGF- β -mediated epithelial-mesenchymal transition. <i>Journal of Biochemistry</i> , 2012, 151, 563-571.	0.9	49
36	TGF- β regulates isoform switching of FGF receptors and epithelial-mesenchymal transition. <i>EMBO Journal</i> , 2011, 30, 783-795.	3.5	205

#	ARTICLE	IF	CITATIONS
37	Glioma-initiating Cells Retain Their Tumorigenicity through Integration of the Sox Axis and Oct4 Protein. <i>Journal of Biological Chemistry</i> , 2011, 286, 41434-41441.	1.6	129
38	RB1CC1 Protein Positively Regulates Transforming Growth Factor- β Signaling through the Modulation of Arkadia E3 Ubiquitin Ligase Activity. <i>Journal of Biological Chemistry</i> , 2011, 286, 32502-32512.	1.6	30
39	Encountering unpredicted off-target effects of pharmacological inhibitors. <i>Journal of Biochemistry</i> , 2011, 150, 1-3.	0.9	14
40	Hepatocyte growth factor activator (HGFA): a serine protease that links tissue injury to activation of hepatocyte growth factor. <i>FEBS Journal</i> , 2010, 277, 2208-2214.	2.2	61
41	Arkadia complexes with clathrin adaptor AP2 and regulates EGF signalling. <i>Journal of Biochemistry</i> , 2010, 148, 733-741.	0.9	15
42	Context-dependent regulation of the expression of c-Ski protein by Arkadia in human cancer cells. <i>Journal of Biochemistry</i> , 2010, 147, 545-554.	0.9	29
43	Smad7 Inhibits Transforming Growth Factor- β Family Type I Receptors through Two Distinct Modes of Interaction. <i>Journal of Biological Chemistry</i> , 2010, 285, 30804-30813.	1.6	51
44	SKI and MEL1 Cooperate to Inhibit Transforming Growth Factor- β Signal in Gastric Cancer Cells. <i>Journal of Biological Chemistry</i> , 2009, 284, 3334-3344.	1.6	74
45	VEGFR2-PLC β 3 axis is essential for endothelial specification of VEGFR2+ vascular progenitor cells. <i>Journal of Cell Science</i> , 2009, 122, 3303-3311.	1.2	39
46	Arkadia represses the expression of myoblast differentiation markers through degradation of Ski and the Ski-bound Smad complex in C2C12 myoblasts. <i>Bone</i> , 2009, 44, 53-60.	1.4	13
47	Autocrine TGF- β Signaling Maintains Tumorigenicity of Glioma-Initiating Cells through Sry-Related HMG-Box Factors. <i>Cell Stem Cell</i> , 2009, 5, 504-514.	5.2	503
48	Chromatin Immunoprecipitation on Microarray Analysis of Smad2/3 Binding Sites Reveals Roles of ETS1 and TFAP2A in Transforming Growth Factor β Signaling. <i>Molecular and Cellular Biology</i> , 2009, 29, 172-186.	1.1	179
49	An Id-like molecule, HHM, is a synexpression group-restricted regulator of TGF- β signalling. <i>EMBO Journal</i> , 2008, 27, 2955-2965.	3.5	51
50	Activation of MET receptor tyrosine kinase in ulcer surface epithelial cells undergoing restitution. <i>Pathology International</i> , 2008, 58, 462-464.	0.6	7
51	Ras signaling directs endothelial specification of VEGFR2+ vascular progenitor cells. <i>Journal of Cell Biology</i> , 2008, 181, 131-141.	2.3	42
52	Smurf2 Induces Ubiquitin-dependent Degradation of Smurf1 to Prevent Migration of Breast Cancer Cells. <i>Journal of Biological Chemistry</i> , 2008, 283, 35660-35667.	1.6	73
53	Arkadia Induces Degradation of SnoN and c-Ski to Enhance Transforming Growth Factor- β Signaling. <i>Journal of Biological Chemistry</i> , 2007, 282, 20492-20501.	1.6	148
54	17 β -Estradiol Induces IL-1 β Gene Expression in Rheumatoid Fibroblast-Like Synovial Cells through Estrogen Receptor β (ER β) and Augmentation of Transcriptional Activity of Sp1 by Dissociating Histone Deacetylase 2 from ER β . <i>Journal of Immunology</i> , 2007, 178, 3059-3066.	0.4	39

#	ARTICLE	IF	CITATIONS
55	Transforming Growth Factor- β Promotes Survival of Mammary Carcinoma Cells through Induction of Antiapoptotic Transcription Factor DEC1. <i>Cancer Research</i> , 2007, 67, 9694-9703.	0.4	90
56	Inhibitory effect of c-Met mutants on the formation of branching tubules by a porcine aortic endothelial cell line. <i>Cancer Science</i> , 2006, 97, 1343-1350.	1.7	5
57	Nuclear and cytoplasmic c-Ski differently modulate cellular functions. <i>Genes To Cells</i> , 2006, 11, 1267-1280.	0.5	35
58	NEDD4-2 (neural precursor cell expressed, developmentally down-regulated 4-2) negatively regulates TGF- β (transforming growth factor- β) signalling by inducing ubiquitin-mediated degradation of Smad2 and TGF- β type I receptor. <i>Biochemical Journal</i> , 2005, 386, 461-470.	1.7	187
59	VEGF-A and FGF-2 synergistically promote neoangiogenesis through enhancement of endogenous PDGF- β PDGFR β signaling. <i>Journal of Cell Science</i> , 2005, 118, 3759-3768.	1.2	263
60	Hepatocyte Growth Factor Activator Inhibitor Type 1 (HAI-1) Is Required for Branching Morphogenesis in the Chorioallantoic Placenta. <i>Molecular and Cellular Biology</i> , 2005, 25, 5687-5698.	1.1	89
61	Interaction with Smad4 Is Indispensable for Suppression of BMP Signaling by c-Ski. <i>Molecular Biology of the Cell</i> , 2004, 15, 963-972.	0.9	59
62	c-Ski inhibits the TGF- β signaling pathway through stabilization of inactive Smad complexes on Smad-binding elements. <i>Oncogene</i> , 2004, 23, 5068-5076.	2.6	114
63	Negative regulation of transforming growth factor- β (TGF- β) signaling by WW domain-containing protein 1 (WWP1). <i>Oncogene</i> , 2004, 23, 6914-6923.	2.6	176
64	Activation of hepatocyte growth factor in monkey stomach following gastric mucosal injury. <i>Journal of Gastroenterology</i> , 2004, 39, 133-139.	2.3	6
65	Regeneration of injured intestinal mucosa is impaired in hepatocyte growth factor activator-deficient mice. <i>Gastroenterology</i> , 2004, 127, 1423-1435.	0.6	69
66	Arkadia amplifies TGF- β superfamily signalling through degradation of Smad7. <i>EMBO Journal</i> , 2003, 22, 6458-6470.	3.5	195
67	TGF- β receptor kinase inhibitor enhances growth and integrity of embryonic stem cell-derived endothelial cells. <i>Journal of Cell Biology</i> , 2003, 163, 1303-1311.	2.3	172
68	Two Short Segments of Smad3 Are Important for Specific Interaction of Smad3 with c-Ski and SnoN. <i>Journal of Biological Chemistry</i> , 2003, 278, 531-536.	1.6	51
69	Structural Insight into Modest Binding of a Non-PXXP Ligand to the Signal Transducing Adaptor Molecule-2 Src Homology 3 Domain. <i>Journal of Biological Chemistry</i> , 2003, 278, 48162-48168.	1.6	47
70	SB-431542 and Gleevec inhibit transforming growth factor- β -induced proliferation of human osteosarcoma cells. <i>Cancer Research</i> , 2003, 63, 7791-8.	0.4	103
71	Functional Characterization of Kunitz Domains in Hepatocyte Growth Factor Activator Inhibitor Type 1. <i>Journal of Biological Chemistry</i> , 2002, 277, 14053-14059.	1.6	61
72	Impairment of Activation of Hepatocyte Growth Factor Precursor into Its Mature Form in Rats with Liver Cirrhosis. <i>Journal of Surgical Research</i> , 2002, 106, 108-114.	0.8	19

#	ARTICLE	IF	CITATIONS
73	Exogenously Administered HGF Activator Augments Liver Regeneration through the Production of Biologically Active HGF. <i>Biochemical and Biophysical Research Communications</i> , 2002, 290, 475-481.	1.0	25
74	Targets of transcriptional regulation by two distinct type I receptors for transforming growth factor- β in human umbilical vein endothelial cells. <i>Journal of Cellular Physiology</i> , 2002, 193, 299-318.	2.0	204
75	Two major Smad pathways in TGF- β 2 superfamily signalling. <i>Genes To Cells</i> , 2002, 7, 1191-1204.	0.5	594
76	High Intensity ERK Signal Mediates Hepatocyte Growth Factor-induced Proliferation Inhibition of the Human Hepatocellular Carcinoma Cell Line HepG2. <i>Journal of Biological Chemistry</i> , 2001, 276, 40968-40976.	1.6	81
77	Mouse hepatocyte growth factor activator gene: its expression not only in the liver but also in the gastrointestinal tract. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2000, 1491, 295-302.	2.4	30
78	A Deubiquitinating Enzyme UBPY Interacts with the Src Homology 3 Domain of Hrs-binding Protein via a Novel Binding Motif PX(V/I)(D/N)RXKP. <i>Journal of Biological Chemistry</i> , 2000, 275, 37481-37487.	1.6	199
79	Hepatocyte Growth Factor Activator Inhibitor Type 1 Is a Specific Cell Surface Binding Protein of Hepatocyte Growth Factor Activator (HGFA) and Regulates HGFA Activity in the Pericellular Microenvironment. <i>Journal of Biological Chemistry</i> , 2000, 275, 40453-40462.	1.6	114
80	Structural organization and chromosomal localization of the human hepatocyte growth factor activator gene. Phylogenetic and functional relationship with blood coagulation factor XII, urokinase, and tissue-type plasminogen activator. <i>FEBS Journal</i> , 1998, 258, 355-361.	0.2	31
81	Hepatocyte Growth Factor Activator Inhibitor, a Novel Kunitz-type Serine Protease Inhibitor. <i>Journal of Biological Chemistry</i> , 1997, 272, 6370-6376.	1.6	241
82	Purification and Characterization of a Novel Protease Inhibitor Specific to Hepatocyte Growth Factor Activator. , 1997, , 403-407.		0
83	Dissociation of C-fos Induction and Mitogen-Activated-Protein Kinase Activation from the Hepatocyte-Growth-Factor-Induced Motility Response in Human Gastric Carcinoma Cells. <i>FEBS Journal</i> , 1996, 236, 476-481.	0.2	15
84	Activation of Hepatocyte Growth Factor in the Injured Tissues Is Mediated by Hepatocyte Growth Factor Activator. <i>Journal of Biological Chemistry</i> , 1996, 271, 3615-3618.	1.6	199
85	Hepatocyte growth factor remains as an inactive single chain after partial hepatectomy or unilateral nephrectomy. <i>FEBS Letters</i> , 1995, 362, 220-224.	1.3	20
86	Activation of Hepatocyte Growth Factor by two Homologous Proteases, Blood-Coagulation Factor XIIa and Hepatocyte Growth Factor Activator. <i>FEBS Journal</i> , 1995, 229, 257-261.	0.2	135
87	Tyrosine Phosphorylation of β -Catenin and Plakoglobin Enhanced by Hepatocyte Growth Factor and Epidermal Growth Factor in Human Carcinoma Cells. <i>Cell Adhesion and Communication</i> , 1994, 1, 295-305.	1.7	402
88	Characterization of the promoter region of the rat hepatocyte-growth-factor/scatter-factor gene. <i>FEBS Journal</i> , 1993, 213, 113-119.	0.2	57
89	Stimulation of prostaglandin production by hepatocyte growth factor in human gastric carcinoma cells. <i>FEBS Letters</i> , 1993, 334, 331-334.	1.3	13
90	Functional characterization of human hepatocyte growth factor mutants obtained by deletion of structural domains. <i>Biochemistry</i> , 1992, 31, 9555-9561.	1.2	72

#	ARTICLE	IF	CITATIONS
91	Characterization of hepatocyte-growth-factor receptors on Meth A cells. FEBS Journal, 1992, 204, 857-864.	0.2	40
92	Hepatocyte Growth Factor and Transforming Growth Factor-.BETA. Stimulate both Cell Growth and Migration of Human Gastric Adenocarcinoma Cells.. Cell Structure and Function, 1992, 17, 185-190.	0.5	38
93	An alternatively processed mRNA generated from human hepatocyte growth factor gene. FEBS Journal, 1991, 197, 15-22.	0.2	100
94	Primary structure of rat hepatocyte growth factor and induction of its mRNA during liver regeneration following hepatic injury. FEBS Journal, 1990, 193, 375-381.	0.2	108
95	Molecular cloning and sequence analysis of cDNA for human hepatocyte growth factor. Biochemical and Biophysical Research Communications, 1989, 163, 967-973.	1.0	670