José MarÃ-a González Molinillo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6149630/publications.pdf

Version: 2024-02-01

150 4,957 40
papers citations h-index

63 g-index

161 161 all docs citations

161 times ranked

3725 citing authors

#	Article	IF	CITATIONS
1	Strategies for the synthesis of canonical, non-canonical and analogues of strigolactones, and evaluation of their parasitic weed germination activity. Phytochemistry Reviews, 2022, 21, 1627-1659.	6.5	14
2	Encapsulation of Cynara Cardunculus Guaiane-type Lactones in Fully Organic Nanotubes Enhances Their Phytotoxic Properties. Journal of Agricultural and Food Chemistry, 2022, 70, 3644-3653.	5.2	7
3	Evaluation of the phytotoxic and antifungal activity of <scp>C₁₇</scp> â€sesquiterpenoids as potential biopesticides. Pest Management Science, 2022, 78, 4240-4251.	3.4	2
4	Acyl Derivatives of Eudesmanolides To Boost their Bioactivity: An Explanation of Behavior in the Cell Membrane Using a Molecular Dynamics Approach. ChemMedChem, 2021, 16, 1297-1307.	3.2	7
5	Sesquiterpenes in Fresh Food., 2021,, 477-542.		3
6	One-Step Encapsulation of <i>ortho</i> -Disulfides in Functionalized Zinc MOF. Enabling Metalâ€"Organic Frameworks in Agriculture. ACS Applied Materials & Samp; Interfaces, 2021, 13, 7997-8005.	8.0	14
7	Absorption and Elimination of the Allelochemical MBOA by Weeds during Seedling Growth. Agronomy, 2021, 11, 471.	3.0	4
8	Sunflower Metabolites Involved in Resistance Mechanisms against Broomrape. Agronomy, 2021, 11, 501.	3.0	6
9	An Overview of the Chemical Characteristics, Bioactivity and Achievements Regarding the Therapeutic Usage of Acetogenins from Annona cherimola Mill Molecules, 2021, 26, 2926.	3.8	15
10	Pharmacological Activities of Aminophenoxazinones. Molecules, 2021, 26, 3453.	3.8	8
11	A Study on the Phytotoxic Potential of the Seasoning Herb Marjoram (Origanum majorana L.) Leaves. Molecules, 2021, 26, 3356.	3.8	17
12	Search of New Tools for Weed Control Using <i>Piptocarpha rotundifolia</i> , a Dominant Species in the Cerrado. Journal of Agricultural and Food Chemistry, 2021, 69, 8684-8694.	5.2	2
13	Bioactive Diterpenes from the Brazilian Native Plant (Moquiniastrum pulchrum) and Their Application in Weed Control. Molecules, 2021, 26, 4632.	3.8	1
14	Synthesis of Pertyolides A, B, and C: A Synthetic Procedure to C17-Sesquiterpenoids and a Study of Their Phytotoxic Activity. Journal of Natural Products, 2021, 84, 2295-2302.	3.0	6
15	Allelopathic Activity of Strigolactones on the Germination of Parasitic Plants and Arbuscular Mycorrhizal Fungi Growth. Agronomy, 2021, 11, 2174.	3.0	11
16	Phytochemical Study of Safflower Roots (Carthamus tinctorius) on the Induction of Parasitic Plant Germination and Weed Control. Journal of Chemical Ecology, 2020, 46, 871-880.	1.8	13
17	Synthesis of Active Strigolactone Analogues Based on Eudesmane- and Guaiane-Type Sesquiterpene Lactones. Journal of Agricultural and Food Chemistry, 2020, 68, 9636-9645.	5.2	13
18	Effect of Shading on the Sesquiterpene Lactone Content and Phytotoxicity of Cultivated Cardoon Leaf Extracts. Journal of Agricultural and Food Chemistry, 2020, 68, 11946-11953.	5.2	16

#	Article	IF	CITATIONS
19	Study by NMR of Liquid-Phase Alkylation of Toluene with Hex-1-ene: Effect of Catalyst on Selectivity. Petroleum Chemistry, 2020, 60, 810-817.	1.4	1
20	Bio-Guided Isolation of Acetogenins from Annona cherimola Deciduous Leaves: Production of Nanocarriers to Boost the Bioavailability Properties. Molecules, 2020, 25, 4861.	3.8	11
21	Synthesis of Vlasouliolides: A Pathway toward Guaiane–Eudesmane C ₁₇ /C ₁₅ Dimers by Photochemical and Michael Additions. Journal of Organic Chemistry, 2020, 85, 7322-7332.	3.2	4
22	Selective Ring Opening of Ethylbenzene on Bifunctional Catalyst Pt–Ir over Hierarchical USY Zeolite. Petroleum Chemistry, 2020, 60, 104-112.	1.4	1
23	Allelopathic activity of Thapsia garganica L. leaves on lettuce and weeds, and identification of the active principles. South African Journal of Botany, 2020, 131, 188-194.	2.5	10
24	Toxicity and Anti-promastigote Activity of Benzoxazinoid Analogs Against Leishmania (Viannia) braziliensis and Leishmania (Leishmania) infantum. Advanced Pharmaceutical Bulletin, 2020, 10, 119-124.	1.4	1
25	Evaluation of the Phytotoxicity of <i>Urochloa humidicola</i> Roots by Bioassays and Microscopic Analysis. Characterization of New Compounds. Journal of Agricultural and Food Chemistry, 2020, 68, 4851-4864.	5.2	5
26	Allelopathy: The Chemical Language of Plants. Progress in the Chemistry of Organic Natural Products, 2020, 112, 1-84.	1.1	10
27	Quantification of Strigolactones. Methods in Molecular Biology, 2020, 2083, 199-208.	0.9	1
28	Sesquiterpenes in Fresh Food. , 2020, , 1-66.		1
29	In Situ Eco Encapsulation of Bioactive Agrochemicals within Fully Organic Nanotubes. ACS Applied Materials & Dividing Property (1988) amp; Interfaces, 2019, 11, 41925-41934.	8.0	13
30	Preparation and Phytotoxicity Evaluation of 11,13-Dehydro <i>seco</i> -Guaianolides. Journal of Natural Products, 2019, 82, 2501-2508.	3.0	4
31	Easy Access to Alkoxy, Amino, Carbamoyl, Hydroxy, and Thiol Derivatives of Sesquiterpene Lactones and Evaluation of Their Bioactivity on Parasitic Weeds. Journal of Agricultural and Food Chemistry, 2019, 67, 10764-10773.	5.2	16
32	Recent advances in allelopathy for weed control: from knowledge to applications. Pest Management Science, 2019, 75, 2413-2436.	3.4	168
33	Phytotoxicity Study of Ortho-Disubstituted Disulfides and Their Acyl Derivatives. ACS Omega, 2019, 4, 2362-2368.	3.5	13
34	Influence of Genotype and Harvest Time on the <i>Cynara cardunculus</i> L. Sesquiterpene Lactone Profile. Journal of Agricultural and Food Chemistry, 2019, 67, 6487-6496.	5.2	30
35	Facile synthesis of anhydrojudaicin and 11,13-dehydroanhydrojudaicin, two eudesmanolide-skeleton lactones with potential allelopathic activity. Phytochemistry Letters, 2019, 31, 229-236.	1.2	11
36	Structure (i>†(i) activity relationship studies on naphthoquinone analogs. The search for new herbicides based on natural products. Pest Management Science, 2019, 75, 2517-2529.	3.4	11

#	Article	IF	Citations
37	Effect of flavonoids isolated from Tridax procumbens on the growth and toxin production of Microcystis aeruginos. Aquatic Toxicology, 2019, 211, 81-91.	4.0	18
38	Hydrolysable Tannins and Biological Activities of Meriania hernandoi and Meriania nobilis (Melastomataceae). Molecules, 2019, 24, 746.	3.8	7
39	Resistance modulatory and efflux-inhibitory activities of capsaicinoids and capsinoids. Bioorganic Chemistry, 2019, 82, 378-384.	4.1	14
40	The extraction procedure improves the allelopathic activity of cardoon (Cynara cardunculus var.) Tj ETQq0 0 0 rg	gBT/Qverlo	ock 10 Tf 50 6
41	Synthesis and antimicrobial activity of some benzoxazinoids derivatives of 2-nitrophenol and 3-hydroxy-2-nitropyridine. Synthetic Communications, 2019, 49, 286-296.	2.1	8
42	Provitamin supramolecular polymer micelle with pH responsiveness to control release, bioavailability enhancement and potentiation of cytotoxic efficacy. Colloids and Surfaces B: Biointerfaces, 2019, 173, 85-93.	5.0	13
43	A new UHPLCâ€MS/MS method for the direct determination of strigolactones in root exudates and extracts. Phytochemical Analysis, 2019, 30, 110-116.	2.4	26
44	Ecological Relevance of the Major Allelochemicals in <i>Lycopersicon esculentum</i> Roots and Exudates. Journal of Agricultural and Food Chemistry, 2018, 66, 4638-4644.	5.2	25
45	A Novel Electron Microscopic Characterization of Core/Shell Nanobiostimulator Against Parasitic Plants. ACS Applied Materials & Samp; Interfaces, 2018, 10, 2354-2359.	8.0	12
46	Synthesis of $(\hat{A}\pm)$ -3,4-dimethoxybenzyl-4-methyloctanoate as a novel internal standard for capsinoid determination by HPLC-ESI-MS/MS(QTOF). Open Chemistry, 2018, 16, 87-94.	1.9	2
47	Qualitative Study on the Production of the Allelochemicals Benzoxazinones by Inducing Polyploidy in Gramineae with Colchicine. Journal of Agricultural and Food Chemistry, 2018, 66, 3666-3674.	5 . 2	4
48	Influence of lipophilicity in <i>O</i> â€acyl and <i>O</i> â€alkyl derivatives of juglone and lawsone: a structure–activity relationship study in the search for natural herbicide models. Pest Management Science, 2018, 74, 682-694.	3.4	19
49	(+)-epi-Epoformin, a Phytotoxic Fungal Cyclohexenepoxide: Structure Activity Relationships. Molecules, 2018, 23, 1529.	3.8	13
50	Phytotoxic studies of naphthoquinone intermediates from the synthesis of the natural product Naphthotectone. Research on Chemical Intermediates, 2017, 43, 4387-4400.	2.7	8
51	Preparation and phytotoxicity study of lappalone from dehydrocostuslactone. Phytochemistry Letters, 2017, 20, 66-72.	1.2	14
52	Phytotoxicity Study on <i>Bidens sulphurea</i> Sch. Bip. as a Preliminary Approach for Weed Control. Journal of Agricultural and Food Chemistry, 2017, 65, 5161-5172.	5.2	23
53	Complexation of sesquiterpene lactones with cyclodextrins: synthesis and effects on their activities on parasitic weeds. Organic and Biomolecular Chemistry, 2017, 15, 6500-6510.	2.8	23
54	Bioactivity and quantitative analysis of isohexenylnaphthazarins in root periderm of two Echium spp.: E.Âplantagineum and E.Âgaditanum. Phytochemistry, 2017, 141, 162-170.	2.9	13

#	Article	IF	CITATIONS
55	Gibberellic and kaurenoic hybrid strigolactone mimics for seed germination of parasitic weeds. Pest Management Science, 2017, 73, 2529-2537.	3.4	12
56	Chemical evidence for the effect of <i>Urochloa ruziziensis</i> on glyphosateâ€resistant soybeans. Pest Management Science, 2017, 73, 2071-2078.	3.4	13
57	Allelopathy of Bracken Fern (Pteridium arachnoideum): New Evidence from Green Fronds, Litter, and Soil. PLoS ONE, 2016, 11, e0161670.	2.5	28
58	Phthalimideâ€derived strigolactone mimics as germinating agents for seeds of parasitic weeds. Pest Management Science, 2016, 72, 2069-2081.	3.4	21
59	Enantioselective Total Syntheses of (<i>R</i>)―and (<i>S</i>)â€Naphthotectone, and Stereochemical Assignment of the Natural Product. European Journal of Organic Chemistry, 2016, 2016, 1599-1605.	2.4	5
60	Synthesis, antibacterial and antifungal activities of naphthoquinone derivatives: a structure–activity relationship study. Medicinal Chemistry Research, 2016, 25, 1274-1285.	2.4	72
61	Evaluation of the Allelopathic Potential of Leaf, Stem, and Root Extracts of <i>Ocotea pulchella </i> <scp>Nees et Mart</scp> . Chemistry and Biodiversity, 2016, 13, 1058-1067.	2.1	10
62	The Joint Action of Sesquiterpene Lactones from Leaves as an Explanation for the Activity of <i>Cynara cardunculus</i> . Journal of Agricultural and Food Chemistry, 2016, 64, 6416-6424.	5.2	26
63	Phytotoxicity evaluation of sesquiterpene lactones and diterpenes from species of the Decachaeta , Salvia and Podachaenium genera. Phytochemistry Letters, 2016, 18, 68-76.	1.2	24
64	Phytotoxicity of Triterpenes and Limonoids from the Rutaceae and Meliaceae. 5α,6β,8α,12α-Tetrahydro-28-norisotoonafolin – a Potent Phytotoxin from Toona ciliata. Natural Product Communications, 2015, 10, 1934578X1501000.	0.5	5
65	Isolation of Bioactive Compounds from Sunflower Leaves (<i>Helianthus annuus</i> L.) Extracted with Supercritical Carbon Dioxide. Journal of Agricultural and Food Chemistry, 2015, 63, 6410-6421.	5.2	34
66	Soil biodegradation of a benzoxazinone analog proposed as a natural products-based herbicide. Plant and Soil, 2015, 393, 207-214.	3.7	15
67	Phytotoxins from <i>Tithonia diversifolia</i> . Journal of Natural Products, 2015, 78, 1083-1092.	3.0	44
68	Trends in the Synthesis and Functionalization of Guaianolides. European Journal of Organic Chemistry, 2015, 2015, 2093-2110.	2.4	46
69	Helikaurolides A–D with a Diterpene-Sesquiterpene Skeleton from Supercritical Fluid Extracts of <i>Helianthus annuus</i> L. var. Arianna. Organic Letters, 2015, 17, 4730-4733.	4.6	12
70	Phytotoxic Potential of <i>Onopordum acanthium</i> L. (Asteraceae). Chemistry and Biodiversity, 2014, 11, 1247-1255.	2.1	12
71	Bio-guided optimization of the ultrasound-assisted extraction of compounds from Annona glabra L. leaves using the etiolated wheat coleoptile bioassay. Ultrasonics Sonochemistry, 2014, 21, 1578-1584.	8.2	25
72	Phytotoxicity of alkaloids, coumarins and flavonoids isolated from 11 species belonging to the Rutaceae and Meliaceae families. Phytochemistry Letters, 2014, 8, 226-232.	1.2	46

#	Article	IF	Citations
73	Evidence for an Allelopathic Interaction Between Rye and Wild Oats. Journal of Agricultural and Food Chemistry, 2014, 62, 9450-9457.	5.2	52
74	Synthesis of Bioactive Speciosins G and P fromHexagonia speciosa. Journal of Natural Products, 2014, 77, 2029-2036.	3.0	9
75	Phytotoxicity of Cardoon (<i>Cynara cardunculus</i>) Allelochemicals on Standard Target Species and Weeds. Journal of Agricultural and Food Chemistry, 2014, 62, 6699-6706.	5 . 2	58
76	Benzoxazinoids in Rye Allelopathy - From Discovery to Application in Sustainable Weed Control and Organic Farming. Journal of Chemical Ecology, 2013, 39, 154-174.	1.8	154
77	Practical First Total Synthesis of the Potent Phytotoxic (±)â€Naphthotectone, Isolated from <i>Tectona grandis</i> . European Journal of Organic Chemistry, 2013, 2013, 6175-6180.	2.4	13
78	Allelopathic Potential of <i>Rapanea umbellata</i> Leaf Extracts. Chemistry and Biodiversity, 2013, 10, 1539-1548.	2.1	7
79	Ecological phytochemistry of Cerrado (Brazilian savanna) plants. Phytochemistry Reviews, 2013, 12, 839-855.	6.5	28
80	Guaianolides for Multipurpose Molecular Design. ACS Symposium Series, 2013, , 167-188.	0.5	14
81	Phytotoxic effect of bioactive compounds isolated from Myrcia tomentosa (Myrtaceae) leaves. Biochemical Systematics and Ecology, 2013, 46, 29-35.	1.3	31
82	Soy isoflavones and their relationship with microflora: beneficial effects on human health in equol producers. Phytochemistry Reviews, 2013, 12, 979-1000.	6.5	47
83	Preface: special issue Biocom12. Phytochemistry Reviews, 2013, 12, 579-580.	6.5	O
84	Facile Preparation of Bioactive <i>seco</i> -Guaianolides and Guaianolides from <i>Artemisia gorgonum</i> and Evaluation of Their Phytotoxicity. Journal of Natural Products, 2012, 75, 1967-1973.	3.0	20
85	SAR studies of epoxycurcuphenol derivatives on leukemia CT-CD4 cells. Bioorganic and Medicinal Chemistry, 2012, 20, 6662-6668.	3.0	2
86	Estudo fitoquÃmico de folhas de Solanum lycocarpum A. StHil (Solanaceae) e sua aplicação na alelopatia. Acta Botanica Brasilica, 2012, 26, 607-618.	0.8	15
87	Tectonoelins, new norlignans from a bioactive extract of Tectona grandis. Phytochemistry Letters, 2012, 5, 382-386.	1.2	23
88	Potential allelopathic of the fractions obtained from sunflower leaves using supercritical carbon dioxide. Journal of Supercritical Fluids, 2011, 60, 28-37.	3.2	14
89	Anthratectone and Naphthotectone, Two Quinones from Bioactive Extracts of Tectona grandis. Journal of Chemical Ecology, 2011, 37, 1341-1348.	1.8	30
90	Biotransformation of ethyl 2-(2′-nitrophenoxy)acetate to benzohydroxamic acid (D-DIBOA) by Escherichia coli. Process Biochemistry, 2011, 46, 358-364.	3.7	7

#	Article	IF	Citations
91	Isolation and Phytotoxicity of Terpenes from Tectona grandis. Journal of Chemical Ecology, 2010, 36, 396-404.	1.8	59
92	Multifunctionalised benzoxazinones in the systems Oryza sativa-Echinochloa crus-galli and Triticum aestivum-Avena fatua as natural-product-based herbicide leads. Pest Management Science, 2010, 66, 1137-1147.	3.4	4
93	Metabolites from <i>Withania aristata </i> with Potential Phytotoxic Activity. Natural Product Communications, 2010, 5, 1934578X1000500.	0.5	6
94	Application of Hansch's Model to Capsaicinoids and Capsinoids: A Study Using the Quantitative Structureâ^'Activity Relationship. A Novel Method for the Synthesis of Capsinoids. Journal of Agricultural and Food Chemistry, 2010, 58, 3342-3349.	5.2	57
95	Combined Strategy for Phytotoxicity Enhancement of Benzoxazinones. Journal of Agricultural and Food Chemistry, 2010, 58, 2047-2053.	5.2	18
96	Aneugenic effects of benzoxazinones in cultured human cells. Mutation Research - Genetic Toxicology and Environmental Mutagenesis, 2010, 695, 81-86.	1.7	10
97	Structure–activity relationship of benzoxazinones and related compounds with respect to the growth inhibition and α-amylase activity in cress seedlings. Journal of Plant Physiology, 2010, 167, 1221-1225.	3.5	16
98	Metabolites from Withania aristata with potential phytotoxic activity. Natural Product Communications, 2010, 5, 1043-7.	0.5	5
99	Exudados de la raiz y su relevancia actual en las interacciones alelopaticas. Quimica Nova, 2009, 32, 198-213.	0.3	8
100	Aromaticâ€ringâ€functionalised benzoxazinones in the system <i>Oryza sativa–Echinochloa crusâ€galli</i> as biorational herbicide models. Pest Management Science, 2009, 65, 1104-1113.	3.4	9
101	Rediscovering the bioactivity and ecological role of 1,4-benzoxazinones. Natural Product Reports, 2009, 26, 478.	10.3	106
102	Sesquiterpenes as Immunosuppressants. Transplantation, 2009, 88, S24-S30.	1.0	4
103	Helikauranoside A, a New Bioactive Diterpene. Journal of Chemical Ecology, 2008, 34, 65-69.	1.8	30
104	Bioactive apocarotenoids from Tectona grandis. Phytochemistry, 2008, 69, 2708-2715.	2.9	55
105	A stereoselective route towards heliannuol A. Tetrahedron, 2008, 64, 5502-5508.	1.9	20
106	Modified Benzoxazinones in the System <i>Oryza sativa</i> °' <i>Echinochloa crus-galli</i> : An Approach to the Development of Biorational Herbicide Models. Journal of Agricultural and Food Chemistry, 2008, 56, 9941-9948.	5. 2	7
107	Allelopathyâ€"a natural alternative for weed control. Pest Management Science, 2007, 63, 327-348.	3.4	354
108	Plant biocommunicators: their phytotoxicity, degradation studies and potential use as herbicide models. Phytochemistry Reviews, 2007, 7, 179-194.	6.5	62

#	Article	IF	CITATIONS
109	Sesquiterpene Lactones as Allelochemicals. Journal of Natural Products, 2006, 69, 795-800.	3.0	40
110	Heliespirones B and C:  Two New Plant Heliespiranes with a Novel Spiro Heterocyclic Sesquiterpene Skeleton. Organic Letters, 2006, 8, 4513-4516.	4.6	51
111	Structureâ [^] Activity Relationship (SAR) Studies of Benzoxazinones, Their Degradation Products, and Analogues. Phytotoxicity on Problematic WeedsAvena fatuaL. andLolium rigidumGaud Journal of Agricultural and Food Chemistry, 2006, 54, 1040-1048.	5.2	65
112	New Herbicide Models from Benzoxazinones:Â Aromatic Ring Functionalization Effects. Journal of Agricultural and Food Chemistry, 2006, 54, 9843-9851.	5.2	26
113	Bioactive steroids from Oryza sativa L Steroids, 2006, 71, 603-608.	1.8	65
114	11,16 Oxetane lactones. Spectroscopic evidences and conformational analysis. Tetrahedron, 2006, 62, 7747-7755.	1.9	19
115	Optimization of Benzoxazinones as Natural Herbicide Models by Lipophilicity Enhancement. Journal of Agricultural and Food Chemistry, 2006, 54, 9357-9365.	5.2	42
116	Isolation and Synthesis of Allelochemicals from Gramineae:Â Benzoxazinones and Related Compounds. Journal of Agricultural and Food Chemistry, 2006, 54, 991-1000.	5.2	76
117	Degradation Studies on Benzoxazinoids. Soil Degradation Dynamics of (2R)-2-O-Î ² -d-Glucopyranosyl-4-hydroxy-(2H)- 1,4-benzoxazin-3(4H)-one (DIBOA-Glc) and Its Degradation Products, Phytotoxic Allelochemicals from Gramineae. Journal of Agricultural and Food Chemistry, 2005, 53, 554-561.	5.2	92
118	Structureâ^'Activity Relationship Studies of Benzoxazinones and Related Compounds. Phytotoxicity on Echinochloa crus-galli(L.) P. Beauv Journal of Agricultural and Food Chemistry, 2005, 53, 4373-4380.	5.2	28
119	Structureâ^'Activity Relationships (SAR) Studies of Benzoxazinones, Their Degradation Products and Analogues. Phytotoxicity on Standard Target Species (STS). Journal of Agricultural and Food Chemistry, 2005, 53, 538-548.	5.2	99
120	Helivypolide G. A novel dimeric bioactive sesquiterpene lactone. Tetrahedron Letters, 2004, 45, 6567-6570.	1.4	9
121	Bioactive apocarotenoids annuionones F and G: structural revision of annuionones A, B and E. Phytochemistry, 2004, 65, 3057-3063.	2.9	42
122	Degradation Studies on Benzoxazinoids. Soil Degradation Dynamics of 2,4-Dihydroxy-7-methoxy-(2H)-1,4-benzoxazin-3(4H)-one (DIMBOA) and Its Degradation Products, Phytotoxic Allelochemicals from Gramineae. Journal of Agricultural and Food Chemistry, 2004, 52, 6402-6413.	5.2	125
123	Bioactive Lignans from a Cultivar ofHelianthus annuus. Journal of Agricultural and Food Chemistry, 2004, 52, 6443-6447.	5.2	60
124	Synthesis of heliannane skeletons. Facile preparation of $(\hat{A}\pm)$ -heliannuol D. Tetrahedron, 2003, 59, 1679-1683.	1.9	44
125	Allelopathy as a new strategy for sustainable ecosystems development. Uchu Seibutsu Kagaku, 2003, 17, 18-23.	0.3	62
126	Allelochemicals from sunflowers: chemistry, bioactivity and applications., 2002,, 73-87.		19

#	Article	IF	CITATIONS
127	First total synthesis of (±)-helibisabonol A. Tetrahedron Letters, 2002, 43, 6417-6420.	1.4	17
128	Bioactive terpenoids from sunflower leaves cv. Peredovick®. Phytochemistry, 2002, 61, 687-692.	2.9	108
129	The Use of Allelopathic Studies in the Search for Natural Herbicides. The Journal of Crop Improvement: Innovations in Practiceory and Research, 2001, 4, 237-255.	0.4	50
130	Studies on the Stereostructure of Eudesmanolides from Umbelliferae: Total Synthesis of (+)-Decipienin A. Tetrahedron, 2000, 56, 3409-3414.	1.9	22
131	Dehydrozaluzanin C: a potent plant growth regulator with potential use as a natural herbicide template. Phytochemistry, 2000, 54, 165-171.	2.9	53
132	Title is missing!. Journal of Chemical Ecology, 2000, 26, 2173-2186.	1.8	41
133	Search for a Standard Phytotoxic Bioassay for Allelochemicals. Selection of Standard Target Speciesâ€. Journal of Agricultural and Food Chemistry, 2000, 48, 2512-2521.	5.2	242
134	Allelochemicals from sunflower leaves cv. Peredovick. Phytochemistry, 1999, 52, 613-621.	2.9	80
135	Heliannuol E. A novel bioactive sesquiterpene of the heliannane family. Tetrahedron Letters, 1999, 40, 4725-4728.	1.4	61
136	New Bioactive Plant Heliannuols from Cultivar Sunflower Leaves 1. Journal of Natural Products, 1999, 62, 1636-1639.	3.0	76
137	Heliespirone A. The first member of a novel family of bioactive sesquiterpenes. Tetrahedron Letters, 1998, 39, 427-430.	1.4	54
138	Bioactive norsesquiterpenes from Helianthus annuus with potential allelopathic activity. Phytochemistry, 1998, 48, 631-636.	2.9	88
139	Bioactive flavonoids from Helianthus annuus cultivars. Phytochemistry, 1997, 45, 683-687.	2.9	63
140	Potential allelopathic sesquiterpene lactones from sunflower leaves. Phytochemistry, 1996, 43, 1205-1215.	2.9	78
141	Studies on the stereostructure of eudesmanolides from Umbelliferae: synthesis of $11\hat{l}^2$ -angeloyloxy- $\hat{l}\pm$ -santonin. Tetrahedron, 1994, 50, 5439-5450.	1.9	17
142	Structural Elucidation and Chemistry of a Novel Family of Bioactive Sesquiterpenes: Heliannuols. Journal of Organic Chemistry, 1994, 59, 8261-8266.	3.2	148
143	Potential allelopathic guaianolides from cultivar sunflower leaves, var. SH-222. Phytochemistry, 1993, 34, 669-674.	2.9	71
144	First synthesis of two naturally occurring oxetane lactones: clementein and clementein b. Tetrahedron, 1993, 49, 2499-2508.	1.9	20

#	Article	IF	CITATIONS
145	Novel sesquiterpene from bioactive fractions of cultivar sunflowers. Tetrahedron Letters, 1993, 34, 1999-2002.	1.4	96
146	Melampolides and cis,cis-germacranolides from Lecocarpus lecocarpoides. Phytochemistry, 1992, 32, 127-131.	2.9	3
147	Study of photochemical addition of acyl radical to electron-deficient olefins. Tetrahedron, 1992, 48, 3345-3352.	1.9	18
148	"An efficient and mild entry to 1,4-dicarbonyl compounds via photochemical addition of acyl radical to electron-deficient olefins― Tetrahedron Letters, 1990, 31, 3063-3066.	1.4	17
149	Terpene synthesis. 1. Chemical transformation of deacylsubexpinnatin into the natural oxetane lactone subexpinnatin C. Journal of Organic Chemistry, 1987, 52, 3323-3326.	3.2	20
150	Structure-activity relationship study of diterpenes for treatment of Alzheimer's disease. Quimica Nova, 0, , .	0.3	1