Kensei Tobinai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6156495/publications.pdf

Version: 2024-02-01

76326 25787 12,039 130 40 108 citations h-index g-index papers 132 132 132 10486 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Revised Response Criteria for Malignant Lymphoma. Journal of Clinical Oncology, 2007, 25, 579-586.	1.6	4,061
2	Integrated molecular analysis of adult T cell leukemia/lymphoma. Nature Genetics, 2015, 47, 1304-1315.	21.4	659
3	Defucosylated Anti-CCR4 Monoclonal Antibody (KW-0761) for Relapsed Adult T-Cell Leukemia-Lymphoma: A Multicenter Phase II Study. Journal of Clinical Oncology, 2012, 30, 837-842.	1.6	581
4	Brentuximab vedotin with chemotherapy for CD30-positive peripheral T-cell lymphoma (ECHELON-2): a global, double-blind, randomised, phase 3 trial. Lancet, The, 2019, 393, 229-240.	13.7	517
5	Definition, Prognostic Factors, Treatment, and Response Criteria of Adult T-Cell Leukemia-Lymphoma: A Proposal From an International Consensus Meeting. Journal of Clinical Oncology, 2009, 27, 453-459.	1.6	485
6	VCAP-AMP-VECP Compared With Biweekly CHOP for Adult T-Cell Leukemia-Lymphoma: Japan Clinical Oncology Group Study JCOG9801. Journal of Clinical Oncology, 2007, 25, 5458-5464.	1.6	429
7	Phase I Study of KW-0761, a Defucosylated Humanized Anti-CCR4 Antibody, in Relapsed Patients With Adult T-Cell Leukemia-Lymphoma and Peripheral T-Cell Lymphoma. Journal of Clinical Oncology, 2010, 28, 1591-1598.	1.6	351
8	A new Gâ€CSFâ€supported combination chemotherapy, LSG15, for adult Tâ€cell leukaemiaâ€lymphoma: Japan Clinical Oncology Group Study 9303. British Journal of Haematology, 2001, 113, 375-382.	2.5	335
9	Phase I/II Study of Concurrent Chemoradiotherapy for Localized Nasal Natural Killer/T-Cell Lymphoma: Japan Clinical Oncology Group Study JCOG0211. Journal of Clinical Oncology, 2009, 27, 5594-5600.	1.6	315
10	Peripheral T-cell lymphoma. Blood, 2011, 117, 6756-6767.	1.4	278
10	Peripheral T-cell lymphoma. Blood, 2011, 117, 6756-6767. Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase ⟨scp⟩ll⟨/scp⟩ study. British Journal of Haematology, 2015, 169, 672-682.	2.5	278
	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase <scp>II</scp> study. British Journal of		
11	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase <scp>II</scp> study. British Journal of Haematology, 2015, 169, 672-682. Toxicity Grading Criteria of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology,	2.5	218
11 12	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase ⟨scp⟩ll⟨lscp⟩ study. British Journal of Haematology, 2015, 169, 672-682. Toxicity Grading Criteria of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology, 1993, 23, 250-257.	2.5	218
11 12 13	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase ⟨scp⟩II⟨/scp⟩ study. British Journal of Haematology, 2015, 169, 672-682. Toxicity Grading Criteria of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology, 1993, 23, 250-257. Feasibility and pharmacokinetic study of a chimeric anti-CD20 monoclonal antibody (IDEC-C2B8,) Tj ETQq1 1 0.75 Significance of cyclin D1 overexpression for the diagnosis of mantle cell lymphoma: a clinicopathologic comparison of cyclin D1-positive MCL and cyclin D1-negative MCL-like B-cell	2.5 1.3 843] 4 rgE	218 170 BT /Qyerlock
11 12 13	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tâ€cell leukaemiaâ€lymphoma: a randomized phase ⟨scp⟩ll⟨lscp⟩ study. British Journal of Haematology, 2015, 169, 672-682. Toxicity Grading Criteria of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology, 1993, 23, 250-257. Feasibility and pharmacokinetic study of a chimeric anti-CD20 monoclonal antibody (IDEC-C2B8,) Tj ETQq1 1 0.76. Significance of cyclin D1 overexpression for the diagnosis of mantle cell lymphoma: a clinicopathologic comparison of cyclin D1-positive MCL and cyclin D1-negative MCL-like B-cell lymphoma. Blood, 2000, 95, 2253-61. A Review of Obinutuzumab (GA101), a Novel Type II Anti-CD20 Monoclonal Antibody, for the Treatment	2.5 1.3 843] 4 rgE 1.2	218 170 BT / Overlock 138
11 12 13 14	Doseâ€intensified chemotherapy alone or in combination with mogamulizumab in newly diagnosed aggressive adult Tā€cell leukaemiaâ€lymphoma: a randomized phase ⟨scp⟩ll⟨Jscp⟩ study. British Journal of Haematology, 2015, 169, 672-682. Toxicity Grading Criteria of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology, 1993, 23, 250-257. Feasibility and pharmacokinetic study of a chimeric anti-CD20 monoclonal antibody (IDEC-C2B8,) Tj ETQq1 1 0.7. Significance of cyclin D1 overexpression for the diagnosis of mantle cell lymphoma: a clinicopathologic comparison of cyclin D1-positive MCL and cyclin D1-negative MCL-like B-cell lymphoma. Blood, 2000, 95, 2253-61. A Review of Obinutuzumab (GA101), a Novel Type II Anti-CD20 Monoclonal Antibody, for the Treatment of Patients with B-Cell Malignancies. Advances in Therapy, 2017, 34, 324-356. Factors affecting toxicity, response and progression-free survival in relapsed patients with indolent B-cell lymphoma and mantle cell lymphoma treated with rituximab: a Japanese phase II study. Annals of	2.5 1.3 843] 4 rgE 1.4 2.9	218 170 3T / Overlock 138 128

#	Article	IF	CITATIONS
19	Multicenter Phase II Study of Lenalidomide in Relapsed or Recurrent Adult T-Cell Leukemia/Lymphoma: ATLL-002. Journal of Clinical Oncology, 2016, 34, 4086-4093.	1.6	123
20	A multicentre phase <scp>II</scp> study of vorinostat in patients with relapsed or refractory indolent Bâ€eell nonâ€Hodgkin lymphoma and mantle cell lymphoma. British Journal of Haematology, 2014, 165, 768-776.	2.5	104
21	Targeting Excessive EZH1 and EZH2 Activities for Abnormal Histone Methylation and Transcription Network in Malignant Lymphomas. Cell Reports, 2019, 29, 2321-2337.e7.	6.4	100
22	Japanese multicenter phase II and pharmacokinetic study of rituximab in relapsed or refractory patients with aggressive B-cell lymphoma. Annals of Oncology, 2004, 15, 821-830.	1.2	99
23	Deoxycoformycin-Containing Combination Chemotherapy for Adult T-Cell Leukemia-Lymphoma: Japan Clinical Oncology Group Study (JCOG9109). International Journal of Hematology, 2003, 77, 164-170.	1.6	91
24	Clinical development of antiâ€ <scp>CD</scp> 19 chimeric antigen receptor Tâ€cell therapy for Bâ€cell nonâ€Hodgkin lymphoma. Cancer Science, 2017, 108, 1109-1118.	3.9	91
25	Japan Clinical Oncology Group (JCOG) prognostic index and characterization of long-term survivors of aggressive adult T-cell leukaemia-lymphoma (JCOG0902A). British Journal of Haematology, 2014, 166, 739-748.	2.5	79
26	A Review of New Findings in Adult T-cell Leukemia–Lymphoma: A Focus on Current and Emerging Treatment Strategies. Advances in Therapy, 2018, 35, 135-152.	2.9	72
27	Human T-cell Lymphotropic Virus Type I–Associated Adult T-cell Leukemia–Lymphoma: New Directions in Clinical Research. Clinical Cancer Research, 2014, 20, 5217-5225.	7.0	68
28	Epidemiological and clinical features of adult Tâ€cell leukemia–lymphoma in Japan, 2010–2011: A nationwide survey. Cancer Science, 2017, 108, 2478-2486.	3.9	63
29	Multicenter phase II study of bendamustine for relapsed or refractory indolent Bâ€cell nonâ€Hodgkin lymphoma and mantle cell lymphoma. Cancer Science, 2010, 101, 2059-2064.	3.9	61
30	Toxicity grading criteria of the Japan Clinical Oncology Group. The Clinical Trial Review Committee of the Japan Clinical Oncology Group. Japanese Journal of Clinical Oncology, 1993, 23, 250-7.	1.3	60
31	Phase II/III Study of R-CHOP-21 Versus R-CHOP-14 for Untreated Indolent B-Cell Non-Hodgkin's Lymphoma: JCOG 0203 Trial. Journal of Clinical Oncology, 2011, 29, 3990-3998.	1.6	59
32	Primary Mediastinal Lymphoma. Journal of Computer Assisted Tomography, 2004, 28, 782-789.	0.9	55
33	Phase I study of inotuzumab ozogamicin (CMCâ€544) in Japanese patients with follicular lymphoma pretreated with rituximabâ€based therapy. Cancer Science, 2010, 101, 1840-1845.	3.9	55
34	Mogamulizumab for relapsed adult Tâ€cell leukemia–lymphoma: Updated followâ€up analysis of phase I and <scp>II</scp> studies. Cancer Science, 2017, 108, 2022-2029.	3.9	55
35	Randomized phase II study of biweekly CHOP anddose-escalated CHOP with prophylactic use of lenograstim (glycosylated G-CSF) in aggressive non-Hodgkin's lymphoma: Japan Clinical Oncology Group Study 9505. Annals of Oncology, 2002, 13, 1347-1355.	1.2	48
36	Lenalidomide in relapsed adult T-cell leukaemia-lymphoma or peripheral T-cell lymphoma (ATLL-001): a phase 1, multicentre, dose-escalation study. Lancet Haematology,the, 2016, 3, e107-e118.	4.6	48

#	Article	IF	CITATIONS
37	Primary Hepatic Low-Grade B-Cell Lymphoma of the Mucosa-Associated Lymphoid Tissue Type: A Case Report and Review of the Literature. International Journal of Hematology, 2002, 75, 85-90.	1.6	44
38	Phase II Study of Cladribine (2-Chlorodeoxyadenosine) in Relapsed or Refractory Adult T-Cell Leukemia-Lymphoma. International Journal of Hematology, 2003, 77, 512-517.	1.6	44
39	Re-Treatment of Relapsed Indolent B-Cell Lymphoma With Rituximab. International Journal of Hematology, 2001, 73, 213-221.	1.6	43
40	Targeting Chemokine Receptor CCR4 in Adult T-Cell Leukemia-Lymphoma and Other T-Cell Lymphomas. Current Hematologic Malignancy Reports, 2012, 7, 235-240.	2.3	43
41	Proteasome inhibitor, bortezomib, for myeloma and lymphoma. International Journal of Clinical Oncology, 2007, 12, 318-326.	2.2	42
42	Durable Response but Prolonged Cytopenia after Cladribine Treatment in Relapsed Patients with Indolent non-Hodgkin's Lymphomas: Results of a Japanese Phase II Study. International Journal of Hematology, 2004, 80, 267-277.	1.6	40
43	Histological and immunophenotypic changes in 59 cases of Bâ€cell nonâ€Hodgkin's lymphoma after rituximab therapy. Cancer Science, 2009, 100, 54-61.	3.9	38
44	Phase II Study of Oral Fludarabine Phosphate in Relapsed Indolent B-Cell Non-Hodgkin's Lymphoma. Journal of Clinical Oncology, 2006, 24, 174-180.	1.6	36
45	Japanese phase II study of ⁹⁰ Yâ€ibritumomab tiuxetan in patients with relapsed or refractory indolent Bâ€cell lymphoma. Cancer Science, 2009, 100, 158-164.	3.9	35
46	Deletion of the TNFAIP3/A20gene detected by FICTION analysis in classical Hodgkin lymphoma. BMC Cancer, 2012, 12, 457.	2.6	34
47	Phase I study of YK-176 (2'-deoxycoformycin) in patients with adult T-cell leukemia-lymphoma. The DCF Study Group. Japanese Journal of Clinical Oncology, 1992, 22, 164-71.	1.3	34
48	γδT-cell neoplasms: a clinicopathological study of 11 cases. Annals of Oncology, 2002, 13, 1792-1798.	1.2	33
49	t(14;16)-positive multiple myeloma shows negativity for CD56 expression and unfavorable outcome even in the era of novel drugs. Blood Cancer Journal, 2015, 5, e285-e285.	6.2	33
50	Biology and treatment of HTLV-1 associated T-cell lymphomas. Best Practice and Research in Clinical Haematology, 2013, 26, 3-14.	1.7	32
51	Clinical Features and Current Optimal Management of Natural Killer/T-Cell Lymphoma. Hematology/Oncology Clinics of North America, 2017, 31, 239-253.	2.2	32
52	Phase I study of obinutuzumab (<scp>GA</scp> 101) in <scp>J</scp> apanese patients with relapsed or refractory <scp>B</scp> â€cell nonâ€ <scp>H</scp> odgkin lymphoma. Cancer Science, 2013, 104, 105-110.	3.9	31
53	Pretreatment total serum protein is a significant prognostic factor for the outcome of patients with peripheral T/natural killer-cell lymphomas. Leukemia and Lymphoma, 2010, 51, 813-821.	1.3	30
54	Clinical Trials and Treatment of ATL. Leukemia Research and Treatment, 2012, 2012, 1-12.	2.0	29

#	Article	IF	Citations
55	Chimeric antigen receptor T-cell therapy for B-cell non-Hodgkin lymphoma: opportunities and challenges. Drugs in Context, 2019, 8, 1-14.	2.2	29
56	Randomized phase II study of concurrent and sequential rituximab and CHOP chemotherapy in untreated indolent B-cell lymphoma. Cancer Science, 2006, 97, 305-312.	3.9	28
57	Safety and tolerability of ibrutinib monotherapy in Japanese patients with relapsed/refractory B cell malignancies. International Journal of Hematology, 2016, 103, 86-94.	1.6	28
58	Safety and efficacy of mogamulizumab in patients with adult T-cell leukemia–lymphoma in Japan: interim results of postmarketing all-case surveillance. International Journal of Hematology, 2017, 106, 522-532.	1.6	28
59	Mogamulizumab for the treatment of T-cell lymphoma. Expert Opinion on Biological Therapy, 2017, 17, 1145-1153.	3.1	28
60	Forodesine in the treatment of relapsed/refractory peripheral T-cell lymphoma: an evidence-based review. OncoTargets and Therapy, 2018, Volume 11, 2287-2293.	2.0	27
61	Development of new agents for peripheral T-cell lymphoma. Expert Opinion on Biological Therapy, 2019, 19, 197-209.	3.1	26
62	Interferon Alfa and Zidovudine in Adult T-Cell Leukemia–Lymphoma. New England Journal of Medicine, 1995, 333, 1285-1286.	27.0	25
63	Phase I study of LY2469298, an Fcâ€engineered humanized antiâ€CD20 antibody, in patients with relapsed or refractory follicular lymphoma. Cancer Science, 2011, 102, 432-438.	3.9	25
64	Lymphoma Study Group of JCOG. Japanese Journal of Clinical Oncology, 2012, 42, 85-95.	1.3	25
65	Targeting EZH2 with tazemetostat. Lancet Oncology, The, 2018, 19, 586-587.	10.7	24
66	Monoclonal Antibody Therapy for B-Cell Lymphoma: Clinical Trials of an Anti-CD20 Monoclonal Antibody for B-Cell Lymphoma in Japan. International Journal of Hematology, 2002, 76, 411-419.	1.6	23
67	Phase I study of the oral mammalian target of rapamycin inhibitor everolimus (RAD001) in Japanese patients with relapsed or refractory non-Hodgkin lymphoma. International Journal of Hematology, 2010, 92, 563-570.	1.6	23
68	Rituximab monotherapy with eight weekly infusions for relapsed or refractory patients with indolent B cell nonâ∈Hodgkin lymphoma mostly pretreated with rituximab: A multicenter phase II study. Cancer Science, 2011, 102, 1698-1705.	3.9	21
69	Clinical Trials for Malignant Lymphoma in Japan. Japanese Journal of Clinical Oncology, 2004, 34, 369-378.	1.3	19
70	Adult T-cell leukemia–lymphoma: current treatment strategies and novel immunological approaches. Expert Review of Hematology, 2010, 3, 743-753.	2.2	19
71	Efficacy and safety of ibrutinib in Japanese patients with relapsed or refractory mantle cell lymphoma. Cancer Science, 2016, 107, 1785-1790.	3.9	19
72	Combination Phase I/II Study of Irinotecan Hydrochloride (CPT-11) and Carboplatin in Relapsed or Refractory Non-Hodgkin's Lymphoma. Japanese Journal of Clinical Oncology, 1996, 26, 455-460.	1.3	17

#	Article	IF	CITATIONS
73	Phase I study of radioimmunotherapy with an anti-CD20 murine radioimmunoconjugate (90Y-ibritumomab tiuxetan) in relapsed or refractory indolent B-cell lymphoma. Cancer Science, 2005, 96, 903-910.	3.9	17
74	Clinical features and outcomes of 139 Japanese patients with Hodgkin lymphoma. International Journal of Hematology, 2016, 104, 236-244.	1.6	17
75	Current management of adult T-cell leukemia/lymphoma. Oncology, 2009, 23, 1250-6.	0.5	17
76	Phase I study of cladribine (2-chlorodeoxyadenosine) in lymphoid malignancies. Cladribine Study Group. Japanese Journal of Clinical Oncology, 1997, 27, 146-153.	1.3	16
77	Detection of t(11;18) in MALT-Type Lymphoma With Dual-Color Fluorescence In Situ Hybridization and Reverse Transcriptase–Polymerase Chain Reaction Analysis. Diagnostic Molecular Pathology, 2001, 10, 207-213.	2.1	16
78	Bclâ€2, <scp>B</scp> clâ€6, and the <scp>I</scp> nternational <scp>P</scp> rognostic <scp>I</scp> ndex are prognostic indicators in patients with diffuse large <scp>B</scp> â€cell lymphoma treated with rituximabâ€containing chemotherapy. Cancer Science, 2012, 103, 1898-1904.	3.9	16
79	Antibody therapy targeting CD19 for B-cell non-Hodgkin's lymphoma. Annals of Oncology, 2018, 29, 1086-1089.	1.2	16
80	Unexpected Hepatotoxicities in Patients with Non-Hodgkin's Lymphoma Treated with Irinotecan (CPT-11) and Etoposide. Japanese Journal of Clinical Oncology, 1998, 28, 502-506.	1.3	15
81	Two Entities of Precursor T-Cell Lymphoblastic Leukemia/Lymphoma Based on Radiologic and Immunophenotypic Findings. International Journal of Hematology, 2004, 80, 43-51.	1.6	15
82	A Randomized Controlled Trial Investigating the Survival Benefit of Dose-Intensified Multidrug Combination Chemotherapy (LSG9) for Intermediate- or High-Grade Non-Hodgkin's Lymphoma: Japan Clinical Oncology Group Study 9002. International Journal of Hematology, 2004, 80, 341-350.	1.6	15
83	Phase II study of chemotherapy and stem cell transplantation for adult acute lymphoblastic leukemia or lymphoblastic lymphoma: Japan Clinical Oncology Group study 9004. Cancer Science, 2007, 98, 1350-1357.	3.9	14
84	Randomized phase II study of concurrent and sequential combinations of rituximab plus CHOP (cyclophosphamide, doxorubicin, vincristine and prednisolone) chemotherapy in untreated indolent Bâ€cell nonâ€Hodgkin lymphoma: 7â€year followâ€up results. Cancer Science, 2010, 101, 2579-2585.	3.9	14
85	Long-term Follow-up Results of Adult Patients with Acute Lymphocytic Leukemia or Lymphoblastic Lymphoma Treated with Short-term, Alternating Non-cross-resistant Chemotherapy: Japan Clinical Oncology Group Study 8702. Japanese Journal of Clinical Oncology, 1999, 29, 340-348.	1.3	13
86	Treatment of Indolent Non-Hodgkin's Lymphoma with Cladribine as Single-Agent Therapy and in Combination with Mitoxantrone. International Journal of Hematology, 2004, 79, 311-321.	1.6	13
87	Rituximab and other emerging antibodies as molecular target-based therapy of lymphoma. International Journal of Clinical Oncology, 2003, 8, 212-223.	2.2	12
88	Primary Mediastinal Large B-Cell Lymphoma: A Single-Institution Clinical Study in Japan. International Journal of Hematology, 2004, 79, 465-471.	1.6	11
89	Randomized phase II study of mogamulizumab (KW-0761) plus VCAP-AMP-VECP (mLSG15) versus mLSG15 alone for newly diagnosed aggressive adult T-cell leukemia-lymphoma (ATL) Journal of Clinical Oncology, 2013, 31, 8506-8506.	1.6	11
90	Rituximab and other emerging monoclonal antibody therapies for lymphoma. Expert Opinion on Emerging Drugs, 2002, 7, 289-302.	2.4	10

#	Article	IF	CITATIONS
91	Clinical trials of a mouse-human chimeric anti-CD20 monoclonal antibody (rituximab) for B cell non-Hodgkin's lymphoma in Japan. Cancer Chemotherapy and Pharmacology, 2001, 48, S85-S90.	2.3	9
92	Clinical Trials for Human T-Cell Lymphotropic Virus Type I–Associated Peripheral T-Cell Lymphoma in Japan. Seminars in Hematology, 2010, 47, S5-S7.	3.4	9
93	Putting the Clinical and Biological Heterogeneity of Non-Hodgkin Lymphoma into Context. Clinical Cancer Research, 2014, 20, 5173-5181.	7.0	9
94	Efficacy and safety of obinutuzumab in patients with previously untreated follicular lymphoma: a subgroup analysis of patients enrolled in Japan in the randomized phase III GALLIUM trial. International Journal of Hematology, 2018, 108, 499-509.	1.6	9
95	Safety and Efficacy of Brentuximab Vedotin in the Treatment of Classic Hodgkin Lymphoma OncoTargets and Therapy, 2020, Volume 13, 5993-6009.	2.0	9
96	A randomized phase 2/3 study of R-CHOP vs CHOP combined with dose-dense rituximab for DLBCL: the JCOG0601 trial. Blood Advances, 2021, 5, 984-993.	5 . 2	9
97	Phase II study of oral fludarabine in combination with rituximab for relapsed indolent Bâ€cell nonâ€Hodgkin lymphoma. Cancer Science, 2009, 100, 1951-1956.	3.9	8
98	Adult T-Cell Leukemia-Lymphoma Successfully Treated with 2-Chlorodeoxyadenosine Internal Medicine, 1998, 37, 411-413.	0.7	7
99	Granular lymphocytic leukemia derived from γδT-cell expressing cytotoxic molecules. Leukemia Research, 2001, 25, 259-261.	0.8	7
100	Phase I and pharmacokinetic study of oral fludarabine phosphate in relapsed indolent B-cell non-Hodgkin's lymphoma. Annals of Oncology, 2006, 17, 330-333.	1.2	7
101	PhaseÂl/II and pharmacokinetic study of cladribine with 2â€h infusion in Japanese patients with relapsed indolent Bâ€cell lymphoma mostly pretreated with rituximab. Cancer Science, 2009, 100, 1344-1350.	3.9	7
102	Japanese phase II study of rituximab maintenance for untreated indolent B-cell non-Hodgkin lymphoma with high tumor burden. International Journal of Hematology, 2016, 104, 700-708.	1.6	7
103	Partial deletion of the <scp>ALK</scp> gene in <scp>ALK</scp> â€positive anaplastic large cell lymphoma. Hematological Oncology, 2018, 36, 150-158.	1.7	7
104	EBV-Positive Burkitt Lymphoma as a Late-Onset Posttransplantion Lymphoproliferative Disorder after Allogeneic Stem Cell Transplantation. International Journal of Hematology, 2004, 79, 387-389.	1.6	6
105	Classical Hodgkin lymphoma primary refractory to brentuximab vedotin, with transformation to CD30-positive diffuse large B-cell lymphoma. International Journal of Hematology, 2016, 104, 396-399.	1.6	6
106	Analysis of Japanese patients from the AUGMENT phase III study of lenalidomide + rituximab (R2) vs. rituximab + placebo in relapsed/refractory indolent non-Hodgkin lymphoma. International Journal of Hematology, 2020, 111, 409-416.	1.6	6
107	Safety considerations with targeted therapy drugs for B-cell non-Hodgkin lymphoma. Expert Opinion on Drug Safety, 2020, 19, 1105-1120.	2.4	6
108	4. Antibody Therapy for Malignant Lymphoma. Internal Medicine, 2007, 46, 99-100.	0.7	5

#	Article	IF	CITATIONS
109	Phase II Study of Intensive Post-remission Chemotherapy and Stem Cell Transplantation for Adult Acute Lymphoblastic Leukemia and Lymphoblastic Lymphoma: Japan Clinical Oncology Group Study, JCOG9402. Japanese Journal of Clinical Oncology, 2012, 42, 394-404.	1.3	5
110	Clinical development of voxtalisib: a pan-PI3K/mTOR inhibitor. Lancet Haematology, the, 2018, 5, e134-e135.	4.6	5
111	Diseaseâ€oriented treatment of <scp>T</scp> â€cell lymphoma. Hematological Oncology, 2017, 35, 54-59.	1.7	4
112	Ibrutinib in Japanese patients with relapsed/refractory B-cell malignancies: final analysis of phase I study. International Journal of Hematology, 2019, 109, 366-368.	1.6	4
113	Adult T-Cell Leukemia-Lymphoma. Cancer Treatment and Research, 2019, 176, 145-161.	0.5	4
114	Adult T-Cell Leukemia-Lymphoma. , 2008, , 2425-2441.		4
115	Subcutaneous rituximab: a practical approach?. Lancet Oncology, The, 2014, 15, 254-255.	10.7	3
116	Safety, tolerability and pharmacokinetics of shorter duration of infusion of obinutuzumab in Japanese patients with B-cell non-Hodgkin lymphoma: final results of the phase II GATS study. Japanese Journal of Clinical Oncology, 2018, 48, 736-742.	1.3	3
117	Phase I Study of KW-0761, a Defucosylated Anti-CCR4 Antibody, in Relapsed Patients (Pts) with Adult T-Cell Leukemia-Lymphoma (ATL) or Peripheral T-Cell Lymphoma (PTCL): Updated Results Blood, 2008, 112, 1007-1007.	1.4	3
118	Rituximab biosimilars: introduction into clinical practice. Lancet Haematology,the, 2017, 4, e342-e343.	4.6	2
119	HTLV-1-Associated T-cell Diseases. , 2013, , 113-135.		2
120	Pharmacokinetic (PK) and Pharmacodynamic (PD) Profiles of Bortezomib (B) in Patients (pts) with Relapsed Multiple Myeloma (MM): A Phase I/II Study in Japan Blood, 2006, 108, 5106-5106.	1.4	2
121	Randomized phase II study of concurrent and sequential combinations of rituximab (R) plus CHOP (R-CHOP) in untreated indolent B-NHL: 7-year follow-up results. Journal of Clinical Oncology, 2008, 26, 8616-8616.	1.6	2
122	The effect of vitamin D2 on hypocalcemia in patients under chronic hemodialysis Tohoku Journal of Experimental Medicine, 1980, 131, 249-255.	1.2	1
123	Monoclonal antibodies for the treatment of hematologic malignancies: clinical trials in Japan. Cancer Chemotherapy and Pharmacology, 2003, 52, 90-96.	2.3	1
124	t(11;18)-Bearing Pulmonary Mucosa-Associated Lymphoid Tissue Lymphoma Responding to Cladribine. International Journal of Hematology, 2004, 80, 70-74.	1.6	1
125	Guest editorial: Management of malignant lymphoma is continuously improving. International Journal of Hematology, 2012, 96, 533-534.	1.6	0
126	An evaluation of mogamulizumab for the treatment of peripheral T-cell lymphoma. Expert Opinion on Orphan Drugs, 2014, 2, 735-742.	0.8	0

lF

CITATIONS

127	Clinical characteristics of patients with B-cell lymphoma enrolled in clinical trials for aggressive lymphoma in Japan: Japan Clinical Oncology Group - Lymphoma Study Group study – JCOG0108A. Journal of Clinical and Experimental Hematopathology: JCEH, 2021, 61, 35-41.	0.8	0
128	3.骨髄腫・リンパ腫治ç™,ã¸ã®ãƒ—ãƒãƒ†ã,¢ã,½ãƒ¼ãƒé~»å®³å‰ Bortezomibã®å°Žå…¥ï¼šè—¬(¢%a©å‹•	æΩ.‹ã « 基ã
129	Adult T-cell Leukemia-Lymphoma. , 2014, , 99-110.		O
130	Adult T-Cell Leukemia-Lymphoma. , 2014, , 2076-2092.e4.		0

ARTICLE