

Yan-Hong Zhou

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6157258/yan-hong-zhou-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

94 papers	6,073 citations	42 h-index	76 g-index
97 ext. papers	7,773 ext. citations	6.7 avg, IF	5.74 L-index

#	Paper	IF	Citations
94	High Nitric Oxide Concentration Inhibits Photosynthetic Pigment Biosynthesis by Promoting the Degradation of Transcription Factor HY5 in Tomato. <i>International Journal of Molecular Sciences</i> , 2022 , 23, 6027	6.3	1
93	Light-dependent activation of HY5 promotes mycorrhizal symbiosis in tomato by systemically regulating strigolactone biosynthesis. <i>New Phytologist</i> , 2021 ,	9.8	5
92	Strigolactones positively regulate abscisic acid-dependent heat and cold tolerance in tomato. <i>Horticulture Research</i> , 2021 , 8, 237	7.7	8
91	ELONGATED HYPOCOTYL 5 mediates blue light-induced starch degradation in tomato. <i>Journal of Experimental Botany</i> , 2021 , 72, 2627-2641	7	7
90	Crosstalk between Brassinosteroid and Redox Signaling Contributes to the Activation of CBF Expression during Cold Responses in Tomato. <i>Antioxidants</i> , 2021 , 10,	7.1	7
89	Brassinosteroid signaling integrates multiple pathways to release apical dominance in tomato. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	9
88	The phyB-dependent induction of HY5 promotes iron uptake by systemically activating FER expression. <i>EMBO Reports</i> , 2021 , 22, e51944	6.5	8
87	Ethylene response factors 15 and 16 trigger jasmonate biosynthesis in tomato during herbivore resistance. <i>Plant Physiology</i> , 2021 , 185, 1182-1197	6.6	9
86	Light regulation of horticultural crop nutrient uptake and utilization. <i>Horticultural Plant Journal</i> , 2021 , 7, 367-379	4.3	13
85	Brassinosteroids act as a positive regulator of NBR1-dependent selective autophagy in response to chilling stress in tomato. <i>Journal of Experimental Botany</i> , 2020 , 71, 1092-1106	7	27
84	Crosstalk of PIF4 and DELLA modulates CBF transcript and hormone homeostasis in cold response in tomato. <i>Plant Biotechnology Journal</i> , 2020 , 18, 1041-1055	11.6	31
83	Brassinosteroid-mediated reactive oxygen species are essential for tapetum degradation and pollen fertility in tomato. <i>Plant Journal</i> , 2020 , 102, 931-947	6.9	22
82	Light-induced HY5 Functions as a Systemic Signal to Coordinate the Photoprotective Response to Light Fluctuation. <i>Plant Physiology</i> , 2020 , 184, 1181-1193	6.6	9
81	The HY5 and MYB15 transcription factors positively regulate cold tolerance in tomato via the CBF pathway. <i>Plant, Cell and Environment</i> , 2020 , 43, 2712-2726	8.4	20
80	Systemic Root-Shoot Signaling Drives Jasmonate-Based Root Defense against Nematodes. <i>Current Biology</i> , 2019 , 29, 3430-3438.e4	6.3	43
79	Brassinosteroids Act as a Positive Regulator of Photoprotection in Response to Chilling Stress. <i>Plant Physiology</i> , 2019 , 180, 2061-2076	6.6	46
78	A novel CO ₂ -responsive systemic signaling pathway controlling plant mycorrhizal symbiosis. <i>New Phytologist</i> , 2019 , 224, 106-116	9.8	20

77	Tomato GLR3.3 and GLR3.5 mediate cold acclimation-induced chilling tolerance by regulating apoplastic H ₂ O ₂ production and redox homeostasis. <i>Plant, Cell and Environment</i> , 2019 , 42, 3326-3339	8.4	28
76	Strigolactones positively regulate defense against root-knot nematodes in tomato. <i>Journal of Experimental Botany</i> , 2019 , 70, 1325-1337	7	27
75	SlHY5 Integrates Temperature, Light, and Hormone Signaling to Balance Plant Growth and Cold Tolerance. <i>Plant Physiology</i> , 2019 , 179, 749-760	6.6	39
74	BZR1 Mediates Brassinosteroid-Induced Autophagy and Nitrogen Starvation in Tomato. <i>Plant Physiology</i> , 2019 , 179, 671-685	6.6	53
73	A Plant Phytosulfokine Peptide Initiates Auxin-Dependent Immunity through Cytosolic Ca Signaling in Tomato. <i>Plant Cell</i> , 2018 , 30, 652-667	11.6	72
72	The bZip transcription factor HY5 mediates CRY1a-induced anthocyanin biosynthesis in tomato. <i>Plant, Cell and Environment</i> , 2018 , 41, 1762-1775	8.4	68
71	Heat Shock Factor HsfA1a Is Essential for Gene-Mediated Nematode Resistance and Triggers H ₂ O ₂ Production. <i>Plant Physiology</i> , 2018 , 176, 2456-2471	6.6	30
70	Brassinosteroids act as a positive regulator for resistance against root-knot nematode involving RESPIRATORY BURST OXIDASE HOMOLOG-dependent activation of MAPKs in tomato. <i>Plant, Cell and Environment</i> , 2018 , 41, 1113-1125	8.4	35
69	Tomato CRY1a plays a critical role in the regulation of phytohormone homeostasis, plant development, and carotenoid metabolism in fruits. <i>Plant, Cell and Environment</i> , 2018 , 41, 354-366	8.4	28
68	Glutaredoxin GRXS16 mediates brassinosteroid-induced apoplastic H ₂ O ₂ production to promote pesticide metabolism in tomato. <i>Environmental Pollution</i> , 2018 , 240, 227-234	9.3	22
67	BZR1 Transcription Factor Regulates Heat Stress Tolerance Through FERONIA Receptor-Like Kinase-Mediated Reactive Oxygen Species Signaling in Tomato. <i>Plant and Cell Physiology</i> , 2018 , 59, 2239-2254	4.9	52
66	The role of calcium-dependent protein kinase in hydrogen peroxide, nitric oxide and ABA-dependent cold acclimation. <i>Journal of Experimental Botany</i> , 2018 , 69, 4127-4139	7	47
65	Light Signaling-Dependent Regulation of Photoinhibition and Photoprotection in Tomato. <i>Plant Physiology</i> , 2018 , 176, 1311-1326	6.6	52
64	Brassinosteroid-mediated apoplastic H ₂ O ₂ -glutaredoxin 12/14 cascade regulates antioxidant capacity in response to chilling in tomato. <i>Plant, Cell and Environment</i> , 2018 , 41, 1052-1064	8.4	53
63	HsfA1a upregulates melatonin biosynthesis to confer cadmium tolerance in tomato plants. <i>Journal of Pineal Research</i> , 2017 , 62, e12387	10.4	130
62	Nitric oxide is involved in the oxytetracycline-induced suppression of root growth through inhibiting hydrogen peroxide accumulation in the root meristem. <i>Scientific Reports</i> , 2017 , 7, 43096	4.9	13
61	Crosstalk between Nitric Oxide and MPK1/2 Mediates Cold Acclimation-induced Chilling Tolerance in Tomato. <i>Plant and Cell Physiology</i> , 2017 , 58, 1963-1975	4.9	35
60	24-Epibrassinolide alleviates organic pollutants-retarded root elongation by promoting redox homeostasis and secondary metabolism in <i>Cucumis sativus</i> L. <i>Environmental Pollution</i> , 2017 , 229, 922-931	9.3	38

59	Systemic Induction of Photosynthesis via Illumination of the Shoot Apex Is Mediated Sequentially by Phytochrome B, Auxin and Hydrogen Peroxide in Tomato. <i>Plant Physiology</i> , 2016 , 172, 1259-1272	6.6	46
58	Role of Hormones in Plant Adaptation to Heat Stress 2016 , 1-21		17
57	Interactions between 2-Cys peroxiredoxins and ascorbate in autophagosome formation during the heat stress response in <i>Solanum lycopersicum</i> . <i>Journal of Experimental Botany</i> , 2016 , 67, 1919-33	7	20
56	Phytochrome A and B Function Antagonistically to Regulate Cold Tolerance via Absciscic Acid-Dependent Jasmonate Signaling. <i>Plant Physiology</i> , 2016 , 170, 459-71	6.6	133
55	Genome-Wide Identification and Expression Analysis of Calcium-dependent Protein Kinase in Tomato. <i>Frontiers in Plant Science</i> , 2016 , 7, 469	6.2	36
54	Unraveling Main Limiting Sites of Photosynthesis under Below- and Above-Ground Heat Stress in Cucumber and the Alleviatory Role of Luffa Rootstock. <i>Frontiers in Plant Science</i> , 2016 , 7, 746	6.2	17
53	Melatonin mediates selenium-induced tolerance to cadmium stress in tomato plants. <i>Journal of Pineal Research</i> , 2016 , 61, 291-302	10.4	140
52	Apoplastic H ₂ O ₂ plays a critical role in axillary bud outgrowth by altering auxin and cytokinin homeostasis in tomato plants. <i>New Phytologist</i> , 2016 , 211, 1266-78	9.8	34
51	Microarray and genetic analysis reveals that csa-miR159b plays a critical role in absciscic acid-mediated heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2016 , 39, 1790-804	8.4	36
50	Grafting cucumber onto luffa improves drought tolerance by increasing ABA biosynthesis and sensitivity. <i>Scientific Reports</i> , 2016 , 6, 20212	4.9	37
49	Interplay between mitogen-activated protein kinase and nitric oxide in brassinosteroid-induced pesticide metabolism in <i>Solanum lycopersicum</i> . <i>Journal of Hazardous Materials</i> , 2016 , 316, 221-31	12.8	24
48	DWARF overexpression induces alteration in phytohormone homeostasis, development, architecture and carotenoid accumulation in tomato. <i>Plant Biotechnology Journal</i> , 2016 , 14, 1021-33	11.6	50
47	Overexpression of a brassinosteroid biosynthetic gene Dwarf enhances photosynthetic capacity through activation of Calvin cycle enzymes in tomato. <i>BMC Plant Biology</i> , 2016 , 16, 33	5.3	42
46	Melatonin enhances thermotolerance by promoting cellular protein protection in tomato plants. <i>Journal of Pineal Research</i> , 2016 , 61, 457-469	10.4	146
45	Salicylic acid binding of mitochondrial alpha-ketoglutarate dehydrogenase E2 affects mitochondrial oxidative phosphorylation and electron transport chain components and plays a role in basal defense against tobacco mosaic virus in tomato. <i>New Phytologist</i> , 2015 , 205, 1296-1307	9.8	45
44	Brassinosteroids play a critical role in the regulation of pesticide metabolism in crop plants. <i>Scientific Reports</i> , 2015 , 5, 9018	4.9	73
43	RNA-seq analysis reveals the role of red light in resistance against <i>Pseudomonas syringae</i> pv. tomato DC3000 in tomato plants. <i>BMC Genomics</i> , 2015 , 16, 120	4.5	52
42	Interplay between reactive oxygen species and hormones in the control of plant development and stress tolerance. <i>Journal of Experimental Botany</i> , 2015 , 66, 2839-56	7	401

41	Enhanced photosynthetic capacity and antioxidant potential mediate brassinosteroid-induced phenanthrene stress tolerance in tomato. <i>Environmental Pollution</i> , 2015 , 201, 58-66	9.3	29
40	Antagonism between phytohormone signalling underlies the variation in disease susceptibility of tomato plants under elevated CO ₂ . <i>Journal of Experimental Botany</i> , 2015 , 66, 1951-63	7	77
39	High atmospheric carbon dioxide-dependent alleviation of salt stress is linked to RESPIRATORY BURST OXIDASE 1 (RBOH1)-dependent H ₂ O ₂ production in tomato (<i>Solanum lycopersicum</i>). <i>Journal of Experimental Botany</i> , 2015 , 66, 7391-404	7	34
38	Light-induced systemic resistance in tomato plants against root-knot nematode <i>Meloidogyne incognita</i> . <i>Plant Growth Regulation</i> , 2015 , 76, 167-175	3.2	14
37	Tomato- <i>Pseudomonas syringae</i> interactions under elevated CO ₂ concentration: the role of stomata. <i>Journal of Experimental Botany</i> , 2015 , 66, 307-16	7	24
36	Guard cell hydrogen peroxide and nitric oxide mediate elevated CO ₂ -induced stomatal movement in tomato. <i>New Phytologist</i> , 2015 , 208, 342-53	9.8	63
35	Involvement of nitric oxide in the jasmonate-dependent basal defense against root-knot nematode in tomato plants. <i>Frontiers in Plant Science</i> , 2015 , 6, 193	6.2	37
34	Melatonin mitigates cadmium phytotoxicity through modulation of phytochelatin biosynthesis, vacuolar sequestration, and antioxidant potential in <i>Solanum lycopersicum</i> L. <i>Frontiers in Plant Science</i> , 2015 , 6, 601	6.2	174
33	Tomato HsfA1a plays a critical role in plant drought tolerance by activating ATG genes and inducing autophagy. <i>Autophagy</i> , 2015 , 11, 2033-2047	10.2	109
32	Role of brassinosteroid in plant adaptation to abiotic stresses and its interplay with other hormones. <i>Current Protein and Peptide Science</i> , 2015 , 16, 462-73	2.8	67
31	Role of H ₂ O ₂ dynamics in brassinosteroid-induced stomatal closure and opening in <i>Solanum lycopersicum</i> . <i>Plant, Cell and Environment</i> , 2014 , 37, 2036-50	8.4	113
30	Hydrogen peroxide mediates abscisic acid-induced HSP70 accumulation and heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2014 , 37, 2768-80	8.4	95
29	The sub/supra-optimal temperature-induced inhibition of photosynthesis and oxidative damage in cucumber leaves are alleviated by grafting onto figleaf gourd/luffa rootstocks. <i>Physiologia Plantarum</i> , 2014 , 152, 571-84	4.6	23
28	Effects of <i>Fusarium oxysporum</i> on rhizosphere microbial communities of two cucumber genotypes with contrasting <i>Fusarium</i> wilt resistance under hydroponic condition. <i>European Journal of Plant Pathology</i> , 2014 , 140, 643-653	2.1	10
27	H ₂ O ₂ mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses. <i>Journal of Experimental Botany</i> , 2014 , 65, 4371-83	7	204
26	RBOH1-dependent H ₂ O ₂ production and subsequent activation of MPK1/2 play an important role in acclimation-induced cross-tolerance in tomato. <i>Journal of Experimental Botany</i> , 2014 , 65, 595-607	7	103
25	Chloroplastic thioredoxin-f and thioredoxin-m1/4 play important roles in brassinosteroids-induced changes in CO ₂ assimilation and cellular redox homeostasis in tomato. <i>Journal of Experimental Botany</i> , 2014 , 65, 4335-47	7	23
24	Brassinosteroids accelerate recovery of photosynthetic apparatus from cold stress by balancing the electron partitioning, carboxylation and redox homeostasis in cucumber. <i>Physiologia Plantarum</i> , 2013 , 148, 133-45	4.6	79

23	Glutathione biosynthesis and regeneration play an important role in the metabolism of chlorothalonil in tomato. <i>Chemosphere</i> , 2013 , 90, 2563-70	8.4	32
22	Brassinosteroid alleviates polychlorinated biphenyls-induced oxidative stress by enhancing antioxidant enzymes activity in tomato. <i>Chemosphere</i> , 2013 , 90, 2645-53	8.4	68
21	Silencing of tomato RBOH1 and MPK2 abolishes brassinosteroid-induced H ₂ O ₂ generation and stress tolerance. <i>Plant, Cell and Environment</i> , 2013 , 36, 789-803	8.4	100
20	Brassinosteroids-Induced Systemic Stress Tolerance was Associated with Increased Transcripts of Several Defence-Related Genes in the Phloem in Cucumis sativus. <i>PLoS ONE</i> , 2013 , 8, e66582	3.7	36
19	Cytokinin-induced parthenocarpic fruit development in tomato is partly dependent on enhanced gibberellin and auxin biosynthesis. <i>PLoS ONE</i> , 2013 , 8, e70080	3.7	54
18	Temperature effects on the reactive oxygen species formation and antioxidant defence in roots of two cucurbit species with contrasting root zone temperature optima. <i>Acta Physiologiae Plantarum</i> , 2012 , 34, 713-720	2.6	13
17	Brassinosteroid improves seed germination and early development of tomato seedling under phenanthrene stress. <i>Plant Growth Regulation</i> , 2012 , 68, 87-96	3.2	25
16	Cellular glutathione redox homeostasis plays an important role in the brassinosteroid-induced increase in CO ₂ assimilation in Cucumis sativus. <i>New Phytologist</i> , 2012 , 194, 932-943	9.8	106
15	Induction of systemic stress tolerance by brassinosteroid in Cucumis sativus. <i>New Phytologist</i> , 2011 , 191, 706-720	9.8	113
14	Effects of nitrogen form on growth, CO ₂ assimilation, chlorophyll fluorescence, and photosynthetic electron allocation in cucumber and rice plants. <i>Journal of Zhejiang University: Science B</i> , 2011 , 12, 126-34	4.5	35
13	Functional analysis of the Arabidopsis PAL gene family in plant growth, development, and response to environmental stress. <i>Plant Physiology</i> , 2010 , 153, 1526-38	6.6	450
12	Brassinosteroids promote photosynthesis and growth by enhancing activation of Rubisco and expression of photosynthetic genes in Cucumis sativus. <i>Planta</i> , 2009 , 230, 1185-96	4.7	188
11	Grafting of Cucumis sativus onto Cucurbita ficifolia leads to improved plant growth, increased light utilization and reduced accumulation of reactive oxygen species in chilled plants. <i>Journal of Plant Research</i> , 2009 , 122, 529-40	2.6	34
10	Effects of light quality on CO ₂ assimilation, chlorophyll-fluorescence quenching, expression of Calvin cycle genes and carbohydrate accumulation in Cucumis sativus. <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2009 , 96, 30-7	6.7	159
9	Impact of light variation on development of photoprotection, antioxidants, and nutritional value in Lactuca sativa L. <i>Journal of Agricultural and Food Chemistry</i> , 2009 , 57, 5494-500	5.7	42
8	Reactive oxygen species are involved in brassinosteroid-induced stress tolerance in cucumber. <i>Plant Physiology</i> , 2009 , 150, 801-14	6.6	507
7	Putrescine enhancement of tolerance to root-zone hypoxia in Cucumis sativus: a role for increased nitrate reduction. <i>Functional Plant Biology</i> , 2008 , 35, 337-345	2.7	17
6	Inhibition of photosynthesis and energy dissipation induced by water and high light stresses in rice. <i>Journal of Experimental Botany</i> , 2007 , 58, 1207-17	7	170

5	Adaptation of cucurbit species to changes in substrate temperature: Root growth, antioxidants, and peroxidation 2007 , 50, 527-532		13
4	Chill-induced decrease in capacity of RuBP carboxylation and associated H ₂ O ₂ accumulation in cucumber leaves are alleviated by grafting onto figleaf gourd. <i>Annals of Botany</i> , 2007 , 100, 839-48	4.1	77
3	Genotypic variation of Rubisco expression, photosynthetic electron flow and antioxidant metabolism in the chloroplasts of chill-exposed cucumber plants. <i>Plant and Cell Physiology</i> , 2006 , 47, 192-9	4.9	82
2	Effects of pollination and N-(2-chloro-4-pyridyl)-N-phenylurea on the expression of acid invertase in ovaries of <i>Lagenaria leucantha</i> . <i>Plant Growth Regulation</i> , 2004 , 42, 263-270	3.2	1
1	Greenhouse and field cucumber genotypes use different mechanisms to protect against dark chilling. <i>Functional Plant Biology</i> , 2004 , 31, 1215-1223	2.7	14