

Fabiola Sepulveda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6164834/publications.pdf>

Version: 2024-02-01

13

papers

178

citations

1163117

8

h-index

1125743

13

g-index

13

all docs

13

docs citations

13

times ranked

209

citing authors

#	ARTICLE	IF	CITATIONS
1	Metazoan parasites in wild fish and farmed salmon from aquaculture sites in southern Chile. Aquaculture, 2004, 235, 89-100.	3.5	68
2	Spatio-temporal patterns of genetic variations in populations of yellowtail kingfish <i>< i>Seriola lalandi</i></i> from the south-eastern Pacific Ocean and potential implications for its fishery management. Journal of Fish Biology, 2017, 90, 249-264.	1.6	15
3	Two New Species of <i>< i>Encotyllabe</i></i> (Monogenea: Capsalidae) Based on Morphometric and Molecular Evidence: Parasites of Two Inshore Fish Species of Northern Chile. Journal of Parasitology, 2014, 100, 344-349.	0.7	13
4	The feasibility of using Udonella sp. (Platyhelminthes: Udonellidae) as a biological control for the sea louse <i>Caligus rogercresseyi</i> , Boxshall and Bravo 2000, (Copepoda: Caligidae) in southern Chile. Aquaculture, 2002, 208, 11-21.	3.5	12
5	Udonella australis n. sp. (Monogenea), an epibiont on sea-lice from native fish off southern Chile. Systematic Parasitology, 2002, 52, 67-74.	1.1	11
6	Potential role of ectoparasites (<i>< i>Zeuxapta seriolae</i></i> and <i>< i>Caligus lalandei</i></i>) in the transmission of pathogenic bacteria in yellowtail kingfish <i>< i>Seriola lalandi</i></i> , inferred from cultivable microbiota and molecular analyses. Journal of Fish Diseases, 2017, 40, 979-985.	1.9	11
7	<i>< i>Acanthocotyle gurgesiella</i></i> n. sp. (Monogenea: Acanthocotylidae) from the deep-sea skate <i>< i>Gurgesiella furvescens</i></i> (Rajidae) in the south-eastern Pacific. Journal of Helminthology, 2018, 92, 223-227.	1.0	11
8	Integrative systematics of the genus Limacia O. F. Müller, 1781 (Gastropoda, Heterobranchia,) Tj ETQq0 0 0 rgBT _{1.0} /Overlock ₁₀ Tf 50 4		
9	DNA barcoding evidence for the first recorded transmission of <i>Neobenedenia</i> sp. from wild fish species to <i>Seriola lalandi</i> cultured in an open recirculating system on the Coast of Northern Chile. Aquaculture, 2019, 501, 239-246.	3.5	8
10	Regional population genetics and global phylogeography of the endangered highly migratory shark <i>< i>Lamna nasus</i></i> : Implications for fishery management and conservation. Aquatic Conservation: Marine and Freshwater Ecosystems, 2021, 31, 620-634.	2.0	7
11	Genetic homogeneity coupled with morphometric variability suggests high phenotypic plasticity in the sea louse <i>< i>Caligus rogercresseyi</i></i> (Boxshall and Bravo, 2000), infecting farmed salmon (<i>< i>Salmo</i></i>) Tj ETQq1 1 0.984314rgBT /Ov		
12	Calicotyle hydrolagi n. sp. (Monogenea: Monocotylidae) infecting the deep-sea Eastern Pacific black ghost shark <i>Hydrolagus melanophasma</i> from the Atacama Trench, with comments on host specificity of <i>Calicotyle</i> spp.. Parasitology International, 2020, 75, 102025.	1.3	5
13	Morphological and molecular characterisation of digenetic parasites of the Galápagos sheephead <i>Semicossyphus darwini</i> (Jenyns) with the re-description of <i>Labriter secundus</i> Manter, 1940 (Lepidapedidae) from the Humboldt Current Large Marine Ecosystem. Systematic Parasitology, 2018, 95, 391-401.	1.1	3