

Philippe A Melas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6171158/publications.pdf>

Version: 2024-02-01

27
papers

1,081
citations

516710

16
h-index

526287

27
g-index

27
all docs

27
docs citations

27
times ranked

1813
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic and epigenetic associations of MAOA and NR3C1 with depression and childhood adversities. <i>International Journal of Neuropsychopharmacology</i> , 2013, 16, 1513-1528.	2.1	182
2	Epigenetic aberrations in leukocytes of patients with schizophrenia: association of global DNA methylation with antipsychotic drug treatment and disease onset. <i>FASEB Journal</i> , 2012, 26, 2712-2718.	0.5	170
3	Antidepressant treatment is associated with epigenetic alterations in the promoter of P11 in a genetic model of depression. <i>International Journal of Neuropsychopharmacology</i> , 2012, 15, 669-679.	2.1	114
4	Antidepressant-Like Effect of Sodium Butyrate is Associated with an Increase in TET1 and in 5-Hydroxymethylation Levels in the Bdnf Gene. <i>International Journal of Neuropsychopharmacology</i> , 2015, 18, pyu032-pyu032.	2.1	111
5	Cannabidiol as a Potential Treatment for Anxiety and Mood Disorders: Molecular Targets and Epigenetic Insights from Preclinical Research. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1863.	4.1	60
6	Examining the public refusal to consent to DNA biobanking: empirical data from a Swedish population-based study. <i>Journal of Medical Ethics</i> , 2010, 36, 93-98.	1.8	59
7	NR3C1 hypermethylation in depressed and bullied adolescents. <i>Translational Psychiatry</i> , 2018, 8, 121.	4.8	46
8	Prior alcohol use enhances vulnerability to compulsive cocaine self-administration by promoting degradation of HDAC4 and HDAC5. <i>Science Advances</i> , 2017, 3, e1701682.	10.3	45
9	Cannabinoid exposure in rat adolescence reprograms the initial behavioral, molecular, and epigenetic response to cocaine. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 9991-10002.	7.1	39
10	Hypomethylation of MAOA's first exon region in depression: A replication study. <i>Psychiatry Research</i> , 2015, 226, 389-391.	3.3	37
11	PreproNPY Pro7 protects against depression despite exposure to environmental risk factors. <i>Journal of Affective Disorders</i> , 2009, 118, 124-130.	4.1	33
12	Cannabinoid Modulation of Eukaryotic Initiation Factors (eIF2 β and eIF2B1) and Behavioral Cross-Sensitization to Cocaine in Adolescent Rats. <i>Cell Reports</i> , 2018, 22, 2909-2923.	6.4	23
13	Adolescent cannabinoid exposure induces irritability-like behavior and cocaine cross-sensitization without affecting the escalation of cocaine self-administration in adulthood. <i>Scientific Reports</i> , 2018, 8, 13893.	3.3	23
14	MicroRNA 101b Is Downregulated in the Prefrontal Cortex of a Genetic Model of Depression and Targets the Glutamate Transporter SLC1A1 (EAAT3) <i>in Vitro</i> . <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, pyw069.	2.1	22
15	Mood Stabilizers and the Influence on Global Leukocyte DNA Methylation in Bipolar Disorder. <i>Molecular Neuropsychiatry</i> , 2015, 1, 76-81.	2.9	20
16	Neuropeptide Y: Identification of a novel rat mRNA splice-variant that is downregulated in the hippocampus and the prefrontal cortex of a depression-like model. <i>Peptides</i> , 2012, 35, 49-55.	2.4	19
17	The serotonin transporter promoter variant (5-HTTLPR) and childhood adversity are associated with the personality trait openness to experience. <i>Psychiatry Research</i> , 2017, 257, 322-326.	3.3	16
18	Information Related to Prenatal Genetic Counseling: Interpretation by Adolescents, Effects on Risk Perception and Ethical Implications. <i>Journal of Genetic Counseling</i> , 2012, 21, 536-546.	1.6	11

#	ARTICLE	IF	CITATIONS
19	Deep brain stimulation of the nucleus accumbens shell attenuates cocaine withdrawal but increases cocaine self-administration, cocaine-induced locomotor activity, and GluR1/GluA1 in the central nucleus of the amygdala in male cocaine-dependent rats. <i>Brain Stimulation</i> , 2022, 15, 13-22.	1.6	11
20	Neuropeptide Y, stressful life events and personality trait conscientiousness: Preliminary associations from a Swedish longitudinal study. <i>Psychiatry Research</i> , 2018, 263, 48-53.	3.3	10
21	The cannabinoid receptor-1 gene interacts with stressful life events to increase the risk for problematic alcohol use. <i>Scientific Reports</i> , 2022, 12, 4963.	3.3	9
22	Single-nucleotide polymorphism in the human TIA1 gene interacts with stressful life events to predict the development of pathological anxiety symptoms in a Swedish population. <i>Journal of Affective Disorders</i> , 2020, 260, 597-603.	4.1	6
23	DNA methylation of the glucocorticoid receptor gene predicts substance use in adolescence: longitudinal data from over 1000 young individuals. <i>Translational Psychiatry</i> , 2021, 11, 477.	4.8	6
24	The monoamine stabilizer OSU6162 has anxiolytic-like properties and reduces voluntary alcohol intake in a genetic rat model of depression. <i>Scientific Reports</i> , 2021, 11, 11856.	3.3	3
25	Physical exercise is associated with a reduction in plasma levels of fractalkine, TGF- β 21, eotaxin-1 and IL-6 in younger adults with mobility disability. <i>PLoS ONE</i> , 2022, 17, e0263173.	2.5	3
26	Physical exercise is associated with a reduction in inflammatory biomarkers in first-episode psychosis: A pilot study of CRP, SAA, sICAM-1 and sVCAM-1. <i>Schizophrenia Research</i> , 2021, 228, 316-318.	2.0	2
27	The Class II Histone Deacetylase Hypothesis of Addiction. <i>Biological Psychiatry</i> , 2018, 84, 165-166.	1.3	1