

Ingvild Saltvedt

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6175065/publications.pdf>

Version: 2024-02-01

92
papers

6,806
citations

145106

33
h-index

87275

74
g-index

107
all docs

107
docs citations

107
times ranked

10418
citing authors

#	ARTICLE	IF	CITATIONS
1	Geriatric assessment with management for older patients with cancer receiving radiotherapy. Protocol of a Norwegian cluster-randomised controlled pilot study. <i>Journal of Geriatric Oncology</i> , 2022, 13, 363-373.	0.5	5
2	Estimation of recurrent atherosclerotic cardiovascular event risk in patients with established cardiovascular disease: the updated SMART2 algorithm. <i>European Heart Journal</i> , 2022, 43, 1715-1727.	1.0	40
3	Is Frailty Index a better predictor than pre-stroke modified Rankin Scale for neurocognitive outcomes 3-months post-stroke?. <i>BMC Geriatrics</i> , 2022, 22, 139.	1.1	5
4	Physical Performance and Cognition as Predictors of Instrumental Activities of Daily Living After Stroke: A Prospective Multicenter Cohort Study. <i>Archives of Physical Medicine and Rehabilitation</i> , 2022, 103, 1320-1326.	0.5	2
5	Cerebrospinal fluid catecholamines in Alzheimer's disease patients with and without biological disease. <i>Translational Psychiatry</i> , 2022, 12, 151.	2.4	16
6	New insights into the genetic etiology of Alzheimer's disease and related dementias. <i>Nature Genetics</i> , 2022, 54, 412-436.	9.4	700
7	Use of lipid-lowering therapy after ischaemic stroke and expected benefit from intensification of treatment. <i>Open Heart</i> , 2022, 9, e001972.	0.9	2
8	Brain Morphometric Correlates of Depressive Symptoms among Patients with and without Dementia. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2022, 12, 107-114.	0.6	1
9	The relationship of acute delirium with cognitive and psychiatric symptoms after stroke: a longitudinal study. <i>BMC Neurology</i> , 2022, 22, .	0.8	6
10	Using Polygenic Hazard Scores to Predict Age at Onset of Alzheimer's Disease in Nordic Populations. <i>Journal of Alzheimer's Disease</i> , 2022, 88, 1533-1544.	1.2	3
11	Vascular risk factor control and adherence to secondary preventive medication after ischaemic stroke. <i>Journal of Internal Medicine</i> , 2021, 289, 355-368.	2.7	11
12	Test Accuracy of the Montreal Cognitive Assessment in Screening for Early Poststroke Neurocognitive Disorder. <i>Stroke</i> , 2021, 52, 317-320.	1.0	25
13	Clinically accessible neuroimaging predictors of post-stroke neurocognitive disorder: a prospective observational study. <i>BMC Neurology</i> , 2021, 21, 89.	0.8	18
14	Associations between post-stroke motor and cognitive function: a cross-sectional study. <i>BMC Geriatrics</i> , 2021, 21, 103.	1.1	46
15	Orthogeriatrics prevents functional decline in hip fracture patients: report from two randomized controlled trials. <i>BMC Geriatrics</i> , 2021, 21, 208.	1.1	9
16	Association between in-hospital frailty and health-related quality of life after stroke: the Nor-COAST study. <i>BMC Neurology</i> , 2021, 21, 100.	0.8	15
17	Abstract P66: Prediction of Early Post-Stroke Major Neurocognitive Disorder Using Support Vector Machines. <i>Stroke</i> , 2021, 52, .	1.0	0
18	The delirium screening tool 4AT in routine clinical practice: prediction of mortality, sensitivity and specificity. <i>European Geriatric Medicine</i> , 2021, 12, 793-800.	1.2	5

#	ARTICLE	IF	CITATIONS
19	Common variants in Alzheimer's disease and risk stratification by polygenic risk scores. <i>Nature Communications</i> , 2021, 12, 3417.	5.8	140
20	Pre-stroke cognitive impairment is associated with vascular imaging pathology: a prospective observational study. <i>BMC Geriatrics</i> , 2021, 21, 362.	1.1	9
21	Genome-wide association identifies the first risk loci for psychosis in Alzheimer disease. <i>Molecular Psychiatry</i> , 2021, 26, 5797-5811.	4.1	30
22	Predicting the Emergence of Major Neurocognitive Disorder Within Three Months After a Stroke. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 705889.	1.7	7
23	Investigating novel biomarkers of immune activation and modulation in the context of sedentary behaviour: a multicentre prospective ischemic stroke cohort study. <i>BMC Neurology</i> , 2021, 21, 318.	0.8	3
24	The Impact of Vascular Risk Factors on Post-stroke Cognitive Impairment: The Nor-COAST Study. <i>Frontiers in Neurology</i> , 2021, 12, 678794.	1.1	10
25	Risk Stratification in Patients with Ischemic Stroke and Residual Cardiovascular Risk with Current Secondary Prevention. <i>Clinical Epidemiology</i> , 2021, Volume 13, 813-823.	1.5	9
26	A genome-wide association study with 1,126,563 individuals identifies new risk loci for Alzheimer's disease. <i>Nature Genetics</i> , 2021, 53, 1276-1282.	9.4	430
27	Neuroimaging improves the prediction of post-stroke major neurocognitive disorder. <i>Journal of the Neurological Sciences</i> , 2021, 429, 118294.	0.3	0
28	Gait, physical function, and physical activity in three groups of home-dwelling older adults with different severity of cognitive impairment – a cross-sectional study. <i>BMC Geriatrics</i> , 2021, 21, 670.	1.1	10
29	Neopterin and kynurenic acid as predictors of stroke recurrence and mortality: a multicentre prospective cohort study on biomarkers of inflammation measured three months after ischemic stroke. <i>BMC Neurology</i> , 2021, 21, 476.	0.8	1
30	The Association Between Circulating Inflammatory Markers and the Progression of Alzheimer Disease in Norwegian Memory Clinic Patients With Mild Cognitive Impairment or Dementia. <i>Alzheimer Disease and Associated Disorders</i> , 2020, 34, 47-53.	0.6	7
31	Post-stroke Cognitive Impairment – Impact of Follow-Up Time and Stroke Subtype on Severity and Cognitive Profile: The Nor-COAST Study. <i>Frontiers in Neurology</i> , 2020, 11, 699.	1.1	51
32	Is long-bout sedentary behaviour associated with long-term glucose levels 3 months after acute ischaemic stroke? A prospective observational cohort study. <i>BMJ Open</i> , 2020, 10, e037475.	0.8	2
33	Cerebrospinal fluid sTREM2 in Alzheimer's disease: comparisons between clinical presentation and AT classification. <i>Scientific Reports</i> , 2020, 10, 15886.	1.6	23
34	A high cerebrospinal fluid soluble TREM2 level is associated with slow clinical progression of Alzheimer's disease. <i>Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring</i> , 2020, 12, e12128.	1.2	16
35	<p>The Risk of Selection Bias in a Clinical Multi-Center Cohort Study. Results from the Norwegian Cognitive Impairment After Stroke (Nor-COAST) Study</p>. <i>Clinical Epidemiology</i> , 2020, Volume 12, 1327-1336.	1.5	27
36	Client, caregiver, volunteer, and therapist views on a voluntary supported group exercise programme for older adults with dementia. <i>BMC Geriatrics</i> , 2020, 20, 235.	1.1	5

#	ARTICLE	IF	CITATIONS
37	Impact of different methods defining post-stroke neurocognitive disorder: The Nor-COAST study. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2020, 6, e12000.	1.8	32
38	The Human Brain Representation of Odor Identification in Amnesic Mild Cognitive Impairment and Alzheimer's Dementia of Mild Degree. <i>Frontiers in Neurology</i> , 2020, 11, 607566.	1.1	15
39	Moderately increased albuminuria, chronic kidney disease and incident dementia: the HUNT study. <i>BMC Nephrology</i> , 2019, 20, 261.	0.8	23
40	Examining the association between genetic liability for schizophrenia and psychotic symptoms in Alzheimer's disease. <i>Translational Psychiatry</i> , 2019, 9, 273.	2.4	36
41	Modifiable factors affecting older patients' quality of life and physical function during cancer treatment. <i>Journal of Geriatric Oncology</i> , 2019, 10, 904-912.	0.5	38
42	Delirium motor subtypes and prognosis in hospitalized geriatric patients – A prospective observational study. <i>Journal of Psychosomatic Research</i> , 2019, 122, 24-28.	1.2	13
43	GBA and APOE ϵ 4 associate with sporadic dementia with Lewy bodies in European genome wide association study. <i>Scientific Reports</i> , 2019, 9, 7013.	1.6	53
44	Interdisciplinary care of hip fractures.. <i>Best Practice and Research in Clinical Rheumatology</i> , 2019, 33, 205-226.	1.4	24
45	My husband is not ill; he has memory loss - caregivers' perspectives on health care services for persons with dementia. <i>BMC Geriatrics</i> , 2019, 19, 75.	1.1	14
46	Motor activity across delirium motor subtypes in geriatric patients assessed using body-worn sensors: a Norwegian cross-sectional study. <i>BMJ Open</i> , 2019, 9, e026401.	0.8	10
47	Short and long-term clinical effectiveness and cost-effectiveness of a late-phase community-based balance and gait exercise program following hip fracture. <i>The EVA-Hip Randomised Controlled Trial. PLoS ONE</i> , 2019, 14, e0224971.	1.1	25
48	Team Approach: Multidisciplinary Treatment of Hip Fractures in Elderly Patients. <i>JBS Reviews</i> , 2019, 7, e6-e6.	0.8	5
49	Cortisol levels among older people with and without depression and dementia. <i>International Psychogeriatrics</i> , 2019, 31, 597-601.	0.6	19
50	Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. <i>Nature Genetics</i> , 2019, 51, 404-413.	9.4	1,625
51	Progression of Alzheimer's Disease: A Longitudinal Study in Norwegian Memory Clinics. <i>Journal of Alzheimer's Disease</i> , 2018, 61, 1221-1232.	1.2	44
52	The Norwegian Cognitive impairment after stroke study (Nor-COAST): study protocol of a multicentre, prospective cohort study. <i>BMC Neurology</i> , 2018, 18, 193.	0.8	39
53	Environmental factors and risk of delirium in geriatric patients: an observational study. <i>BMC Geriatrics</i> , 2018, 18, 282.	1.1	14
54	Meta-analysis of Alzheimer's disease on 9,751 samples from Norway and IGAP study identifies four risk loci. <i>Scientific Reports</i> , 2018, 8, 18088.	1.6	47

#	ARTICLE	IF	CITATIONS
55	Association between vascular comorbidity and progression of Alzheimer's disease: a two-year observational study in Norwegian memory clinics. <i>BMC Geriatrics</i> , 2018, 18, 120.	1.1	26
56	The association of high sensitivity C-reactive protein and incident Alzheimer disease in patients 60 years and older: The HUNT study, Norway. <i>Immunity and Ageing</i> , 2018, 15, 4.	1.8	27
57	Association between blood pressure and Alzheimer disease measured up to 27 years prior to diagnosis: the HUNT Study. <i>Alzheimer's Research and Therapy</i> , 2017, 9, 37.	3.0	66
58	Comprehensive geriatric assessment for older adults admitted to hospital. <i>The Cochrane Library</i> , 2017, CD006211.	1.5	383
59	Patterns of drug prescriptions in an orthogeriatric ward as compared to orthopaedic ward: results from the Trondheim Hip Fracture Trial—a randomised clinical trial. <i>European Journal of Clinical Pharmacology</i> , 2017, 73, 937-947.	0.8	12
60	Visual Evaluation of Medial Temporal Lobe Atrophy as a Clinical Marker of Conversion from Mild Cognitive Impairment to Dementia and for Predicting Progression in Patients with Mild Cognitive Impairment and Mild Alzheimer's Disease. <i>Dementia and Geriatric Cognitive Disorders</i> , 2017, 44, 12-24.	0.7	13
61	Trajectories of depressive symptoms and their relationship to the progression of dementia. <i>Journal of Affective Disorders</i> , 2017, 222, 146-152.	2.0	39
62	Factors that influence the levels of cerebrospinal fluid biomarkers in memory clinic patients. <i>BMC Geriatrics</i> , 2017, 17, 210.	1.1	9
63	MRI-assessed atrophy subtypes in Alzheimer's disease and the cognitive reserve hypothesis. <i>PLoS ONE</i> , 2017, 12, e0186595.	1.1	51
64	One-year health and care costs after hip fracture for home-dwelling elderly patients in Norway: Results from the Trondheim Hip Fracture Trial. <i>Scandinavian Journal of Public Health</i> , 2016, 44, 791-798.	1.2	22
65	Who benefits from orthogeriatric treatment? Results from the Trondheim hip-fracture trial. <i>BMC Geriatrics</i> , 2016, 16, 49.	1.1	38
66	The long-term effect of comprehensive geriatric care on gait after hip fracture: the Trondheim Hip Fracture Trial—a randomised controlled trial. <i>Osteoporosis International</i> , 2016, 27, 933-942.	1.3	55
67	The long-term effect of being treated in a geriatric ward compared to an orthopaedic ward on six measures of free-living physical behavior 4 and 12 months after a hip fracture - a randomised controlled trial. <i>BMC Geriatrics</i> , 2015, 15, 160.	1.1	28
68	Comprehensive geriatric care for patients with hip fractures: a prospective, randomised, controlled trial. <i>Lancet, The</i> , 2015, 385, 1623-1633.	6.3	449
69	Alcohol consumption and risk of dementia up to 27 years later in a large, population-based sample: the HUNT study, Norway. <i>European Journal of Epidemiology</i> , 2015, 30, 1049-1056.	2.5	72
70	Effectiveness of Task Specific Gait and Balance Exercise 4 Months After Hip Fracture: Protocol of a Randomized Controlled Trial—the Eva Hip Study. <i>Physiotherapy Research International</i> , 2015, 20, 87-99.	0.7	9
71	Cohort Profile: The Health and Memory Study (HMS): a dementia cohort linked to the HUNT study in Norway. <i>International Journal of Epidemiology</i> , 2014, 43, 1759-1768.	0.9	21
72	Physical Behavior and Function Early After Hip Fracture Surgery in Patients Receiving Comprehensive Geriatric Care or Orthopedic Care—A Randomized Controlled Trial. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2014, 69A, 338-345.	1.7	84

#	ARTICLE	IF	CITATIONS
73	The brain structural and cognitive basis of odor identification deficits in mild cognitive impairment and Alzheimer's disease. <i>BMC Neurology</i> , 2014, 14, 168.	0.8	64
74	Mental health and wellbeing in spouses of persons with dementia: the Nord-Trøndelag health study. <i>BMC Public Health</i> , 2014, 14, 413.	1.2	36
75	The effect of a pre- and postoperative orthogeriatric service on cognitive function in patients with hip fracture: randomized controlled trial (Oslo Orthogeriatric Trial). <i>BMC Medicine</i> , 2014, 12, 63.	2.3	134
76	The effect of a pre- and post-operative orthogeriatric service on cognitive function in patients with hip fracture. The protocol of the Oslo Orthogeriatrics Trial. <i>BMC Geriatrics</i> , 2012, 12, 36.	1.1	40
77	Development and delivery of patient treatment in the Trondheim Hip Fracture Trial. A new geriatric in-hospital pathway for elderly patients with hip fracture. <i>BMC Research Notes</i> , 2012, 5, 355.	0.6	37
78	Demens og nevropsykiatriske symptomer hos sykehjemspasienter i Nord-Trøndelag. <i>Tidsskrift for Den Norske Lægeforening</i> , 2012, 132, 1956-1959.	0.2	35
79	Effect of in-hospital comprehensive geriatric assessment (CGA) in older people with hip fracture. The protocol of the Trondheim Hip Fracture Trial. <i>BMC Geriatrics</i> , 2011, 11, 18.	1.1	47
80	Unwanted incidents during transition of geriatric patients from hospital to home: a prospective observational study. <i>BMC Health Services Research</i> , 2010, 10, 1.	0.9	442
81	Validation of Doloplus-2 among nonverbal nursing home patients - an evaluation of Doloplus-2 in a clinical setting. <i>BMC Geriatrics</i> , 2010, 10, 9.	1.1	26
82	Risk-reducing effect of education in Alzheimer's disease. <i>International Journal of Geriatric Psychiatry</i> , 2008, 23, 1156-1162.	1.3	73
83	APOE ϵ 4 lowers age at onset and is a high risk factor for Alzheimer's disease; A case control study from central Norway. <i>BMC Neurology</i> , 2008, 8, 9.	0.8	196
84	Long-chain n^{-3} fatty acids and mortality in elderly patients. <i>American Journal of Clinical Nutrition</i> , 2008, 88, 722-729.	2.2	26
85	Doloplus-2, a valid tool for behavioural pain assessment?. <i>BMC Geriatrics</i> , 2007, 7, 29.	1.1	41
86	Randomised Trial of In-Hospital Geriatric Intervention: Impact on Function and Morale. <i>Gerontology</i> , 2006, 52, 223-230.	1.4	24
87	Patterns of drug prescription in a geriatric evaluation and management unit as compared with the general medical wards: a randomised study. <i>European Journal of Clinical Pharmacology</i> , 2005, 61, 921-928.	0.8	64
88	The Norwegian Doloplus-2, a tool for behavioural pain assessment: translation and pilot-validation in nursing home patients with cognitive impairment. <i>Palliative Medicine</i> , 2005, 19, 411-417.	1.3	45
89	Acute geriatric intervention increases the number of patients able to live at home. A prospective randomized study. <i>Aging Clinical and Experimental Research</i> , 2004, 16, 300-306.	1.4	34
90	Which cancer patients die in nursing homes? Quality of life, medical and sociodemographic characteristics. <i>Palliative Medicine</i> , 2003, 17, 433-444.	1.3	51

#	ARTICLE	IF	CITATIONS
91	Reduced Mortality in Treating Acutely Sick, Frail Older Patients in a Geriatric Evaluation and Management Unit. A Prospective Randomized Trial. Journal of the American Geriatrics Society, 2002, 50, 792-798.	1.3	162
92	Longitudinal Brain Changes After Stroke and the Association With Cognitive Decline. Frontiers in Neurology, 0, 13, .	1.1	7