Patrik Danielson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6179385/publications.pdf

Version: 2024-02-01

33 papers 1,110 citations

430874 18 h-index 434195 31 g-index

34 all docs

34 docs citations

34 times ranked 1199 citing authors

#	Article	IF	Citations
1	Microstructured collagen films for 3D corneal stroma modelling. Connective Tissue Research, 2022, 63, 443-452.	2.3	2
2	Activation of NF- \hat{l}° B signaling via cytosolic mitochondrial RNA sensing in kerotocytes with mitochondrial DNA common deletion. Scientific Reports, 2021, 11, 7360.	3.3	6
3	Sustained Release of TPCAâ€1 from Silk Fibroin Hydrogels Preserves Keratocyte Phenotype and Promotes Corneal Regeneration by Inhibiting Interleukinâ€1 <i>β</i> Signaling. Advanced Healthcare Materials, 2020, 9, e2000591.	7.6	26
4	Regulation of Keratocyte Phenotype and Cell Behavior by Substrate Stiffness. ACS Biomaterials Science and Engineering, 2020, 6, 5162-5171.	5.2	22
5	Acetylcholine decreases formation of myofibroblasts and excessive extracellular matrix production in an in vitro human corneal fibrosis model. Journal of Cellular and Molecular Medicine, 2020, 24, 4850-4862.	3.6	9
6	Substance P induces fibrotic changes through activation of the RhoA/ROCK pathway in an in vitro human corneal fibrosis model. Journal of Molecular Medicine, 2019, 97, 1477-1489.	3.9	13
7	Induction of Fibroblast Senescence During Mouse Corneal Wound Healing. , 2019, 60, 3669.		34
8	Mechanical stress potentiates the differentiation of periodontal ligament stem cells into keratocytes. British Journal of Ophthalmology, 2018, 102, 562-569.	3.9	18
9	Ascorbic Acid Promotes the Stemness of Corneal Epithelial Stem/Progenitor Cells and Accelerates Epithelial Wound Healing in the Cornea. Stem Cells Translational Medicine, 2017, 6, 1356-1365.	3.3	53
10	Surface Topography and Mechanical Strain Promote Keratocyte Phenotype and Extracellular Matrix Formation in a Biomimetic 3D Corneal Model. Advanced Healthcare Materials, 2017, 6, 1601238.	7.6	38
11	Glutamate signaling through the NMDA receptor reduces the expression of scleraxis in plantaris tendon derived cells. BMC Musculoskeletal Disorders, 2017, 18, 218.	1.9	7
12	Corneal Epithelium-Derived Neurotrophic Factors Promote Nerve Regeneration., 2017, 58, 4695.		70
13	Substance P and patterned silk biomaterial stimulate periodontal ligament stem cells to form corneal stroma in a bioengineered three-dimensional model. Stem Cell Research and Therapy, 2017, 8, 260.	5.5	14
14	The effects of substance P and acetylcholine on human tenocyte proliferation converge mechanistically via TGF- \hat{l}^21 . PLoS ONE, 2017, 12, e0174101.	2.5	16
15	Antiapoptotic Effect of Acetylcholine in Fas-Induced Apoptosis in Human Keratocytes., 2016, 57, 5892.		8
16	Ciliary Neurotrophic Factor Promotes the Migration of Corneal Epithelial Stem/progenitor Cells by Up-regulation of MMPs through the Phosphorylation of Akt. Scientific Reports, 2016, 6, 25870.	3.3	35
17	Transforming Growth Factor Beta 1 Modulates the Functional Expression of the Neurokinin-1 Receptor in Human Keratocytes. Current Eye Research, 2016, 41, 1035-1043.	1.5	7
18	Substance P Enhances Keratocyte Migration and Neutrophil Recruitment through Interleukin-8. Molecular Pharmacology, 2016, 89, 215-225.	2.3	56

#	Article	IF	Citations
19	Expression Profiles of Neuropeptides, Neurotransmitters, and Their Receptors in Human Keratocytes In Vitro and In Situ. PLoS ONE, 2015, 10, e0134157.	2.5	41
20	Acetylcholine enhances keratocyte proliferation through muscarinic receptor activation. International Immunopharmacology, 2015, 29, 57-62.	3.8	16
21	Substance P reduces TNF-î±-induced apoptosis in human tenocytes through NK-1 receptor stimulation. British Journal of Sports Medicine, 2014, 48, 1414-1420.	6.7	28
22	Substance P Promotes Diabetic Corneal Epithelial Wound Healing Through Molecular Mechanisms Mediated via the Neurokinin-1 Receptor. Diabetes, 2014, 63, 4262-4274.	0.6	141
23	93â€Evolving Inflammatory Cell Populations In The Overused Rabbit Achilles Tendon. British Journal of Sports Medicine, 2014, 48, A60.2-A61.	6.7	1
24	Substance P enhances collagen remodeling and MMPâ€3 expression by human tenocytes. Journal of Orthopaedic Research, 2013, 31, 91-98.	2.3	34
25	Aktâ€mediated antiâ€apoptotic effects of substance P in Antiâ€Fasâ€induced apoptosis of human tenocytes. Journal of Cellular and Molecular Medicine, 2013, 17, 723-733.	3.6	24
26	An Emerging Role for Angiogenesis in Tendinopathy. European Musculoskeletal Review, 2009, 4, 75-76.	0.0	5
27	Marked sympathetic component in the perivascular innervation of the dorsal paratendinous tissue of the patellar tendon in arthroscopically treated tendinosis patients. Knee Surgery, Sports Traumatology, Arthroscopy, 2008, 16, 621-626.	4.2	50
28	Presence of substance P and the neurokinin-1 receptor in tenocytes of the human Achilles tendon. Regulatory Peptides, 2008, 150, 81-87.	1.9	80
29	Extensive expression of markers for acetylcholine synthesis and of M2 receptors in tenocytes in therapy-resistant chronic painful patellar tendon tendinosis — a pilot study. Life Sciences, 2007, 80, 2235-2238.	4.3	34
30	Studies on the importance of sympathetic innervation, adrenergic receptors, and a possible local catecholamine production in the development of patellar tendinopathy (tendinosis) in man. Microscopy Research and Technique, 2007, 70, 310-324.	2.2	56
31	In situ hybridization studies confirming recent findings of the existence of a local nonneuronal catecholamine production in human patellar tendinosis. Microscopy Research and Technique, 2007, 70, 908-911.	2.2	41
32	Distribution of general (PGP 9.5) and sensory (substance P/CGRP) innervations in the human patellar tendon. Knee Surgery, Sports Traumatology, Arthroscopy, 2006, 14, 125-132.	4.2	55
33	Immunohistochemical and histochemical findings favoring the occurrence of autocrine/paracrine as well as nerveâ€related cholinergic effects in chronic painful patellar tendon tendinosis. Microscopy Research and Technique, 2006, 69, 808-819.	2.2	69