## Frederic Fumeron

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6183870/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Polymorphism in the fractalkine receptor CX3CR1 as a genetic risk factor for coronary artery disease. Blood, 2001, 97, 1925-1928.	0.6	314
2	Alcohol intake modulates the effect of a polymorphism of the cholesteryl ester transfer protein gene on plasma high density lipoprotein and the risk of myocardial infarction Journal of Clinical Investigation, 1995, 96, 1664-1671.	3.9	272
3	New loci for body fat percentage reveal link between adiposity and cardiometabolic disease risk. Nature Communications, 2016, 7, 10495.	5.8	245
4	Predicting Diabetes: Clinical, Biological, and Genetic Approaches. Diabetes Care, 2008, 31, 2056-2061.	4.3	215
5	Extensive association analysis between the CETP gene and coronary heart disease phenotypes reveals several putative functional polymorphisms and gene-environment interaction. Genetic Epidemiology, 2000, 19, 64-80.	0.6	205
6	Identification of novel risk loci for restless legs syndrome in genome-wide association studies in individuals of European ancestry: a meta-analysis. Lancet Neurology, The, 2017, 16, 898-907.	4.9	191
7	Adiponectin Gene Polymorphisms and Adiponectin Levels Are Independently Associated With the Development of Hyperglycemia During a 3-Year Period: The Epidemiologic Data on the Insulin Resistance Syndrome Prospective Study. Diabetes, 2004, 53, 1150-1157.	0.3	185
8	Association of the G-2548A polymorphism in the 5′ region of the LEP gene with overweight. Annals of Human Genetics, 2000, 64, 391-394.	0.3	178
9	Plasma Dihydroceramides Are Diabetes Susceptibility Biomarker Candidates in Mice and Humans. Cell Reports, 2017, 18, 2269-2279.	2.9	168
10	Novel polymorphisms in the 5' region of the LEP gene: association with leptin levels and response to low-calorie diet in human obesity. Diabetes, 1998, 47, 487-489.	0.3	156
11	ACDC/Adiponectin Polymorphisms Are Associated With Severe Childhood and Adult Obesity. Diabetes, 2006, 55, 545-550.	0.3	154
12	Impact of Common Type 2 Diabetes Risk Polymorphisms in the DESIR Prospective Study. Diabetes, 2008, 57, 244-254.	0.3	146
13	Association of the G-2548A polymorphism in the 5' region of the LEP gene with overweight. Annals of Human Genetics, 2000, 64, 391-394.	0.3	145
14	Dairy Consumption and the Incidence of Hyperglycemia and the Metabolic Syndrome. Diabetes Care, 2011, 34, 813-817.	4.3	136
15	Lipoprotein lipase gene polymorphisms: associations with myocardial infarction and lipoprotein levels, the ECTIM study. Etude Cas Témoin sur l'Infarctus du Myocarde Journal of Lipid Research, 1996, 36, 2141-2146.	2.0	125
16	Ferritin and Transferrin Are Both Predictive of the Onset of Hyperglycemia in Men and Women Over 3 Years: The Data from an Epidemiological Study on the Insulin Resistance Syndrome (DESIR) study. Diabetes Care, 2006, 29, 2090-2094.	4.3	123
17	Ferritin and Transferrin Are Associated With Metabolic Syndrome Abnormalities and Their Change Over Time in a General Population: Data from an Epidemiological Study on the Insulin Resistance Syndrome (DESIR). Diabetes Care, 2007, 30, 1795-1801.	4.3	121
18	Nine-year incident diabetes is predicted by fatty liver indices: the French D.E.S.I.R. study. BMC Gastroenterology, 2010, 10, 56.	0.8	120

#	Article	IF	CITATIONS
19	Comparison Between Copeptin and Vasopressin in a Population From the Community and in People With Chronic Kidney Disease. Journal of Clinical Endocrinology and Metabolism, 2014, 99, 4656-4663.	1.8	110
20	Lipoprotein lipase gene polymorphisms: associations with myocardial infarction and lipoprotein levels, the ECTIM study. Etude Cas Témoin sur l'Infarctus du Myocarde. Journal of Lipid Research, 1995, 36, 2141-6.	2.0	104
21	Increased plasma adiponectin concentrations are associated with microangiopathy in type 1 diabetic subjects. Diabetologia, 2005, 48, 1088-1092.	2.9	99
22	Endogenous opiates and energy balance. Science, 1982, 215, 1536-1538.	6.0	98
23	HLA and longevity. Tissue Antigens, 1982, 19, 168-173.	1.0	92
24	Bardet-Biedl Syndrome Gene Variants Are Associated With Both Childhood and Adult Common Obesity in French Caucasians. Diabetes, 2006, 55, 2876-2882.	0.3	87
25	LEP R gene polymorphisms: associations with overweight, fat mass and response to diet in women. European Journal of Clinical Investigation, 2001, 31, 398-404.	1.7	79
26	Serotonin Transporter Gene Polymorphism and Myocardial Infarction. Circulation, 2002, 105, 2943-2945.	1.6	77
27	Association of a functional 5-HT transporter gene polymorphism with anorexia nervosa and food intake. Molecular Psychiatry, 2001, 6, 9-10.	4.1	75
28	Plasma Copeptin and Renal Outcomes in Patients With Type 2 Diabetes and Albuminuria. Diabetes Care, 2013, 36, 3639-3645.	4.3	73
29	5-HT2A receptor gene polymorphism is associated with food and alcohol intake in obese people. International Journal of Obesity, 2000, 24, 920-924.	1.6	72
30	Analysis of novel risk loci for type 2 diabetes in a general French population: the D.E.S.I.R. study. Journal of Molecular Medicine, 2008, 86, 341-348.	1.7	71
31	Influence of the saci apo C-III gene polymorphism on the insulin response to an oral glucose tolerance test in obese subjects. Atherosclerosis, 1999, 144, 109-110.	0.4	70
32	Prognostic Value of the Insertion/Deletion Polymorphism of the <i>ACE</i> Gene in Type 2 Diabetic Subjects. Diabetes Care, 2008, 31, 1847-1852.	4.3	66
33	Weight change and changes in the metabolic syndrome as the French population moves towards overweight: The D.E.S.I.R. Cohort. International Journal of Epidemiology, 2006, 35, 190-196.	0.9	63
34	The PPARG Pro12Ala Polymorphism Is Associated With a Decreased Risk of Developing Hyperglycemia Over 6 Years and Combines With the Effect of the APM1 G-11391A Single Nucleotide Polymorphism: The Data From an Epidemiological Study on the Insulin Resistance Syndrome (DESIR) Study. Diabetes, 2006, 55, 1157-1162.	0.3	63
35	Dairy product consumption, calcium intakes, and metabolic syndrome–related factors over 5 years in the STANISLAS study. Nutrition, 2013, 29, 519-524.	1.1	60
36	GUESS-ing Polygenic Associations with Multiple Phenotypes Using a GPU-Based Evolutionary Stochastic Search Algorithm. PLoS Genetics, 2013, 9, e1003657.	1.5	58

#	Article	IF	CITATIONS
37	Plasma Copeptin, <i>AVP</i> Gene Variants, and Incidence of Type 2 Diabetes in a Cohort From the Community. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 2432-2439.	1.8	58
38	nâ^'3 Polyunsaturated fatty acids raise low-density lipoproteins, high-density lipoprotein 2, and plasminogen-activator inhibitor in healthy young men. American Journal of Clinical Nutrition, 1991, 54, 118-122.	2.2	56
39	Polymorphisms of the Tissue Factor Pathway Inhibitor (TFPI) Gene in Patients With Acute Coronary Syndromes and in Healthy Subjects. Arteriosclerosis, Thrombosis, and Vascular Biology, 1999, 19, 862-869.	1.1	55
40	A New T-287C Polymorphism in the 5' Regulatory Region of the Tissue Factor Pathway Inhibitor Gene. Thrombosis and Haemostasis, 2000, 84, 244-249.	1.8	53
41	Lowering of HDL2-cholesterol and lipoprotein A-1 particle levels by increasing the ratio of polyunsaturated to saturated fatty acids. American Journal of Clinical Nutrition, 1991, 53, 655-659.	2.2	51
42	Dietary fat intake and polymorphisms at the PPARG locus modulate BMI and type 2 diabetes risk in the D.E.S.I.R. prospective study. International Journal of Obesity, 2012, 36, 218-224.	1.6	51
43	Plasma Copeptin, Kidney Outcomes, Ischemic Heart Disease, and All-Cause Mortality in People With Long-standing Type 1 Diabetes. Diabetes Care, 2016, 39, 2288-2295.	4.3	51
44	<i>ABCA1</i> Single Nucleotide Polymorphisms on Highâ€Density Lipoproteinâ€Cholesterol and Overweight: the D.E.S.I.R. Study. Obesity, 2006, 14, 1874-1879.	1.5	49
45	The loss-of-function PCSK9 p.R46L genetic variant does not alter glucose homeostasis. Diabetologia, 2015, 58, 2051-2055.	2.9	49
46	Presence of the French Canadian Deletion in a French Patient with Familial Hypercholesterolemia. New England Journal of Medicine, 1992, 326, 69-69.	13.9	48
47	Plasma extracellular superoxide dismutase concentration, allelic variations in the SOD3 gene and risk of myocardial infarction and all-cause mortality in people with type 1 and type 2 diabetes. Cardiovascular Diabetology, 2015, 14, 845.	2.7	47
48	The Functional c2G>C Variant of the Mineralocorticoid Receptor Modulates Blood Pressure, Renin, and Aldosterone Levels. Hypertension, 2010, 56, 995-1002.	1.3	46
49	Dairy Products and the Metabolic Syndrome in a Prospective Study, DESIR. Journal of the American College of Nutrition, 2011, 30, 454S-463S.	1.1	45
50	Intrauterine Growth Retardation, Perinatal Death, and Maternal Homocysteine Levels. New England Journal of Medicine, 1992, 326, 69-70.	13.9	44
51	The Common â^'866G>A Variant in the Promoter of UCP2 Is Associated With Decreased Risk of Coronary Artery Disease in Type 2 Diabetic Men. Diabetes, 2008, 57, 1063-1068.	0.3	44
52	Plasma Copeptin and Decline in Renal Function in a Cohort from the Community: The Prospective D.E.S.I.R. Study. American Journal of Nephrology, 2015, 42, 107-114.	1.4	43
53	Death, end-stage renal disease and renal function decline in patients with diabetic nephropathy in French cohorts of type 1 and type 2 diabetes. Diabetologia, 2016, 59, 208-216.	2.9	42
54	The â^'33T → C Polymorphism in Intron 7 of the TFPI Gene Influences the Risk of Venous Thromboembolism, Independently of the Factor V Leiden and Prothrombin Mutations. Thrombosis and Haemostasis, 2002, 88, 195-199.	1.8	41

#	Article	IF	CITATIONS
55	High Baseline Insulin Levels Associated With 6-Year Incident Observed Sleep Apnea. Diabetes Care, 2010, 33, 1044-1049.	4.3	41
56	Association of Birth Weight With Type 2 Diabetes and Glycemic Traits. JAMA Network Open, 2019, 2, e1910915.	2.8	41
57	Sex hormone-binding globulin predicts the incidence of hyperglycemia in women: interactions with adiponectin levels. European Journal of Endocrinology, 2009, 161, 81-85.	1.9	40
58	Allelic variations of the vitamin D receptor (VDR) gene are associated with increased risk of coronary artery disease in type 2 diabetics: The DIABHYCAR prospective study. Diabetes and Metabolism, 2013, 39, 263-270.	1.4	40
59	Relationships between common polymorphisms of adenosine triphosphate–binding cassette transporter A1 and high-density lipoprotein cholesterol and coronary heart disease in a population with type 2 diabetes mellitus. Metabolism: Clinical and Experimental, 2009, 58, 74-79.	1.5	39
60	Lack of association between genetic variations of apo A-l–C-III–A-IV gene cluster and myocardial infarction in a sample of European male: ECTIM study. Atherosclerosis, 1999, 145, 187-195.	0.4	38
61	Dynamic Changes in Renal Function Are Associated With Major Cardiovascular Events in Patients With Type 2 Diabetes. Diabetes Care, 2016, 39, 1259-1266.	4.3	38
62	Glutathione peroxidase-1 gene (GPX1) variants, oxidative stress and risk of kidney complications in people with type 1 diabetes. Metabolism: Clinical and Experimental, 2016, 65, 12-19.	1.5	37
63	Effect of a moderate alcohol intake on the lipoproteins of normotriglyceridemic obese subjects compared with normoponderal controls. Metabolism: Clinical and Experimental, 1992, 41, 856-861.	1.5	36
64	Allelic variations in superoxide dismutase-1 (SOD1) gene and renal and cardiovascular morbidity and mortality in type 2 diabetic subjects. Molecular Genetics and Metabolism, 2012, 106, 359-365.	0.5	36
65	Plasma copeptin, kidney disease, and risk for cardiovascular morbidity and mortality in two cohorts of type 2 diabetes. Cardiovascular Diabetology, 2018, 17, 110.	2.7	35
66	Beneficial effects of a moderate consumption of red wine on cellular cholesterol efflux in young men. Nutrition, Metabolism and Cardiovascular Diseases, 2000, 10, 63-9.	1.1	35
67	Allelic variations in superoxide dismutase-1 (SOD1) gene are associated with increased risk of diabetic nephropathy in type 1 diabetic subjects. Molecular Genetics and Metabolism, 2011, 104, 654-660.	0.5	34
68	Association of vitronectin and plasminogen activator inhibitor-1 levels with the risk of metabolic syndrome and type 2 diabetes mellitus. Thrombosis and Haemostasis, 2011, 106, 416-422.	1.8	34
69	Dairy Consumption and Body Mass Index Among Adults: Mendelian Randomization Analysis of 184802 Individuals from 25 Studies. Clinical Chemistry, 2018, 64, 183-191.	1.5	34
70	The lactase persistence genotype is associated with body mass index and dairy consumption in the D.E.S.I.R. study. Metabolism: Clinical and Experimental, 2013, 62, 1323-1329.	1.5	33
71	Association of ADIPOQ genetic variants and plasma adiponectin isoforms with the risk of incident renal events in type 2 diabetes. Nephrology Dialysis Transplantation, 2010, 25, 2231-2237.	0.4	32
72	Plasma copeptin and chronic kidney disease risk in 3 European cohorts from the general population. JCI Insight, 2018, 3, .	2.3	32

#	Article	IF	CITATIONS
73	Fine-scale human genetic structure in Western France. European Journal of Human Genetics, 2015, 23, 831-836.	1.4	31
74	Manganese Superoxide Dismutase (SOD2) Polymorphisms, Plasma Advanced Oxidation Protein Products (AOPP) Concentration and Risk of Kidney Complications in Subjects with Type 1 Diabetes. PLoS ONE, 2014, 9, e96916.	1,1	31
75	Polymorphism of the 5-HT2A receptor gene and food intakes in children and adolescents: the Stanislas Family Study. American Journal of Clinical Nutrition, 2005, 82, 467-470.	2.2	30
76	Association of apolipoprotein ɛ4 allele with hypertriglyceridemia in obesity. Clinical Genetics, 1988, 34, 258-264.	1.0	30
77	Adiponectin gene variants, adiponectin isoforms and cardiometabolic risk in type 2 diabetic patients. Journal of Diabetes Investigation, 2014, 5, 192-198.	1.1	30
78	Contribution of the low-frequency, loss-of-function p.R270H mutation in <i>FFAR4</i> ( <i>GPR120</i> ) to increased fasting plasma glucose levels. Journal of Medical Genetics, 2015, 52, 595-598.	1.5	29
79	Decreased insulin secretion and increased risk of type 2 diabetes associated with allelic variations of the WFS1 gene: the Data from Epidemiological Study on the Insulin Resistance Syndrome (DESIR) prospective study. Diabetologia, 2011, 54, 554-562.	2.9	28
80	Sexâ€dependent Associations of Leptin With Metabolic Syndrome–related Variables: The Stanislas Study. Obesity, 2010, 18, 196-201.	1.5	24
81	Evidence for distinct effects of GH and IGF-I in the metabolic syndrome. Diabetic Medicine, 2007, 24, 1012-1018.	1.2	23
82	Association Between Endothelin Receptor B Nonsynonymous Variants and Melanoma Risk. Journal of the National Cancer Institute, 2005, 97, 1297-1301.	3.0	22
83	Associations of the -344 T>C and the 3097 G>A Polymorphisms of CYP11B2 Gene With Hypertension, Type 2 Diabetes, and Metabolic Syndrome in a French Population. American Journal of Hypertension, 2010, 23, 660-667.	1.0	22
84	Low high density lipoproteina <sup>2</sup> concentrations in obese male subjects. Atherosclerosis, 1988, 73, 57-61.	0.4	21
85	HLA genotypes in familial Hodgkin's disease. Excess of HLA identical affected sibs. European Journal of Cancer, 1980, 16, 809-815.	1.0	20
86	Interindividual variability in the cholesterol-lowering effect of supplementation with plant sterols or stanols. Nutrition Reviews, 2017, 75, 134-145.	2.6	20
87	Hind III polymorphism of the lipoprotein lipase gene and plasma lipid response to low calorie diet. International Journal of Obesity, 1997, 21, 280-283.	1.6	19
88	Adiponectin gene and cardiovascular risk in type 2 diabetic patients: a review of evidences. Arquivos Brasileiros De Endocrinologia E Metabologia, 2007, 51, 153-159.	1.3	19
89	Potential risks associated with increased plasma plant-sterol levels. Diabetes and Metabolism, 2015, 41, 76-81.	1.4	19
90	Adiponectin Multimers and ADIPOQ T45G in Coronary Artery Disease in Caribbean Type 2 Diabetic Subjects of African Descent. Obesity, 2010, 18, 1466-1468.	1.5	18

#	Article	IF	CITATIONS
91	Apolipoprotein B signal peptide polymorphism and plasma LDL-cholesterol response to low-calorie diet. International Journal of Obesity, 2004, 28, 902-905.	1.6	17
92	Plasma proproteinâ€convertaseâ€subtilisin/kexin type 9 (PCSK9) and cardiovascular events in type 2 diabetes. Diabetes, Obesity and Metabolism, 2018, 20, 943-953.	2.2	17
93	The CETP <i>Taq</i> IB Polymorphism Is Associated With the Risk of Sudden Death in Type 2 Diabetic Patients. Diabetes Care, 2007, 30, 2863-2867.	4.3	16
94	Genetic Variability at the Six Transmembrane Protein of Prostate 2 Locus and the Metabolic Syndrome: The Data from an Epidemiological Study on the Insulin Resistance Syndrome (DESIR) Study. Journal of Clinical Endocrinology and Metabolism, 2010, 95, 2942-2947.	1.8	16
95	Comparative effects of several simple carbohydrates on erythrocyte insulin receptors in obese subjects. Pharmacology Biochemistry and Behavior, 1986, 25, 681-688.	1.3	15
96	Nonâ€severe hypoglycaemia is associated with weight gain in patients with type 1 diabetes: Results from the Diabetes Control and Complication Trial. Diabetes, Obesity and Metabolism, 2018, 20, 1289-1292.	2.2	15
97	Prognostic value of plasma MR-proADM vs NT-proBNP for heart failure in people with type 2 diabetes: the SURDIAGENE prospective study. Diabetologia, 2018, 61, 2643-2653.	2.9	15
98	New roles for prokineticin 2 in feeding behavior, insulin resistance and type 2 diabetes: Studies in mice and humans. Molecular Metabolism, 2019, 29, 182-196.	3.0	15
99	Plasma Copeptin and Risk of Lower-Extremity Amputation in Type 1 and Type 2 Diabetes. Diabetes Care, 2019, 42, 2290-2297.	4.3	15
100	Catalase activity, allelic variations in the catalase gene and risk of kidney complications in patients with type 1 diabetes. Diabetologia, 2013, 56, 2733-2742.	2.9	14
101	Allelic variations in the CYBA gene of NADPH oxidase and risk of kidney complications in patients with type 1 diabetes. Free Radical Biology and Medicine, 2015, 86, 16-24.	1.3	14
102	Antidepressant medication use and trajectories of fasting plasma glucose, glycated haemoglobin, β-cell function and insulin sensitivity: a 9-year longitudinal study of the D.E.S.I.R. cohort. International Journal of Epidemiology, 2015, 44, 1927-1940.	0.9	14
103	T-cadherin gene variants are associated with type 2 diabetes and the Fatty Liver Index in the French population. Diabetes and Metabolism, 2017, 43, 33-39.	1.4	14
104	Effects of acyl-coenzyme A binding protein (ACBP)/diazepam-binding inhibitor (DBI) on body mass index. Cell Death and Disease, 2021, 12, 599.	2.7	13
105	Lipoprotein lipase gene polymorphisms: associations with hypertriglyceridemia and body mass index in obese people. , 1995, 19, 270-4.		13
106	Polymorphism of the 5-HT2A receptor gene and food intakes in children and adolescents: the Stanislas Family Study. American Journal of Clinical Nutrition, 2005, 82, 467-470.	2.2	12
107	Plasma Adrenomedullin and Allelic Variation in the <i>ADM</i> Gene and Kidney Disease in People With Type 2 Diabetes. Diabetes, 2015, 64, 3262-3272.	0.3	12
108	Urinary Sodium Concentration Is an Independent Predictor of All-Cause and Cardiovascular Mortality in a Type 2 Diabetes Cohort Population. Journal of Diabetes Research, 2017, 2017, 1-10.	1.0	12

#	Article	IF	CITATIONS
109	Plasma Apelin and Risk of Type 2 Diabetes in a Cohort From the Community. Diabetes Care, 2020, 43, e15-e16.	4.3	12
110	Genetics of macrovascular complications in diabetes. Current Diabetes Reports, 2006, 6, 162-168.	1.7	11
111	ABCG8 polymorphisms and renal disease in type 2 diabetic patients. Metabolism: Clinical and Experimental, 2015, 64, 713-719.	1.5	11
112	A new T-287C polymorphism in the 5' regulatory region of the tissue factor pathway inhibitor gene. Association study of the T-287C and C-399T polymorphisms with coronary artery disease and plasma TFPI levels. Thrombosis and Haemostasis, 2000, 84, 244-9.	1.8	11
113	Longitudinal association of antidepressant medication use with metabolic syndrome: Results of a 9-year follow-up of the D.E.S.I.R. cohort study. Psychoneuroendocrinology, 2016, 74, 34-45.	1.3	10
114	Are the Same Clinical Risk Factors Relevant for Incident Diabetes Defined by Treatment, Fasting Plasma Glucose, and HbA1c?. Diabetes Care, 2011, 34, 957-959.	4.3	9
115	Interaction between GPR120 p.R270H loss-of-function variant and dietary fat intake on incident type 2 diabetes risk in the D.E.S.I.R. study. Nutrition, Metabolism and Cardiovascular Diseases, 2016, 26, 931-936.	1.1	9
116	Mendelian randomization analysis does not support causal associations of birth weight with hypertension risk and blood pressure in adulthood. European Journal of Epidemiology, 2020, 35, 685-697.	2.5	9
117	The -33T>C polymorphism in intron 7 of the TFPI gene influences the risk of venous thromboembolism, independently of the factor V Leiden and prothrombin mutations. Thrombosis and Haemostasis, 2002, 88, 195-9.	1.8	9
118	Lack of association between dietary alcohol and HDL-cholesterol concentrations in obese women. Atherosclerosis, 1990, 81, 119-125.	0.4	8
119	Lack of plasmatic beta-endorphin response to a gastronomic meal in healthy humans. Physiology and Behavior, 1991, 49, 1217-1221.	1.0	8
120	Hyperadiponectinemia Is Independent of Kidney Function, Diabetes Duration, and Control in Type 1 Diabetic Patients without Microangiopathy. Journal of Clinical Endocrinology and Metabolism, 2011, 96, E485-E487.	1.8	8
121	Association between HLA-B18 and the Familial-Obesity Syndrome. New England Journal of Medicine, 1981, 305, 645-645.	13.9	7
122	Intensifying glycaemic control with insulin reduces adiponectin and its HMW isoform moderately in type 2, but not in type 1, diabetes. Diabetes and Metabolism, 2011, 37, 259-261.	1.4	7
123	Effects of Slight Plasma Glucose Decrease on Glycosylated Hemoglobin in Healthy Subjects during Caloric Restriction. New England Journal of Medicine, 1985, 313, 958-959.	13.9	6
124	Patterns of Alcohol Consumption and Cardiovascular Risk in Northern Ireland and France. Annals of Epidemiology, 2007, 17, S75-S80.	0.9	6
125	Tissue kallikrein deficiency, insulin resistance, and diabetes in mouse and man. Journal of Endocrinology, 2014, 221, 297-308.	1.2	6
126	Polymorphisms of the lamina maturation pathway and their association with the metabolic syndrome: the DESIR prospective study. Journal of Molecular Medicine, 2010, 88, 193-201.	1.7	5

#	Article	IF	CITATIONS
127	Association of common variants in NPPA and NPPB with blood pressure does not translate into kidney damage in a general population study. Journal of Hypertension, 2010, 28, 1230-1233.	0.3	5
128	Association of ACACB gene polymorphism (rs2268388, G > A) with type 2 diabetes and end stage renal disease in Pakistani Punjabi population. Meta Gene, 2017, 12, 109-112.	0.3	3
129	T-cadherin gene variants are associated with nephropathy in subjects with type 1 diabetes. Nephrology Dialysis Transplantation, 2017, 32, 2144-2144.	0.4	3
130	Dairy consumption is associated with lower plasma dihydroceramides in women from the D.E.S.I.R. cohort. Diabetes and Metabolism, 2020, 46, 144-149.	1.4	3
131	Dietary fat, energy density and BMI: a case of a missing flower?. International Journal of Obesity, 1998, 22, 1032-1033.	1.6	2
132	Studies on an Isolated West Indies Population: I. Analysis of HLA Genotypes. Tissue Antigens, 1981, 17, 338-342.	1.0	2
133	T-cadherin gene variants are associated with nephropathy in subjects with type 1 diabetes. Nephrology Dialysis Transplantation, 2017, 32, 1987-1993.	0.4	2
134	Lipoprotein Lipase Gene Variants and Coronary Risk. Circulation, 1998, 97, 2588-2588.	1.6	1
135	Obésité : d'un syndrome monogénique exceptionnel aux interactions entre gènes multiples et environnement nutritionnel. Oleagineux Corps Gras Lipides, 2003, 10, 109-114.	0.2	1
136	Comment on: Park et al. Association of Serum Ferritin and the Development of Metabolic Syndrome in Middle-Aged Korean Men: A 5-Year Follow-up Study. Diabetes Care 2012;35:2521-2526. Diabetes Care, 2012, 35, e92-e92.	4.3	1
137	Phytostérols : un point sur les recommandations de l'ANSES. OCL - Oilseeds and Fats, Crops and Lipids, 2015, 22, D205.	0.6	1
138	Plasma total adiponectin and changes in renal function in a cohort from the community: the prospective Data from an Epidemiological Study on the Insulin Resistance Syndrome study. Nephrology Dialysis Transplantation, 2020, 36, 2058-2065.	0.4	1
139	Relationship between renal capacity to reabsorb glucose and renal status in patients with diabetes. Diabetes and Metabolism, 2020, 46, 488-495.	1.4	1
140	De l'obésité au diabète de type 2 : épidémiologie et physiopathologie. Sciences Des Aliments, 2005, 25, 339-347.	0.2	1
141	Circadian rhythm of energy intake and corpulence status in adults. , 1990, 14, 387-93.		1
142	BIRTHWEIGHT DIFFERENCE IN MONOZYGOUS TWINS FOLLOWED BY DIFFERENCE IN DEVELOPMENT OF BODY WEIGHT. Lancet, The, 1982, 320, 274.	6.3	0
143	Iron intake in obese menstruating women. Diabetes Research and Clinical Practice, 1990, 10, S287-S290.	1.1	0
144	Genetics of the human obesities. Obesite, 2013, 8, 22-33.	0.1	0

#	Article	IF	CITATIONS
145	Genetics of the Human Obesities. , 2013, , 351-372.		Ο
146	Génétique des obésités humaines. , 2013, , 359-380.		0
147	La T-cadhérine, troisième récepteur de l'adiponectine : structure et rÃ1e en santé humaine. Obesite, 2017, 12, 267-276.	0.1	Ο
148	L'adiponectine. Sciences Des Aliments, 2003, 23, 3-12.	0.2	0
149	Réponse aux lipides de l'alimentation et profil génétique. Sciences Des Aliments, 2008, 28, 143-149.	0.2	Ο
150	Xmn1 restriction polymorphism of apolipoprotein AI gene and lipoprotein levels in obesity. , 1994, 18, 460-4.		0