

Frederic Fumeron

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6183870/frederic-fumeron-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

150
papers

5,857
citations

43
h-index

72
g-index

165
ext. papers

6,440
ext. citations

8.3
avg, IF

4.76
L-index

#	Paper	IF	Citations
150	Plasma total adiponectin and changes in renal function in a cohort from the community: the prospective Data from an Epidemiological Study on the Insulin Resistance Syndrome study. <i>Nephrology Dialysis Transplantation</i> , 2021 , 36, 2058-2065	4.3	0
149	Effects of acyl-coenzyme A binding protein (ACBP)/diazepam-binding inhibitor (DBI) on body mass index. <i>Cell Death and Disease</i> , 2021 , 12, 599	9.8	5
148	Mendelian randomization analysis does not support causal associations of birth weight with hypertension risk and blood pressure in adulthood. <i>European Journal of Epidemiology</i> , 2020 , 35, 685-697	12.1	2
147	Relationship between renal capacity to reabsorb glucose and renal status in patients with diabetes. <i>Diabetes and Metabolism</i> , 2020 , 46, 488-495	5.4	1
146	Dairy consumption is associated with lower plasma dihydroceramides in women from the D.E.S.I.R. cohort. <i>Diabetes and Metabolism</i> , 2020 , 46, 144-149	5.4	1
145	Marian Apfelbaum 1931-2020. <i>Cahiers De Nutrition Et De Dietetique</i> , 2020 , 55, 56	0.2	
144	Plasma Apelin and Risk of Type 2 Diabetes in a Cohort From the Community. <i>Diabetes Care</i> , 2020 , 43, e15-e16	14.6	8
143	Association of Birth Weight With Type 2 Diabetes and Glycemic Traits: A Mendelian Randomization Study. <i>JAMA Network Open</i> , 2019 , 2, e1910915	10.4	14
142	New roles for prokineticin 2 in feeding behavior, insulin resistance and type 2 diabetes: Studies in mice and humans. <i>Molecular Metabolism</i> , 2019 , 29, 182-196	8.8	11
141	Plasma Copeptin and Risk of Lower-Extremity Amputation in Type 1 and Type 2 Diabetes. <i>Diabetes Care</i> , 2019 , 42, 2290-2297	14.6	10
140	Non-severe hypoglycaemia is associated with weight gain in patients with type 1 diabetes: Results from the Diabetes Control and Complication Trial. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 1289-1292	6.7	6
139	Plasma copeptin, kidney disease, and risk for cardiovascular morbidity and mortality in two cohorts of type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2018 , 17, 110	8.7	19
138	Plasma proprotein-convertase-subtilisin/kexin type 9 (PCSK9) and cardiovascular events in type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 943-953	6.7	9
137	Plasma copeptin and chronic kidney disease risk in 3 European cohorts from the general population. <i>JCI Insight</i> , 2018 , 3,	9.9	20
136	Dairy Consumption and Body Mass Index Among Adults: Mendelian Randomization Analysis of 184802 Individuals from 25 Studies. <i>Clinical Chemistry</i> , 2018 , 64, 183-191	5.5	24
135	Prognostic value of plasma MR-proADM vs NT-proBNP for heart failure in people with type 2 diabetes: the SURDIAGENE prospective study. <i>Diabetologia</i> , 2018 , 61, 2643-2653	10.3	6
134	T-cadherin gene variants are associated with type 2 diabetes and the Fatty Liver Index in the French population. <i>Diabetes and Metabolism</i> , 2017 , 43, 33-39	5.4	13

133	Association of ACACB gene polymorphism (rs2268388, G > A) with type 2 diabetes and end stage renal disease in Pakistani Punjabi population. <i>Meta Gene</i> , 2017 , 12, 109-112	0.7	2
132	Plasma Dihydroceramides Are Diabetes Susceptibility Biomarker Candidates in Mice and Humans. <i>Cell Reports</i> , 2017 , 18, 2269-2279	10.6	108
131	T-cadherin gene variants are associated with nephropathy in subjects with type 1 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2017 , 32, 1987-1993	4.3	2
130	Identification of novel risk loci for restless legs syndrome in genome-wide association studies in individuals of European ancestry: a meta-analysis. <i>Lancet Neurology</i> , 2017 , 16, 898-907	24.1	121
129	Urinary Sodium Concentration Is an Independent Predictor of All-Cause and Cardiovascular Mortality in a Type 2 Diabetes Cohort Population. <i>Journal of Diabetes Research</i> , 2017 , 2017, 5327352	3.9	8
128	La T-cadhřine, troisiřme rřcepteur de lřdiponectine : structure et rřle en santř humaine. <i>Obesite</i> , 2017 , 12, 267-276	0.1	
127	Interindividual variability in the cholesterol-lowering effect of supplementation with plant sterols or stanols. <i>Nutrition Reviews</i> , 2017 , 75, 134-145	6.4	14
126	Longitudinal association of antidepressant medication use with metabolic syndrome: Results of a 9-year follow-up of the D.E.S.I.R. cohort study. <i>Psychoneuroendocrinology</i> , 2016 , 74, 34-45	5	8
125	Death, end-stage renal disease and renal function decline in patients with diabetic nephropathy in French cohorts of type 1 and type 2 diabetes. <i>Diabetologia</i> , 2016 , 59, 208-216	10.3	31
124	New loci for body fat percentage reveal link between adiposity and cardiometabolic disease risk. <i>Nature Communications</i> , 2016 , 7, 10495	17.4	180
123	Glutathione peroxidase-1 gene (GPX1) variants, oxidative stress and risk of kidney complications in people with type 1 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 12-9	12.7	25
122	Les aliments fonctionnels contenant des phytostřfols ou des phytostřnols : quels břnřces, quels risques?. <i>Pratiques En Nutrition</i> , 2016 , 12, 32-35	0	1
121	Dynamic Changes in Renal Function Are Associated With Major Cardiovascular Events in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2016 , 39, 1259-66	14.6	26
120	Interaction between GPR120 p.R270H loss-of-function variant and dietary fat intake on incident type 2 diabetes risk in the D.E.S.I.R. study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 931-6	4.5	8
119	Plasma Copeptin, AVP Gene Variants, and Incidence of Type 2 Diabetes in a Cohort From the Community. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 2432-9	5.6	52
118	Plasma Copeptin, Kidney Outcomes, Ischemic Heart Disease, and All-Cause Mortality in People With Long-standing Type 1 Diabetes. <i>Diabetes Care</i> , 2016 , 39, 2288-2295	14.6	41
117	Plasma extracellular superoxide dismutase concentration, allelic variations in the SOD3 gene and risk of myocardial infarction and all-cause mortality in people with type 1 and type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 845	8.7	34
116	Allelic variations in the CYBA gene of NADPH oxidase and risk of kidney complications in patients with type 1 diabetes. <i>Free Radical Biology and Medicine</i> , 2015 , 86, 16-24	7.8	12

115	The loss-of-function PCSK9 p.R46L genetic variant does not alter glucose homeostasis. <i>Diabetologia</i> , 2015 , 58, 2051-5	10.3	36
114	ABCG8 polymorphisms and renal disease in type 2 diabetic patients. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 713-9	12.7	9
113	Phytostfols : un point sur les recommandations de l'ANSES. <i>Cahiers De Nutrition Et De Dietetique</i> , 2015 , 50, 209-214	0.2	2
112	Plasma Adrenomedullin and Allelic Variation in the ADM Gene and Kidney Disease in People With Type 2 Diabetes. <i>Diabetes</i> , 2015 , 64, 3262-72	0.9	7
111	Antidepressant medication use and trajectories of fasting plasma glucose, glycated haemoglobin, β -cell function and insulin sensitivity: a 9-year longitudinal study of the D.E.S.I.R. cohort. <i>International Journal of Epidemiology</i> , 2015 , 44, 1927-40	7.8	13
110	Potential risks associated with increased plasma plant-sterol levels. <i>Diabetes and Metabolism</i> , 2015 , 41, 76-81	5.4	16
109	Phytostfols : un point sur les recommandations de l'ANSES. <i>OCL - Oilseeds and Fats, Crops and Lipids</i> , 2015 , 22, D205	1.5	1
108	Contribution of the low-frequency, loss-of-function p.R270H mutation in FFAR4 (GPR120) to increased fasting plasma glucose levels. <i>Journal of Medical Genetics</i> , 2015 , 52, 595-8	5.8	22
107	Plasma Copeptin and Decline in Renal Function in a Cohort from the Community: The Prospective D.E.S.I.R. Study. <i>American Journal of Nephrology</i> , 2015 , 42, 107-14	4.6	28
106	Fine-scale human genetic structure in Western France. <i>European Journal of Human Genetics</i> , 2015 , 23, 831-6	5.3	22
105	Phytostfols, phytostanols, et risque cardiovasculaire. <i>Medecine Des Maladies Metaboliques</i> , 2014 , 8, 483-488	0.1	1
104	Tissue kallikrein deficiency, insulin resistance, and diabetes in mouse and man. <i>Journal of Endocrinology</i> , 2014 , 221, 297-308	4.7	4
103	Comparison between copeptin and vasopressin in a population from the community and in people with chronic kidney disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 4656-63	5.6	87
102	Adiponectin gene variants, adiponectin isoforms and cardiometabolic risk in type 2 diabetic patients. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 192-8	3.9	26
101	Manganese superoxide dismutase (SOD2) polymorphisms, plasma advanced oxidation protein products (AOPP) concentration and risk of kidney complications in subjects with type 1 diabetes. <i>PLoS ONE</i> , 2014 , 9, e96916	3.7	23
100	Genetics of the human obesities. <i>Obesity</i> , 2013 , 8, 22-33	0.1	
99	Catalase activity, allelic variations in the catalase gene and risk of kidney complications in patients with type 1 diabetes. <i>Diabetologia</i> , 2013 , 56, 2733-42	10.3	9
98	Allelic variations of the vitamin D receptor (VDR) gene are associated with increased risk of coronary artery disease in type 2 diabetics: the DIABHYCAR prospective study. <i>Diabetes and Metabolism</i> , 2013 , 39, 263-70	5.4	33

97	Dairy product consumption, calcium intakes, and metabolic syndrome-related factors over 5 years in the STANISLAS study. <i>Nutrition</i> , 2013 , 29, 519-24	4.8	47
96	Genetics of the Human Obesities 2013 , 351-372		
95	The lactase persistence genotype is associated with body mass index and dairy consumption in the D.E.S.I.R. study. <i>Metabolism: Clinical and Experimental</i> , 2013 , 62, 1323-9	12.7	29
94	Gh̃tique des ob̃sit̃s humaines 2013 , 359-380		
93	GUESS-ing polygenic associations with multiple phenotypes using a GPU-based evolutionary stochastic search algorithm. <i>PLoS Genetics</i> , 2013 , 9, e1003657	6	45
92	Plasma copeptin and renal outcomes in patients with type 2 diabetes and albuminuria. <i>Diabetes Care</i> , 2013 , 36, 3639-45	14.6	59
91	Dietary fat intake and polymorphisms at the PPARG locus modulate BMI and type 2 diabetes risk in the D.E.S.I.R. prospective study. <i>International Journal of Obesity</i> , 2012 , 36, 218-24	5.5	47
90	Allelic variations in superoxide dismutase-1 (SOD1) gene and renal and cardiovascular morbidity and mortality in type 2 diabetic subjects. <i>Molecular Genetics and Metabolism</i> , 2012 , 106, 359-65	3.7	30
89	Comment on: Park et al. Association of serum ferritin and the development of metabolic syndrome in middle-aged Korean men: a 5-year follow-up study. <i>Diabetes Care</i> 2012;35:2521-2526. <i>Diabetes Care</i> , 2012 , 35, e92	14.6	1
88	Intensifying glycaemic control with insulin reduces adiponectin and its HMW isoform moderately in type 2, but not in type 1, diabetes. <i>Diabetes and Metabolism</i> , 2011 , 37, 259-61	5.4	6
87	Allelic variations in superoxide dismutase-1 (SOD1) gene are associated with increased risk of diabetic nephropathy in type 1 diabetic subjects. <i>Molecular Genetics and Metabolism</i> , 2011 , 104, 654-60	3.7	31
86	Dairy consumption and the incidence of hyperglycemia and the metabolic syndrome: results from a french prospective study, Data from the Epidemiological Study on the Insulin Resistance Syndrome (DESIR). <i>Diabetes Care</i> , 2011 , 34, 813-7	14.6	125
85	Association of vitronectin and plasminogen activator inhibitor-1 levels with the risk of metabolic syndrome and type 2 diabetes mellitus. Results from the D.E.S.I.R. prospective cohort. <i>Thrombosis and Haemostasis</i> , 2011 , 106, 416-22	7	22
84	Decreased insulin secretion and increased risk of type 2 diabetes associated with allelic variations of the WFS1 gene: the Data from Epidemiological Study on the Insulin Resistance Syndrome (DESIR) prospective study. <i>Diabetologia</i> , 2011 , 54, 554-62	10.3	25
83	Dairy products and the metabolic syndrome in a prospective study, DESIR. <i>Journal of the American College of Nutrition</i> , 2011 , 30, 454S-63S	3.5	38
82	Hyperadiponectinemia is independent of kidney function, diabetes duration, and control in type 1 diabetic patients without microangiopathy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E485-7	5.6	5
81	Are the same clinical risk factors relevant for incident diabetes defined by treatment, fasting plasma glucose, and HbA1c?. <i>Diabetes Care</i> , 2011 , 34, 957-9	14.6	7
80	Sex-dependent associations of leptin with metabolic syndrome-related variables: the Stanislas study. <i>Obesity</i> , 2010 , 18, 196-201	8	20

79	Adiponectin multimers and ADIPOQ T45G in coronary artery disease in Caribbean type 2 diabetic subjects of African descent. <i>Obesity</i> , 2010 , 18, 1466-8	8	18
78	High baseline insulin levels associated with 6-year incident observed sleep apnea. <i>Diabetes Care</i> , 2010 , 33, 1044-9	14.6	31
77	The functional c.-2G>C variant of the mineralocorticoid receptor modulates blood pressure, renin, and aldosterone levels. <i>Hypertension</i> , 2010 , 56, 995-1002	8.5	40
76	Association of ADIPOQ genetic variants and plasma adiponectin isoforms with the risk of incident renal events in type 2 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2010 , 25, 2231-7	4.3	29
75	Associations of the -344 T>C and the 3097 G>A polymorphisms of CYP11B2 gene with hypertension, type 2 diabetes, and metabolic syndrome in a French population. <i>American Journal of Hypertension</i> , 2010 , 23, 660-7	2.3	16
74	Genetic variability at the six transmembrane protein of prostate 2 locus and the metabolic syndrome: the data from an epidemiological study on the Insulin Resistance Syndrome (DESIR) study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 2942-7	5.6	14
73	Polymorphisms of the lamina maturation pathway and their association with the metabolic syndrome: the DESIR prospective study. <i>Journal of Molecular Medicine</i> , 2010 , 88, 193-201	5.5	5
72	Nine-year incident diabetes is predicted by fatty liver indices: the French D.E.S.I.R. study. <i>BMC Gastroenterology</i> , 2010 , 10, 56	3	97
71	Association of common variants in NPPA and NPPB with blood pressure does not translate into kidney damage in a general population study. <i>Journal of Hypertension</i> , 2010 , 28, 1230-3	1.9	3
70	Sex hormone-binding globulin predicts the incidence of hyperglycemia in women: interactions with adiponectin levels. <i>European Journal of Endocrinology</i> , 2009 , 161, 81-5	6.5	35
69	Relationships between common polymorphisms of adenosine triphosphate-binding cassette transporter A1 and high-density lipoprotein cholesterol and coronary heart disease in a population with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2009 , 58, 74-9	12.7	35
68	Prognostic value of the insertion/deletion polymorphism of the ACE gene in type 2 diabetic subjects: results from the Non-insulin-dependent Diabetes, Hypertension, Microalbuminuria or Proteinuria, Cardiovascular Events, and Ramipril (DIABHYCAR), Diabete de type 2, Nephropathie et Complications (DIABNEDUCOGENE) and Genetic Diabetes, Hypertension and Complications (GUDDIAGENE) studies. <i>Diabetes Care</i> , 2009 , 32, 103-8	14.6	58
67	The common -866G>A variant in the promoter of UCP2 is associated with decreased risk of coronary artery disease in type 2 diabetic men. <i>Diabetes</i> , 2008 , 57, 1063-8	0.9	42
66	Predicting diabetes: clinical, biological, and genetic approaches: data from the Epidemiological Study on the Insulin Resistance Syndrome (DESIR). <i>Diabetes Care</i> , 2008 , 31, 2056-61	14.6	183
65	Impact of common type 2 diabetes risk polymorphisms in the DESIR prospective study. <i>Diabetes</i> , 2008 , 57, 244-54	0.9	137
64	Analysis of novel risk loci for type 2 diabetes in a general French population: the D.E.S.I.R. study. <i>Journal of Molecular Medicine</i> , 2008 , 86, 341-8	5.5	60
63	Adiponectin gene and cardiovascular risk in type 2 diabetic patients: a review of evidences. <i>Arquivos Brasileiros De Endocrinologia E Metabologia</i> , 2007 , 51, 153-9		15
62	Evidence for distinct effects of GH and IGF-I in the metabolic syndrome. <i>Diabetic Medicine</i> , 2007 , 24, 1013-8		16

61	Ferritin and transferrin are associated with metabolic syndrome abnormalities and their change over time in a general population: Data from an Epidemiological Study on the Insulin Resistance Syndrome (DESIR). <i>Diabetes Care</i> , 2007 , 30, 1795-801	14.6	100
60	The CETP TaqIB polymorphism is associated with the risk of sudden death in type 2 diabetic patients. <i>Diabetes Care</i> , 2007 , 30, 2863-7	14.6	9
59	Patterns of Alcohol Consumption and Cardiovascular Risk in Northern Ireland and France. <i>Annals of Epidemiology</i> , 2007 , 17, S75-S80	6.4	5
58	Genetics of macrovascular complications in diabetes. <i>Current Diabetes Reports</i> , 2006 , 6, 162-8	5.6	9
57	Weight change and changes in the metabolic syndrome as the French population moves towards overweight: the D.E.S.I.R. cohort. <i>International Journal of Epidemiology</i> , 2006 , 35, 190-6	7.8	55
56	The PPARG Pro12Ala polymorphism is associated with a decreased risk of developing hyperglycemia over 6 years and combines with the effect of the APM1 G-11391A single nucleotide polymorphism: the Data From an Epidemiological Study on the Insulin Resistance Syndrome (DESIR)	0.9	53
55	Ferritin and transferrin are both predictive of the onset of hyperglycemia in men and women over 3 years: the data from an epidemiological study on the Insulin Resistance Syndrome (DESIR) study. <i>Diabetes Care</i> , 2006 , 29, 2090-4	14.6	102
54	Bardet-Biedl syndrome gene variants are associated with both childhood and adult common obesity in French Caucasians. <i>Diabetes</i> , 2006 , 55, 2876-82	0.9	68
53	ACDC/adiponectin polymorphisms are associated with severe childhood and adult obesity. <i>Diabetes</i> , 2006 , 55, 545-50	0.9	139
52	ABCA1 single nucleotide polymorphisms on high-density lipoprotein-cholesterol and overweight: the D.E.S.I.R. study. <i>Obesity</i> , 2006 , 14, 1874-9	8	41
51	Increased plasma adiponectin concentrations are associated with microangiopathy in type 1 diabetic subjects. <i>Diabetologia</i> , 2005 , 48, 1088-92	10.3	86
50	Polymorphism of the 5-HT2A receptor gene and food intakes in children and adolescents: the Stanislas Family Study. <i>American Journal of Clinical Nutrition</i> , 2005 , 82, 467-70	7	9
49	Polymorphism of the 5-HT2A receptor gene and food intakes in children and adolescents: the Stanislas Family Study. <i>American Journal of Clinical Nutrition</i> , 2005 , 82, 467-470	7	28
48	Association between endothelin receptor B nonsynonymous variants and melanoma risk. <i>Journal of the National Cancer Institute</i> , 2005 , 97, 1297-301	9.7	21
47	Adiponectin gene polymorphisms and adiponectin levels are independently associated with the development of hyperglycemia during a 3-year period: the epidemiologic data on the insulin resistance syndrome prospective study. <i>Diabetes</i> , 2004 , 53, 1150-7	0.9	172
46	Apolipoprotein B signal peptide polymorphism and plasma LDL-cholesterol response to low-calorie diet. <i>International Journal of Obesity</i> , 2004 , 28, 902-5	5.5	16
45	Obésité: dūn syndrome monogħique exceptionnel aux interactions entre għes multiples et environnement nutritionnel. <i>Oleagineux Corps Gras Lipides</i> , 2003 , 10, 109-114		
44	The B3T -> C Polymorphism in Intron 7 of the TFPI Gene Influences the Risk of Venous Thromboembolism, Independently of the Factor V Leiden and Prothrombin Mutations. <i>Thrombosis and Haemostasis</i> , 2002 , 88, 195-199	7	36

43	Serotonin transporter gene polymorphism and myocardial infarction: Etude Cas-Tmoins de l'Infarctus du Myocarde (ECTIM). <i>Circulation</i> , 2002 , 105, 2943-5	16.7	71
42	The -33T-->C polymorphism in intron 7 of the TFPI gene influences the risk of venous thromboembolism, independently of the factor V Leiden and prothrombin mutations. <i>Thrombosis and Haemostasis</i> , 2002 , 88, 195-9	7	8
41	Polymorphism in the fractalkine receptor CX3CR1 as a genetic risk factor for coronary artery disease. <i>Blood</i> , 2001 , 97, 1925-8	2.2	289
40	LEPR gene polymorphisms: associations with overweight, fat mass and response to diet in women. <i>European Journal of Clinical Investigation</i> , 2001 , 31, 398-404	4.6	65
39	Association of a functional 5-HT transporter gene polymorphism with anorexia nervosa and food intake. <i>Molecular Psychiatry</i> , 2001 , 6, 9-10	15.1	72
38	A New T-287C Polymorphism in the 5'Regulatory Region of the Tissue Factor Pathway Inhibitor Gene. <i>Thrombosis and Haemostasis</i> , 2000 , 84, 244-249	7	47
37	Extensive association analysis between the CETP gene and coronary heart disease phenotypes reveals several putative functional polymorphisms and gene-environment interaction. <i>Genetic Epidemiology</i> , 2000 , 19, 64-80	2.6	192
36	Association of the G-2548A polymorphism in the 5' region of the LEP gene with overweight. <i>Annals of Human Genetics</i> , 2000 , 64, 391-394	2.2	140
35	5-HT2A receptor gene polymorphism is associated with food and alcohol intake in obese people. <i>International Journal of Obesity</i> , 2000 , 24, 920-4	5.5	64
34	Association of the G-2548A polymorphism in the 5' region of the LEP gene with overweight. <i>Annals of Human Genetics</i> , 2000 , 64, 391-4	2.2	147
33	Beneficial effects of a moderate consumption of red wine on cellular cholesterol efflux in young men. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2000 , 10, 63-9	4.5	29
32	A new T-287C polymorphism in the 5' regulatory region of the tissue factor pathway inhibitor gene. Association study of the T-287C and C-399T polymorphisms with coronary artery disease and plasma TFPI levels. <i>Thrombosis and Haemostasis</i> , 2000 , 84, 244-9	7	6
31	Polymorphisms of the tissue factor pathway inhibitor (TFPI) gene in patients with acute coronary syndromes and in healthy subjects : impact of the V264M substitution on plasma levels of TFPI. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999 , 19, 862-9	9.4	49
30	Lack of association between genetic variations of apo A-I-C-III-A-IV gene cluster and myocardial infarction in a sample of European male: ECTIM study. <i>Atherosclerosis</i> , 1999 , 145, 187-95	3.1	35
29	Influence of the saci apo C-III gene polymorphism on the insulin response to an oral glucose tolerance test in obese subjects. <i>Atherosclerosis</i> , 1999 , 144, 109-110	3.1	68
28	Dietary fat, energy density and BMI: a case of a missing flower?. <i>International Journal of Obesity</i> , 1998 , 22, 1032-4	5.5	2
27	Novel polymorphisms in the 5' region of the LEP gene: association with leptin levels and response to low-calorie diet in human obesity. <i>Diabetes</i> , 1998 , 47, 487-9	0.9	135
26	Lipoprotein lipase gene variants and coronary risk. <i>Circulation</i> , 1998 , 97, 2588	16.7	1

25	Hind III polymorphism of the lipoprotein lipase gene and plasma lipid response to low calorie diet. <i>International Journal of Obesity</i> , 1997 , 21, 280-3	5.5	16
24	Lipoprotein lipase gene polymorphisms: associations with myocardial infarction and lipoprotein levels, the ECTIM study. Etude Cas Tmoin sur l'Infarctus du Myocarde.. <i>Journal of Lipid Research</i> , 1996 , 36, 2141-2146	6.3	100
23	Lipoprotein lipase gene polymorphisms: associations with hypertriglyceridemia and body mass index in obese people 1995 , 19, 270-4		13
22	Alcohol intake modulates the effect of a polymorphism of the cholesteryl ester transfer protein gene on plasma high density lipoprotein and the risk of myocardial infarction. <i>Journal of Clinical Investigation</i> , 1995 , 96, 1664-71	15.9	229
21	Lipoprotein lipase gene polymorphisms: associations with myocardial infarction and lipoprotein levels, the ECTIM study. Etude Cas Tmoin sur l'Infarctus du Myocarde. <i>Journal of Lipid Research</i> , 1995 , 36, 2141-6	6.3	98
20	Xmn1 restriction polymorphism of apolipoprotein AI gene and lipoprotein levels in obesity 1994 , 18, 460-4		
19	Presence of the French Canadian deletion in a French patient with familial hypercholesterolemia. <i>New England Journal of Medicine</i> , 1992 , 326, 69	59.2	29
18	Presence of the French Canadian Deletion in a French Patient with Familial Hypercholesterolemia. <i>New England Journal of Medicine</i> , 1992 , 326, 69-69	59.2	48
17	Effect of a moderate alcohol intake on the lipoproteins of normotriglyceridemic obese subjects compared with normoponderal controls. <i>Metabolism: Clinical and Experimental</i> , 1992 , 41, 856-61	12.7	32
16	Les m̄astases h̄patiques ont-elles un r̄le n̄faste sur la positivation du bilan azot̄ sous nutrition ent̄rale chez les patients canc̄reux ?. <i>Nutrition Clinique Et Metabolisme</i> , 1991 , 5, 5-10	0.8	1
15	Lack of plasmic beta-endorphin response to a gastronomic meal in healthy humans. <i>Physiology and Behavior</i> , 1991 , 49, 1217-21	3.5	6
14	Lowering of HDL2-cholesterol and lipoprotein A-I particle levels by increasing the ratio of polyunsaturated to saturated fatty acids. <i>American Journal of Clinical Nutrition</i> , 1991 , 53, 655-9	7	47
13	n-3 polyunsaturated fatty acids raise low-density lipoproteins, high-density lipoprotein 2, and plasminogen-activator inhibitor in healthy young men. <i>American Journal of Clinical Nutrition</i> , 1991 , 54, 118-22	7	51
12	Iron intake in obese menstruating women. <i>Diabetes Research and Clinical Practice</i> , 1990 , 10 Suppl 1, S287-90		
11	Lack of association between dietary alcohol and HDL-cholesterol concentrations in obese women. <i>Atherosclerosis</i> , 1990 , 81, 119-25	3.1	7
10	Circadian rhythm of energy intake and corpulence status in adults 1990 , 14, 387-93		1
9	Association of apolipoprotein epsilon 4 allele with hypertriglyceridemia in obesity. <i>Clinical Genetics</i> , 1988 , 34, 258-64	4	26
8	Low high density lipoprotein-2 concentrations in obese male subjects. <i>Atherosclerosis</i> , 1988 , 73, 57-61	3.1	20

7	Comparative effects of several simple carbohydrates on erythrocyte insulin receptors in obese subjects. <i>Pharmacology Biochemistry and Behavior</i> , 1986 , 25, 681-8	3.9	15
6	Effect of slight plasma glucose decrease on glycosylated hemoglobin in healthy subjects during caloric restriction. <i>New England Journal of Medicine</i> , 1985 , 313, 958-9	59.2	4
5	Endogenous opiates and energy balance. <i>Science</i> , 1982 , 215, 1536-8	33.3	91
4	HLA and longevity. <i>Tissue Antigens</i> , 1982 , 19, 168-73		81
3	Studies on an isolated West Indies population: I. Analysis of HLA genotypes. <i>Tissue Antigens</i> , 1981 , 17, 338-42		2
2	Association between HLA-B18 and the familial-obesity syndrome. <i>New England Journal of Medicine</i> , 1981 , 305, 645	59.2	5
1	HLA genotypes in familial Hodgkin's disease. Excess of HLA identical affected subs. <i>European Journal of Cancer</i> , 1980 , 16, 809-15		16