Erwin E H Van Wegen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6186078/publications.pdf

Version: 2024-02-01

141 papers 10,432 citations

51 h-index 97 g-index

153 all docs

153 docs citations

153 times ranked 9658 citing authors

#	Article	IF	CITATIONS
1	What Is the Evidence for Physical Therapy Poststroke? A Systematic Review and Meta-Analysis. PLoS ONE, 2014, 9, e87987.	2.5	854
2	Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. Journal of Neurology, Neurosurgery and Psychiatry, 2007, 78, 134-140.	1.9	677
3	Early Prediction of Outcome of Activities of Daily Living After Stroke. Stroke, 2011, 42, 1482-1488.	2.0	421
4	Effects of external rhythmical cueing on gait in patients with Parkinson's disease: a systematic review. Clinical Rehabilitation, 2005, 19, 695-713.	2.2	412
5	Effects of Robot-Assisted Therapy for the Upper Limb After Stroke. Neurorehabilitation and Neural Repair, 2017, 31, 107-121.	2.9	398
6	Constraint-induced movement therapy after stroke. Lancet Neurology, The, 2015, 14, 224-234.	10.2	365
7	Standardized measurement of sensorimotor recovery in stroke trials: Consensus-based core recommendations from the Stroke Recovery and Rehabilitation Roundtable. International Journal of Stroke, 2017, 12, 451-461.	5.9	352
8	Presence of Finger Extension and Shoulder Abduction Within 72 Hours After Stroke Predicts Functional Recovery. Stroke, 2010, 41, 745-750.	2.0	334
9	Attending to the task: Interference effects of functional tasks on walking in Parkinsona ** "s disease and the roles of cognition, depression, fatigue, and balance 11No party having a direct interest in the results of the research supporting this article has or will confer a benefit on the author(s) or on any organization with which the author(s) is/are associated Archives of Physical Medicine and	0.9	265
10	Generalizability of the Proportional Recovery Model for the Upper Extremity After an Ischemic Stroke. Neurorehabilitation and Neural Repair, 2015, 29, 614-622.	2.9	250
11	The Effect of External Rhythmic Cues (Auditory and Visual) on Walking During a Functional Task in Homes of People With Parkinson's Disease. Archives of Physical Medicine and Rehabilitation, 2005, 86, 999-1006.	0.9	219
12	On the Functional Aspects of Variability in Postural Control. Exercise and Sport Sciences Reviews, 2002, 30, 177-183.	3.0	201
13	Evidence for motor learning in Parkinson's disease: Acquisition, automaticity and retention of cued gait performance after training with external rhythmical cues. Brain Research, 2010, 1319, 103-111.	2.2	172
14	Exercise therapy for fatigue in multiple sclerosis. The Cochrane Library, 2015, 2015, CD009956.	2.8	163
15	The use of rhythmic auditory cues to influence gait in patients with Parkinson's disease, the differential effect for freezers and non-freezers, an explorative study. Disability and Rehabilitation, 2006, 28, 721-728.	1.8	159
16	Self-report fatigue questionnaires in multiple sclerosis, Parkinson's disease and stroke: a systematic review of measurement properties. Quality of Life Research, 2012, 21, 925-944.	3.1	155
17	Effects of Augmented Exercise Therapy on Outcome of Gait and Gait-Related Activities in the First 6 Months After Stroke. Stroke, 2011, 42, 3311-3315.	2.0	154
18	Postural orientation: Age-related changes in variability and time-to-boundary. Human Movement Science, 2002, 21, 61-84.	1.4	143

#	Article	IF	CITATIONS
19	Effects of Unilateral Upper Limb Training in Two Distinct Prognostic Groups Early After Stroke. Neurorehabilitation and Neural Repair, 2016, 30, 804-816.	2.9	140
20	Standardized Measurement of Sensorimotor Recovery in Stroke Trials: Consensus-Based Core Recommendations from the Stroke Recovery and Rehabilitation Roundtable. Neurorehabilitation and Neural Repair, 2017, 31, 784-792.	2.9	135
21	Measuring fatigue in patients with multiple sclerosis: reproducibility, responsiveness and concurrent validity of three Dutch self-report questionnaires. Disability and Rehabilitation, 2010, 32, 1870-1876.	1.8	131
22	Is Accurate Prediction of Gait in Nonambulatory Stroke Patients Possible Within 72 Hours Poststroke?. Neurorehabilitation and Neural Repair, 2011, 25, 268-274.	2.9	126
23	Measuring gait and gait-related activities in Parkinson's patients own home environment: a reliability, responsiveness and feasibility study. Parkinsonism and Related Disorders, 2005, 11, 19-24.	2.2	123
24	The attentional cost of external rhythmical cues and their impact on gait in Parkinson's disease: effect of cue modality and task complexity. Journal of Neural Transmission, 2007, 114, 1243-1248.	2.8	123
25	On Variability and Stability in Human Movement. Journal of Applied Biomechanics, 2000, 16, 394-406.	0.8	110
26	The Short-Term Effects of Different Cueing Modalities on Turn Speed in People with Parkinson's Disease. Neurorehabilitation and Neural Repair, 2009, 23, 831-836.	2.9	99
27	Exercise-induced increase in brain-derived neurotrophic factor in human Parkinson's disease: a systematic review and meta-analysis. Translational Neurodegeneration, 2018, 7, 7.	8.0	97
28	The effect of rhythmic somatosensory cueing on gait in patients with Parkinson's disease. Journal of the Neurological Sciences, 2006, 248, 210-214.	0.6	94
29	Stability Boundaries and Lateral Postural Control in Parkinson's Disease. Motor Control, 2001, 5, 254-269.	0.6	89
30	Turning in Parkinson's disease patients and controls: The effect of auditory cues. Movement Disorders, 2007, 22, 1871-1878.	3.9	87
31	A comparison of two validated tests for upper limb function after stroke: The Wolf Motor Function Test and the Action Research Arm Test. Journal of Rehabilitation Medicine, 2010, 42, 694-696.	1.1	87
32	Impact of Time on Quality of Motor Control of the Paretic Upper Limb After Stroke. Archives of Physical Medicine and Rehabilitation, 2014, 95, 338-344.	0.9	86
33	Walking speed during single and dual tasks in Parkinson's disease: Which characteristics are important?. Movement Disorders, 2008, 23, 2312-2318.	3.9	84
34	Age-related changes in upper body adaptation to walking speed in human locomotion. Gait and Posture, 2005, 22, 233-239.	1.4	82
35	Constraint-Induced Movement Therapy for the Upper Paretic Limb in Acute or Sub-Acute Stroke: A Systematic Review. International Journal of Stroke, 2011, 6, 425-433.	5.9	82
36	Lifestyle Interventions to Prevent Cardiovascular Events After Stroke and Transient Ischemic Attack. Stroke, 2017, 48, 174-179.	2.0	79

#	Article	IF	CITATIONS
37	Predictive value of the NIHSS for ADL outcome after ischemic hemispheric stroke: Does timing of early assessment matter?. Journal of the Neurological Sciences, 2010, 294, 57-61.	0.6	78
38	Understanding Adaptive Motor Control of the Paretic Upper Limb Early Poststroke. Neurorehabilitation and Neural Repair, 2013, 27, 854-863.	2.9	76
39	Early Supported Discharge by Caregiver-Mediated Exercises and e-Health Support After Stroke. Stroke, 2016, 47, 1885-1892.	2.0	74
40	Diagnostic Accuracy of the Barthel Index for Measuring Activities of Daily Living Outcome After Ischemic Hemispheric Stroke. Stroke, 2011, 42, 342-346.	2.0	71
41	Is impact of fatigue an independent factor associated with physical activity in patients with idiopathic Parkinson's disease?. Movement Disorders, 2009, 24, 1512-1518.	3.9	67
42	Feasibility of external rhythmic cueing with the Google Glass for improving gait in people with Parkinson's disease. Journal of Neurology, 2016, 263, 1156-1165.	3.6	67
43	Postural control of the trunk during unstable sitting in Parkinson's disease. Parkinsonism and Related Disorders, 2006, 12, 492-498.	2.2	61
44	Everyday walking with Parkinson's disease: Understanding personal challenges and strategies. Disability and Rehabilitation, 2008, 30, 1213-1221.	1.8	61
45	Effects of augmented visual feedback during balance training in Parkinson's disease: A pilot randomized clinical trial. Parkinsonism and Related Disorders, 2014, 20, 1352-1358.	2.2	61
46	Does Cueing Training Improve Physical Activity in Patients With Parkinson's Disease?. Neurorehabilitation and Neural Repair, 2010, 24, 469-477.	2.9	59
47	Unraveling the interaction between pathological upper limb synergies and compensatory trunk movements during reach-to-grasp after stroke: a cross-sectional study. Experimental Brain Research, 2012, 221, 251-262.	1.5	59
48	Effectiveness of Botulinum Toxin Treatment for Upper Limb Spasticity Poststroke Over Different ICF Domains: A Systematic Review and Meta-Analysis. Archives of Physical Medicine and Rehabilitation, 2019, 100, 1703-1725.	0.9	59
49	Reliability and structural validity of the Multidimensional Fatigue Inventory (MFI) in patients with idiopathic Parkinson's disease. Parkinsonism and Related Disorders, 2012, 18, 532-536.	2.2	58
50	Usability of Videogame-Based Dexterity Training in the Early Rehabilitation Phase of Stroke Patients: A Pilot Study. Frontiers in Neurology, 2017, 8, 654.	2.4	58
51	The effects of visual rhythms and optic flow on stride patterns of patients with Parkinson's disease. Parkinsonism and Related Disorders, 2006, 12, 21-27.	2.2	57
52	Short-Term Effects of Cerebellar tDCS on Standing Balance Performance in Patients with Chronic Stroke and Healthy Age-Matched Elderly. Cerebellum, 2018, 17, 575-589.	2.5	56
53	Impact of early applied upper limb stimulation: The EXPLICIT-stroke programme design. BMC Neurology, 2008, 8, 49.	1.8	54
54	Caregiver-mediated exercises for improving outcomes after stroke. The Cochrane Library, 2016, 12, CD011058.	2.8	53

#	Article	IF	Citations
55	Caregiver-mediated exercises with e-health support for early supported discharge after stroke (CARE4STROKE): A randomized controlled trial. PLoS ONE, 2019, 14, e0214241.	2.5	53
56	Generalizability of the Maximum Proportional Recovery Rule to Visuospatial Neglect Early Poststroke. Neurorehabilitation and Neural Repair, 2017, 31, 334-342.	2.9	48
57	The effects of cognitive behavioral and mindfulness-based therapies on psychological distress in patients with multiple sclerosis, Parkinson's disease and Huntington's disease: Two meta-analyses. Journal of Psychosomatic Research, 2019, 122, 43-51.	2.6	45
58	Home based training for dexterity in Parkinson's disease: A randomized controlled trial. Parkinsonism and Related Disorders, 2017, 41, 92-98.	2.2	44
59	Gait and gait-related activities and fatigue in Parkinson's disease: What is the relationship?. Disability and Rehabilitation, 2006, 28, 1365-1371.	1.8	43
60	Brain activation is related to smoothness of upper limb movements after stroke. Experimental Brain Research, 2016, 234, 2077-2089.	1.5	43
61	Moving stroke rehabilitation forward: The need to change research. NeuroRehabilitation, 2018, 43, 19-30.	1.3	42
62	Accuracy of Physical Therapists' Early Predictions of Upper-Limb Function in Hospital Stroke Units: The EPOS Study. Physical Therapy, 2013, 93, 460-469.	2.4	41
63	Determination of head conductivity frequency response in vivo with optimized EIT-EEG. Neurolmage, 2016, 127, 484-495.	4.2	41
64	Effect of ketoprofen on muscle function and sEMG activity after eccentric exercise. Medicine and Science in Sports and Exercise, 2001, 33, 702-710.	0.4	39
65	How to design clinical rehabilitation trials for the upper paretic limb early post stroke?. Trials, 2016, 17, 468.	1.6	39
66	Is the proportional recovery rule applicable to the lower limb after a first-ever ischemic stroke? PLoS ONE, 2018, 13, e0189279.	2.5	39
67	When Does Return of Voluntary Finger Extension Occur Post-Stroke? A Prospective Cohort Study. PLoS ONE, 2016, 11, e0160528.	2.5	39
68	Pelvic Floor Muscle Exercise Therapy with Myofeedback for Women with Stress Urinary Incontinence: A Meta-analysis. Physiotherapy, 1996, 82, 107-113.	0.4	38
69	Interventions for fatigue in Parkinson's disease. The Cochrane Library, 2015, 2015, CD010925.	2.8	38
70	Motor Switching and Motor Adaptation Deficits Contribute to Freezing of Gait in Parkinson's Disease. Neurorehabilitation and Neural Repair, 2015, 29, 132-142.	2.9	38
71	How does upper extremity Fugl-Meyer motor score relate to resting-state EEG in chronic stroke? A power spectral density analysis. Clinical Neurophysiology, 2019, 130, 856-862.	1.5	38
72	The association between perceived fatigue and actual level of physical activity in multiple sclerosis. Multiple Sclerosis Journal, 2011, 17, 1231-1237.	3.0	37

#	Article	IF	CITATIONS
73	Slowing of M1 activity in Parkinson's disease during rest and movement – An MEG study. Clinical Neurophysiology, 2011, 122, 789-795.	1.5	36
74	Assessing Longitudinal Change in Coordination of the Paretic Upper Limb Using On-Site 3-Dimensional Kinematic Measurements. Physical Therapy, 2012, 92, 142-151.	2.4	36
75	Novel artefact removal algorithms for co-registered EEG/fMRI based on selective averaging and subtraction. Neurolmage, 2013, 64, 407-415.	4.2	36
76	Is Recovery of Somatosensory Impairment Conditional for Upper-Limb Motor Recovery Early After Stroke?. Neurorehabilitation and Neural Repair, 2020, 34, 403-416.	2.9	36
77	Exergaming-Based Dexterity Training in Persons With Parkinson Disease: A Pilot Feasibility Study. Journal of Neurologic Physical Therapy, 2019, 43, 168-174.	1.4	35
78	Do Patients With Multiple Sclerosis Show Different Daily Physical Activity Patterns From Healthy Individuals?. Neurorehabilitation and Neural Repair, 2014, 28, 516-523.	2.9	34
79	Is gait speed a valid measure to predict community ambulation in patients with Parkinsonââ,¬â"¢s disease?. Journal of Rehabilitation Medicine, 2013, 45, 370-375.	1.1	33
80	E-health Support in People with Parkinson's Disease with Smart Glasses: A Survey of User Requirements and Expectations in the Netherlands. Journal of Parkinson's Disease, 2015, 5, 369-378.	2.8	31
81	Caregiver-mediated exercises with e-health support for early supported discharge after stroke (CARE4STROKE): study protocol for a randomized controlled trial. BMC Neurology, 2015, 15, 193.	1.8	30
82	Body awareness training in the treatment of wearing-off related anxiety in patients with Parkinson's disease: Results from a pilot randomized controlled trial. Journal of Psychosomatic Research, 2017, 103, 1-8.	2.6	30
83	Identifying fallers with Parkinson's disease using homeâ€based tests: Who is at risk?. Movement Disorders, 2008, 23, 2411-2415.	3.9	27
84	Effects of Multidisciplinary Rehabilitation on Chronic Fatigue in Multiple Sclerosis: A Randomized Controlled Trial. PLoS ONE, 2014, 9, e107710.	2.5	27
85	Influence of focus of attention, reinvestment and fall history on elderly gait stability. Physiological Reports, 2017, 5, e13061.	1.7	25
86	Computerised patient-specific prediction of the recovery profile of upper limb capacity within stroke services: the next step. Journal of Neurology, Neurosurgery and Psychiatry, 2021, 92, 574-581.	1.9	25
87	Harnessing Cueing Training for Neuroplasticity in Parkinson Disease. Topics in Geriatric Rehabilitation, 2014, 30, 46-57.	0.4	24
88	Reliability and validity of a new dexterity questionnaire (DextQ-24) in Parkinson's disease. Parkinsonism and Related Disorders, 2016, 33, 78-83.	2.2	23
89	Impact of fatigue on health-related quality of life in patients with Parkinson's disease: a prospective study. Clinical Rehabilitation, 2014, 28, 300-311.	2.2	22
90	Is Resting-State EEG Longitudinally Associated With Recovery of Clinical Neurological Impairments Early Poststroke? A Prospective Cohort Study. Neurorehabilitation and Neural Repair, 2020, 34, 389-402.	2.9	22

#	Article	IF	Citations
91	The effects of augmented visual feedback during balance training in Parkinson's disease: study design of a randomized clinical trial. BMC Neurology, 2013, 13, 137.	1.8	21
92	Respiratory muscle training for multiple sclerosis. The Cochrane Library, 2017, 2017, CD009424.	2.8	20
93	Smoothness metrics for reaching performance after stroke. Part 1: which one to choose?. Journal of NeuroEngineering and Rehabilitation, 2021, 18, 154.	4.6	20
94	Interlimb Coupling Patterns in Human Locomotion: Are We Bipeds or Quadrupeds?. Annals of the New York Academy of Sciences, 1998, 860, 539-542.	3.8	19
95	Are early measured resting-state EEG parameters predictive for upper limb motor impairment six months poststroke?. Clinical Neurophysiology, 2021, 132, 56-62.	1.5	19
96	Quantifying Quality of Reaching Movements Longitudinally Post-Stroke: A Systematic Review. Neurorehabilitation and Neural Repair, 2022, 36, 183-207.	2.9	19
97	Exercise-induced increase in blood-based brain-derived neurotrophic factor (BDNF) in people with multiple sclerosis: A systematic review and meta-analysis of exercise intervention trials. PLoS ONE, 2022, 17, e0264557.	2.5	19
98	The effects of visual feedback during a rhythmic weight-shifting task in patients with Parkinson's disease. Gait and Posture, 2016, 48, 140-145.	1.4	18
99	Quantification of task-dependent cortical activation evoked by robotic continuous wrist joint manipulation in chronic hemiparetic stroke. Journal of NeuroEngineering and Rehabilitation, 2017, 14, 30.	4.6	18
100	Effects of attentional focus on walking stability in elderly. Gait and Posture, 2017, 55, 94-99.	1.4	17
101	Description of the <scp>CARE4STROKE</scp> programme: A caregiverâ€mediated exercises intervention with eâ€health support for stroke patients. Physiotherapy Research International, 2018, 23, e1719.	1.5	16
102	Measurement Properties of the NeuroFlexor Device for Quantifying Neural and Non-neural Components of Wrist Hyper-Resistance in Chronic Stroke. Frontiers in Neurology, 2019, 10, 730.	2.4	16
103	Prospectively Classifying Community Walkers After Stroke: Who Are They?. Archives of Physical Medicine and Rehabilitation, 2019, 100, 2113-2118.	0.9	16
104	Smoothness metric during reach-to-grasp after stroke: part 2. longitudinal association with motor impairment. Journal of NeuroEngineering and Rehabilitation, 2021, 18, 144.	4.6	16
105	How Reproducible Is Home-Based 24-Hour Ambulatory Monitoring of Motor Activity in Patients With Multiple Sclerosis?. Archives of Physical Medicine and Rehabilitation, 2010, 91, 1537-1541.	0.9	15
106	Characterizing the Protocol for Early Modified Constraintâ€induced Movement Therapy in the EXPLICITâ€Stroke Trial. Physiotherapy Research International, 2013, 18, 1-15.	1.5	15
107	Partnered Dancing to Improve Mobility for People With Parkinson's Disease. Frontiers in Neuroscience, 2015, 9, 444.	2.8	15
108	Getting into a "Flow―state: a systematic review of flow experience in neurological diseases. Journal of NeuroEngineering and Rehabilitation, 2021, 18, 65.	4.6	15

#	Article	IF	CITATIONS
109	Experiences of patients with stroke and their caregivers with caregiver-mediated exercises during the CARE4STROKE trial. Disability and Rehabilitation, 2020, 42, 698-704.	1.8	13
110	Sensor assisted self-management in Parkinson's disease: A feasibility study of ambulatory posture detection and feedback to treat stooped posture. Parkinsonism and Related Disorders, 2018, 46, S57-S61.	2.2	13
111	Tablet App Based Dexterity Training in Multiple Sclerosis (TAD-MS): Research Protocol of a Randomized Controlled Trial. Frontiers in Neurology, 2019, 10, 61.	2.4	10
112	The association between freezing of gait, fear of falling and anxiety in Parkinson's disease: a longitudinal analysis. Neurodegenerative Disease Management, 2020, 10, 159-168.	2.2	10
113	Invited Commentary on Comparison of Robotics, Functional Electrical Stimulation, and Motor Learning Methods for Treatment of Persistent Upper Extremity Dysfunction After Stroke: A Randomized ControlledÂTrial. Archives of Physical Medicine and Rehabilitation, 2015, 96, 991-993.	0.9	9
114	The Subjective Experience of Living with Parkinson's Disease: A Meta-Ethnography of Qualitative Literature. Journal of Parkinson's Disease, 2021, 11, 139-151.	2.8	9
115	Feasibility of a Home-Based Tablet App for Dexterity Training in Multiple Sclerosis: Usability Study. JMIR MHealth and UHealth, 2020, 8, e18204.	3.7	9
116	Constraint-induced movement therapy improves upper extremity motor function after stroke. Australian Journal of Physiotherapy, 2007, 53, 132.	0.9	7
117	BEWARE: Body awareness training in the treatment of wearing-off related anxiety in patients with Parkinson's disease: study protocol for a randomized controlled trial. Trials, 2015, 16, 283.	1.6	7
118	Position-Cortical Coherence as a Marker of Afferent Pathway Integrity Early Poststroke: A Prospective Cohort Study. Neurorehabilitation and Neural Repair, 2020, 34, 344-359.	2.9	7
119	Agreement and differences regarding family functioning between patients with acquired brain injury and their partners. Brain Injury, 2020, 34, 489-495.	1.2	7
120	Family-delivered rehabilitation services at home: is the glass empty?. Lancet, The, 2017, 390, 538-539.	13.7	6
121	Incongruent visual feedback during a postural task enhances cortical alpha and beta modulation in patients with Parkinson's disease. Clinical Neurophysiology, 2018, 129, 1357-1365.	1.5	5
122	Portable Gait Lab: Zero Moment Point for Minimal Sensing of Gait., 2019, 2019, 2077-2081.		5
123	Self-monitoring of Physical Activity After Hospital Discharge in Patients Who Have Undergone Gastrointestinal or Lung Cancer Surgery: Mixed Methods Feasibility Study. JMIR Cancer, 2022, 8, e35694.	2.4	5
124	High-Intensity Interval Cycle Ergometer Training in Parkinson's Disease: Protocol for Identifying Individual Response Patterns Using a Single-Subject Research Design. Frontiers in Neurology, 2020, 11, 569880.	2.4	4
125	Time Course of Wrist Hyper-Resistance in Relation to Upper Limb Motor Recovery Early Post Stroke. Neurorehabilitation and Neural Repair, 2020, 34, 690-701.	2.9	4
126	Can telerehabilitation services combined with caregiver-mediated exercises improve early supported discharge services poststroke? A study protocol for a multicentre, observer-blinded, randomized controlled trial. BMC Neurology, 2022, 22, 29.	1.8	4

#	Article	IF	CITATIONS
127	Is it possible to accurately predict outcome of a drop-foot in patients admitted to a hospital stroke unit?. International Journal of Rehabilitation Research, 2013, 36, 346-353.	1.3	3
128	Quantifying neural and non-neural components of wrist hyper-resistance after stroke: Comparing two instrumented assessment methods. Medical Engineering and Physics, 2021, 98, 57-64.	1.7	3
129	The effect of cerebellar transcranial direct current stimulation to improve standing balance performance early post-stroke, study protocol of a randomized controlled trial. International Journal of Stroke, 2019, 14, 650-657.	5.9	2
130	The effect of botulinum toxinâ€A on neural and nonâ€neural components of wrist hyperâ€resistance in adults with stroke or cerebral palsy. PM and R, 2021, , .	1.6	2
131	In-Home Falls Risk Assessment in Parkinson Disease: A Guide for Clinicians. Archives of Physical Medicine and Rehabilitation, 2021, 102, 2051-2054.	0.9	2
132	Symptom dimensions of anxiety in Parkinson's disease: Replication study in a neuropsychiatric patient population. Clinical Parkinsonism & Related Disorders, 2021, 5, 100117.	0.9	2
133	Physical activity dimensions after stroke: patterns and relation with lower limb motor function. Journal of NeuroEngineering and Rehabilitation, 2021, 18, 171.	4.6	2
134	*Poster 63: Presence of Finger Extension and Shoulder Abduction Within 72 Hours Poststroke Predicts Functional Recovery. Archives of Physical Medicine and Rehabilitation, 2010, 91, e24.	0.9	1
135	Clinimetrics: The Neurological Fatigue Index for Multiple Sclerosis. Journal of Physiotherapy, 2019, 65, 241.	1.7	1
136	The Cortical Response Evoked by Robotic Wrist Perturbations Reflects Level of Proprioceptive Impairment After Stroke. Frontiers in Human Neuroscience, 2021, 15, 695366.	2.0	1
137	Graded peak cycle ergometer test for cognitively impaired patients with Parkinson's disease: a pilot study. Physiotherapy Theory and Practice, 2023, 39, 1249-1256.	1.3	1
138	Response to Letter by Corea et al. Stroke, 2010, 41, .	2.0	0
139	Allied Rehabilitation Using Web-Based Caregiver MEDiated Exercises for STROKE: The ARMED4STROKE Trial Design. Biosystems and Biorobotics, 2022, , 427-431.	0.3	0
140	7 Effecten van fysiotherapie en het gebruik van externe ritmische cues bij patiënten met de ziekte van Parkinson. , 2010, , 106-123.		0
141	Comparing two identically protocolized, multicentre, randomized controlled trials on caregiver-mediated exercises poststroke: Any differences across countries?. PLoS ONE, 2022, 17, e0263013.	2.5	0