

# Jerzy Romaszko

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6190197/publications.pdf>

Version: 2024-02-01

26  
papers

480  
citations

840776

11  
h-index

713466

21  
g-index

26  
all docs

26  
docs citations

26  
times ranked

703  
citing authors

#	ARTICLE	IF	CITATIONS
1	Are the levels of uric acid associated with biometeorological conditions?. Science of the Total Environment, 2022, 819, 152020.	8.0	1
2	Universal Climate Thermal Index as a prognostic tool in medical science in the context of climate change: A systematic review. Science of the Total Environment, 2022, 828, 154492.	8.0	19
3	The association between the biometeorological indicators and emergency interventions due to fainting: A retrospective cohort study. Science of the Total Environment, 2021, 770, 145376.	8.0	3
4	Assessment of the Consultation Rate with General Practitioners in the Initial Phase of the COVID-19 Pandemic. International Journal of Environmental Research and Public Health, 2020, 17, 7710.	2.6	9
5	Relationship between biometeorological factors and the number of hospitalizations due to asthma. Scientific Reports, 2020, 10, 9593.	3.3	24
6	UTCI as a bio-meteorological tool in the assessment of cold-induced stress as a risk factor for hypertension. Science of the Total Environment, 2019, 688, 970-975.	8.0	20
7	Applicability of the universal thermal climate index for predicting the outbreaks of respiratory tract infections: a mathematical modeling approach. International Journal of Biometeorology, 2019, 63, 1231-1241.	3.0	15
8	Are the Levels of Lipid Parameters Associated with Biometeorological Conditions?. International Journal of Environmental Research and Public Health, 2019, 16, 4636.	2.6	12
9	Spirometry testing among the homeless. Advances in Clinical and Experimental Medicine, 2018, 27, 689-693.	1.4	0
10	Mortality among the homeless: Causes and meteorological relationships. PLoS ONE, 2017, 12, e0189938.	2.5	55
11	Socioeconomic Characteristics, Health Risk Factors and Alcohol Consumption among the Homeless in North-Eastern Part of Poland. Central European Journal of Public Health, 2017, 25, 29-34.	1.1	9
12	Coping Styles and Alcohol Dependence among Homeless People. PLoS ONE, 2016, 11, e0162381.	2.5	22
13	Active Case Finding Among Homeless People as a Means of Reducing the Incidence of Pulmonary Tuberculosis in General Population. Advances in Experimental Medicine and Biology, 2016, 911, 67-76.	1.6	4
14	The impact of red cabbage fermentation on bioavailability of anthocyanins and antioxidant capacity of human plasma. Food Chemistry, 2016, 190, 730-740.	8.2	55
15	Exposure of breastfed infants to quercetin after consumption of a single meal rich in quercetin by their mothers. Molecular Nutrition and Food Research, 2014, 58, 221-228.	3.3	22
16	Artificial neural networks approach to early lung cancer detection. Open Medicine (Poland), 2014, 9, 632-641.	1.3	13
17	Prediction of antimicrobial activity of imidazole derivatives by artificial neural networks. Open Medicine (Poland), 2013, 8, 1-15.	1.3	9
18	An assessment of the duration of breastfeeding in north-eastern Poland. Open Medicine (Poland), 2013, 8, 75-79.	1.3	0

#	ARTICLE	IF	CITATIONS
19	The incidence of pulmonary tuberculosis among the homeless in north-eastern Poland. <i>Open Medicine (Poland)</i> , 2013, 8, 283-285.	1.3	9
20	Identification of clinical risk factors of atrial fibrillation in congestive heart failure. <i>Cardiology Journal</i> , 2013, 20, 364-369.	1.2	2
21	ANN as a prognostic tool after treatment of non-seminoma testicular cancer. <i>Open Medicine (Poland)</i> , 2012, 7, 672-679.	1.3	1
22	Indapamide-induced hyponatremia or the syndrome of inappropriate antidiuretic hormone secretion: a case report. <i>Open Medicine (Poland)</i> , 2011, 6, 407-410.	1.3	2
23	Evaluation of selected lipid parameters and blood pressure in ethnically-homogenous population of middle-aged persons, depending on gender, age and body mass. <i>Open Medicine (Poland)</i> , 2010, 5, 486-494.	1.3	1
24	Antihypertensive efficacy and tolerability of aliskiren/hydrochlorothiazide (HCT) single-pill combinations in patients who are non-responsive to HCT 25 mg alone. <i>Current Medical Research and Opinion</i> , 2009, 25, 903-910.	1.9	30
25	Quercetin from Shallots ( <i>Allium cepa</i> L. var. <i>aggregatum</i> ) Is More Bioavailable Than Its Glucosides , ,3. <i>Journal of Nutrition</i> , 2008, 138, 885-888.	2.9	141
26	Analysis of selected risk factors of coronary artery disease in a healthy population aged 35-55 years. <i>Kardiologia Polska</i> , 2007, 65, 1216-22; discussion 1223-4.	0.6	2