

Tetsuhiro Kasamatsu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6196975/publications.pdf>

Version: 2024-02-01

19
papers

206
citations

1163117

8
h-index

1125743

13
g-index

19
all docs

19
docs citations

19
times ranked

377
citing authors

#	ARTICLE	IF	CITATIONS
1	DNA-double strand breaks enhance the expression of major histocompatibility complex class II through the ATM-NF- κ B-IRF1-CLITA pathway. <i>Cancer Gene Therapy</i> , 2022, 29, 225-240.	4.6	10
2	IDO2 rs10109853 polymorphism affects the susceptibility to multiple myeloma. <i>Clinical and Experimental Medicine</i> , 2021, 21, 323-329.	3.6	5
3	PDCD1 and PDCD1LG1 polymorphisms affect the susceptibility to multiple myeloma. <i>Clinical and Experimental Medicine</i> , 2020, 20, 51-62.	3.6	15
4	Long Noncoding RNA PVT1 Is Regulated by Bromodomain Protein BRD4 in Multiple Myeloma and Is Associated with Disease Progression. <i>International Journal of Molecular Sciences</i> , 2020, 21, 7121.	4.1	16
5	PARP1 V762A polymorphism affects the prognosis of myelodysplastic syndromes. <i>European Journal of Haematology</i> , 2020, 104, 526-537.	2.2	5
6	<i>PDCD1</i> and <i>CTLA4</i> polymorphisms affect the susceptibility to, and clinical features of, chronic immune thrombocytopenia. <i>British Journal of Haematology</i> , 2018, 180, 705-714.	2.5	16
7	<i>IL17A</i> and <i>IL23R</i> gene polymorphisms affect the clinical features and prognosis of patients with multiple myeloma. <i>Hematological Oncology</i> , 2018, 36, 196-201.	1.7	10
8	Unsuppressed serum albumin levels may jeopardize the clinical relevance of the international staging system to patients with light chain myeloma. <i>Hematological Oncology</i> , 2018, 36, 792-800.	1.7	1
9	Association between OGG1 S326C CC genotype and elevated relapse risk in acute myeloid leukemia. <i>International Journal of Hematology</i> , 2018, 108, 246-253.	1.6	9
10	The cytokine polymorphisms affecting Th1/Th2 increase the susceptibility to, and severity of, chronic ITP. <i>BMC Immunology</i> , 2017, 18, 26.	2.2	37
11	Long non-coding RNA <i>MALAT1</i> is an inducible stress response gene associated with extramedullary spread and poor prognosis of multiple myeloma. <i>British Journal of Haematology</i> , 2017, 179, 449-460.	2.5	68
12	Polymorphism of IL-10 receptor β^2 affects the prognosis of multiple myeloma patients treated with thalidomide and/or bortezomib. <i>Hematological Oncology</i> , 2017, 35, 711-718.	1.7	7
13	Polymorphisms of <i>IL10</i> affect the severity and prognosis of myelodysplastic syndrome. <i>European Journal of Haematology</i> , 2016, 96, 245-251.	2.2	7
14	Elemental analysis of erythrocytes in patients with chronic liver diseases positive for hepatitis C virus. <i>International Journal of PIXE</i> , 2015, 25, 65-72.	0.4	0
15	Differences in Expression Patterns of DNMTs and TSG Proteins in Lymphoid Tissue Section Play an Important Role in Their Association. <i>Blood</i> , 2015, 126, 2657-2657.	1.4	0
16	The Influence of Polymorphisms of Interleukin-17F Genes on the Susceptibility and Clinical Significance of Chronic Idiopathic Thrombocytopenic Purpura. <i>Blood</i> , 2008, 112, 3428-3428.	1.4	0
17	IL-10-592 Polymorphism Predicts the Clinical Outcome of Japanese Patients with Multiple Myeloma and MGUS. <i>Blood</i> , 2007, 110, 4759-4759.	1.4	0
18	Interleukin-10 Gene Polymorphism Reflects the Severity of Chronic Idiopathic Thrombocytopenic Purpura. <i>Blood</i> , 2007, 110, 2111-2111.	1.4	0

#	ARTICLE	IF	CITATIONS
19	IL-10 Promoter Region -592A/C Genotype Increases the Risk of Multiple Myeloma in Japanese Patients.. Blood, 2006, 108, 5015-5015.	1.4	0