Helena Erlandsson-Harris

List of Publications by Citations

 $\textbf{Source:} \ https://exaly.com/author-pdf/6200670/helena-erlandsson-harris-publications-by-citations.pdf$

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

34 2,777 16 38 g-index

38 3,137 6.1 4.91 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
34	High mobility group 1 protein (HMG-1) stimulates proinflammatory cytokine synthesis in human monocytes. <i>Journal of Experimental Medicine</i> , 2000 , 192, 565-70	16.6	1194
33	HMGB1: a multifunctional alarmin driving autoimmune and inflammatory disease. <i>Nature Reviews Rheumatology</i> , 2012 , 8, 195-202	8.1	469
32	Mini-review: The nuclear protein HMGB1 as a proinflammatory mediator. <i>European Journal of Immunology</i> , 2004 , 34, 1503-12	6.1	311
31	Alarmin(g) news about danger: workshop on innate danger signals and HMGB1. <i>EMBO Reports</i> , 2006 , 7, 774-8	6.5	226
30	A systematic nomenclature for the redox states of high mobility group box (HMGB) proteins. <i>Molecular Medicine</i> , 2014 , 20, 135-7	6.2	83
29	A novel high mobility group box 1 neutralizing chimeric antibody attenuates drug-induced liver injury and postinjury inflammation in mice. <i>Hepatology</i> , 2016 , 64, 1699-1710	11.2	76
28	The immune response of the human brain to abdominal surgery. <i>Annals of Neurology</i> , 2017 , 81, 572-582	9.4	58
27	High systemic levels of the cytokine-inducing HMGB1 isoform secreted in severe macrophage activation syndrome. <i>Molecular Medicine</i> , 2015 , 20, 538-47	6.2	40
26	HMGB1 levels are increased in patients with juvenile idiopathic arthritis, correlate with early onset of disease, and are independent of disease duration. <i>Journal of Rheumatology</i> , 2013 , 40, 1604-13	4.1	40
25	Brain-released alarmins and stress response synergize in accelerating atherosclerosis progression after stroke. <i>Science Translational Medicine</i> , 2018 , 10,	17.5	37
24	Systemic TNF blockade does not modulate synovial expression of the pro-inflammatory mediator HMGB1 in rheumatoid arthritis patientsa prospective clinical study. <i>Arthritis Research and Therapy</i> , 2008 , 10, R33	5.7	30
23	Neuroinflammation in Response to Intracerebral Injections of Different HMGB1 Redox Isoforms. Journal of Innate Immunity, 2018 , 10, 215-227	6.9	28
22	Antibodies against High Mobility Group Box protein-1 (HMGB1) versus other anti-nuclear antibody fine-specificities and disease activity in systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , 2015 , 17, 338	5.7	26
21	Role of selectins in experimental Staphylococcus aureus-induced arthritis. <i>European Journal of Immunology</i> , 2000 , 30, 1606-13	6.1	25
20	Characterization of the Inflammatory Properties of Actively Released HMGB1 in Juvenile Idiopathic Arthritis. <i>Antioxidants and Redox Signaling</i> , 2016 , 24, 605-19	8.4	21
19	Ligation of free HMGB1 to TLR2 in the absence of ligand is negatively regulated by the C-terminal tail domain. <i>Molecular Medicine</i> , 2018 , 24, 19	6.2	18
18	Proteasome inhibitor MG132 modulates inflammatory pain by central mechanisms in adjuvant arthritis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 25-32	2.3	14

LIST OF PUBLICATIONS

17	Sex-dependent role of microglia in disulfide high mobility group box 1 protein-mediated mechanical hypersensitivity. <i>Pain</i> , 2021 , 162, 446-458	8	13
16	Neuroinflammatory markers associate with cognitive decline after major surgery: Findings of an explorative study. <i>Annals of Neurology</i> , 2020 , 87, 370-382	9.4	12
15	Fragmented hyaluronan has no alarmin function assessed in arthritis synovial fibroblast and chondrocyte cultures. <i>Innate Immunity</i> , 2018 , 24, 131-141	2.7	10
14	Sex- and cell-dependent contribution of peripheral high mobility group box 1 and TLR4 in arthritis-induced pain. <i>Pain</i> , 2021 , 162, 459-470	8	10
13	Differential bone turnover in an angulated fracture model in the rat. <i>Calcified Tissue International</i> , 2004 , 75, 50-9	3.9	7
12	Redox modifications of cysteine residues regulate the cytokine activity of HMGB1. <i>Molecular Medicine</i> , 2021 , 27, 58	6.2	7
11	The impact of damage-associated molecular patterns on the neurotransmitter release and gene expression in the ex vivo rat carotid body. <i>Experimental Physiology</i> , 2020 , 105, 1634-1647	2.4	5
10	Therapeutic administration of etoposide coincides with reduced systemic HMGB1 levels in macrophage activation syndrome. <i>Molecular Medicine</i> , 2021 , 27, 48	6.2	3
9	Disulfide and Fully Reduced HMGB1 Induce Different Macrophage Polarization and Migration Patterns. <i>Biomolecules</i> , 2021 , 11,	5.9	3
8	Cleavage of HMGB1 by Proteolytic Enzymes Associated with Inflammatory Conditions. <i>Frontiers in Immunology</i> , 2020 , 11, 448262	8.4	2
7	The pro-inflammatory effect of HMGB1, a mediator of inflammation in arthritis, is dependent on the redox status of the protein. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, A81.2-A82	2.4	1
6	Juvenile idiopathic arthritis patients have a distinct cartilage and bone biomarker profile that differs from healthy and knee-injured children. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 355-3	6 ² 5 ²	1
5	Immunoprofiling of active and inactive systemic juvenile idiopathic arthritis reveals distinct biomarkers: a single-center study <i>Pediatric Rheumatology</i> , 2021 , 19, 173	3.5	1
4	Transcriptomic Profiling Reveals That HMGB1 Induces Macrophage Polarization Different from Classical M1. <i>Biomolecules</i> , 2022 , 12, 779	5.9	O
3	Successful therapy with anti-HMGB1 monoclonal antibodies in two separate experimental arthritis models. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, A77-A78	2.4	
2	Crosstalk between nitrosative stress and altered Ca2+ handling in arthritis-induced skeletal muscle dysfunction. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, A43.3-A44	2.4	
1	HMGB1 mediates muscle fatigue via TLR4 - a possible mechanism for muscle fatigue in patients with inflammatory myopathies. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, A42.2-A43	2.4	