

Kwan-Woo Lee

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6207572/publications.pdf>

Version: 2024-02-01

37
papers

762
citations

687363

13
h-index

552781

26
g-index

38
all docs

38
docs citations

38
times ranked

1279
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Characteristics and Clinical Course of Diabetes of the Exocrine Pancreas: A Nationwide Population-Based Cohort Study. <i>Diabetes Care</i> , 2022, 45, 1141-1150. | 8.6 | 3 |
| 2 | Visfatin exacerbates hepatic inflammation and fibrosis in a methionineâ€cholineâ€deficient diet mouse model. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021, 36, 2592-2600. | 2.8 | 14 |
| 3 | Efficacy and Safety of Switching to Tenziglipitin in Patients with Type 2 Diabetes Inadequately Controlled with Dipeptidyl Peptidase-4 Inhibitors: 52-Week Results from a Prospective Observational Study. <i>Diabetes Therapy</i> , 2021, 12, 2907-2920. | 2.5 | 3 |
| 4 | Pathways Linking Health Literacy to Self-Management in People with Type 2 Diabetes. <i>Healthcare (Switzerland)</i> , 2021, 9, 1734. | 2.0 | 5 |
| 5 | A New Objective Health Numeracy Test for Patients with Type 2 Diabetes: Development and Evaluation of Psychometric Properties. <i>Asian Nursing Research</i> , 2020, 14, 66-72. | 1.4 | 4 |
| 6 | Clinical Characteristics and Prevalence of Comorbidities according to Metformin Use in Korean Patients with Type 2 Diabetes. <i>International Journal of Endocrinology</i> , 2020, 2020, 1-7. | 1.5 | 5 |
| 7 | A New Self-management Scale with a Hierarchical Structure for Patients with Type 2 Diabetes. <i>Asian Nursing Research</i> , 2020, 14, 249-256. | 1.4 | 3 |
| 8 | CCL20 induced by visfatin in macrophages via the NF- κ B and MKK3/6-p38 signaling pathways contributes to hepatic stellate cell activation. <i>Molecular Biology Reports</i> , 2020, 47, 4285-4293. | 2.3 | 10 |
| 9 | Visfatin Induces Inflammation and Insulin Resistance via the NF- κ B and STAT3 Signaling Pathways in Hepatocytes. <i>Journal of Diabetes Research</i> , 2019, 2019, 1-11. | 2.3 | 89 |
| 10 | Efficacy and Safety of Switching to Tenziglipitin in Patients with Type 2 Diabetes Inadequately Controlled with Dipeptidyl Peptidase-4 Inhibitors: A 12-Week Interim Report. <i>Diabetes Therapy</i> , 2019, 10, 1271-1282. | 2.5 | 8 |
| 11 | Iron overload by transferrin receptor protein 1 regulation plays an important role in palmitateâ€induced insulin resistance in human skeletal muscle cells. <i>FASEB Journal</i> , 2019, 33, 1771-1786. | 0.5 | 23 |
| 12 | Risk of Dementia in Older Patients with Type 2 Diabetes on Dipeptidyl-Peptidase IV Inhibitors Versus Sulfonylureas: A Real-World Population-Based Cohort Study. <i>Journal of Clinical Medicine</i> , 2019, 8, 28. | 2.4 | 23 |
| 13 | Predictive Factors for Efficacy of AST-120 Treatment in Diabetic Nephropathy: a Prospective Single-Arm, Open-Label, Multi-Center Study. <i>Journal of Korean Medical Science</i> , 2019, 34, e117. | 2.5 | 7 |
| 14 | Failure of monotherapy in clinical practice in patients with type 2 diabetes: The Korean National Diabetes Program. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1144-1152. | 2.4 | 10 |
| 15 | Importance of family history of diabetes in computing a diabetes risk score in Korean prediabetic population. <i>Scientific Reports</i> , 2018, 8, 15958. | 3.3 | 9 |
| 16 | Blood adiponectin levels are not associated with risk of cardiovascular events in patients with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 571-575. | 2.0 | 2 |
| 17 | Dipeptidyl Peptidase-4 Inhibitors and the Risk of Pancreatitis in Patients with Type 2 Diabetes Mellitus: A Population-Based Cohort Study. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-10. | 2.3 | 10 |
| 18 | A new comprehensive diabetes health literacy scale: Development and psychometric evaluation. <i>International Journal of Nursing Studies</i> , 2018, 88, 1-8. | 5.6 | 29 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Efficacy and safety of ipragliflozin as an add-on therapy to sitagliptin and metformin in Korean patients with inadequately controlled type 2 diabetes mellitus: A randomized controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 2408-2415. | 4.4 | 30 |
| 20 | Effects of foot complications in patients with Type 2 diabetes mellitus on public healthcare: An analysis based on the Korea National Diabetes Program Cohort. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 375-380. | 2.3 | 3 |
| 21 | Association between insulin resistance and impairment of FGF21 signal transduction in skeletal muscles. <i>Endocrine</i> , 2016, 53, 97-106. | 2.3 | 18 |
| 22 | Impact of Cadmium Exposure on the Association between Lipopolysaccharide and Metabolic Syndrome. <i>International Journal of Environmental Research and Public Health</i> , 2015, 12, 11396-11409. | 2.6 | 14 |
| 23 | A Korean Version of the Appraisal of Diabetes Scale (ADS-K). <i>Journal of Transcultural Nursing</i> , 2015, 26, 270-278. | 1.3 | 8 |
| 24 | Clinical and Economic Outcomes in Medication-Adherent and -Nonadherent Patients With Type 2 Diabetes Mellitus in the Republic of Korea. <i>Clinical Therapeutics</i> , 2014, 36, 245-254. | 2.5 | 10 |
| 25 | IL-6 induction of TLR-4 gene expression via STAT3 has an effect on insulin resistance in human skeletal muscle. <i>Acta Diabetologica</i> , 2013, 50, 189-200. | 2.5 | 59 |
| 26 | A compound (DW1182v) protecting high glucose/palmitate-induced glucolipotoxicity to INS-1 beta cells preserves islet integrity and improves hyperglycemia in obese db/db mouse. <i>European Journal of Pharmacology</i> , 2012, 696, 187-193. | 3.5 | 4 |
| 27 | Association between Nutrient Intake and Obesity in Type 2 Diabetic Patients from the Korean National Diabetes Program: A Cross-Sectional Study. <i>Journal of Korean Medical Science</i> , 2012, 27, 1188. | 2.5 | 13 |
| 28 | Supplementation of pyruvate prevents palmitate-induced impairment of glucose uptake in C2 myotubes. <i>Molecular and Cellular Endocrinology</i> , 2011, 345, 79-87. | 3.2 | 5 |
| 29 | Intake of brown rice lees reduces waist circumference and improves metabolic parameters in type 2 diabetes. <i>Nutrition Research</i> , 2011, 31, 131-138. | 2.9 | 24 |
| 30 | Capsaicin attenuates palmitate-induced expression of macrophage inflammatory protein 1 and interleukin 8 by increasing palmitate oxidation and reducing c-Jun activation in THP-1 (human acute) Tj ETQq0 0 0 r2BT /Overlock 10 Tf 5 | 2.9 | 4 |
| 31 | Fermented kimchi reduces body weight and improves metabolic parameters in overweight and obese patients. <i>Nutrition Research</i> , 2011, 31, 436-443. | 2.9 | 131 |
| 32 | Outcome Research in Diabetes. <i>Journal of Korean Diabetes</i> , 2011, 12, 2. | 0.3 | 3 |
| 33 | Prevalence of Chronic Complications in Korean Patients with Type 2 Diabetes Mellitus Based on the Korean National Diabetes Program. <i>Diabetes and Metabolism Journal</i> , 2011, 35, 504. | 4.7 | 56 |
| 34 | Comparison of the Efficacy of Glimepiride, Metformin, and Rosiglitazone Monotherapy in Korean Drug-Naïve Type 2 Diabetic Patients: The Practical Evidence of Antidiabetic Monotherapy Study. <i>Diabetes and Metabolism Journal</i> , 2011, 35, 26. | 4.7 | 32 |
| 35 | Atherosclerosis induced by a high-fat diet is alleviated by lithium chloride via reduction of VCAM expression in ApoE-deficient mice. <i>Vascular Pharmacology</i> , 2010, 53, 264-272. | 2.1 | 45 |
| 36 | Comparison of the Efficacy and Safety of Glimepiride/Metformin Fixed Combination Versus Free Combination in Patients with Type 2 Diabetes: Multicenter, Randomized, Controlled Trial. <i>The Journal of Korean Diabetes Association</i> , 2006, 30, 466. | 0.1 | 3 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Efficacy Evaluation of Atorvastatin in Korean Hyperlipidemic Patients with Type 2 Diabetes Mellitus. The Journal of Korean Diabetes Association, 2006, 30, 292. | 0.1 | 3 |