

Susumu Itoh

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6208077/susumu-itoh-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

95
papers

8,694
citations

37
h-index

93
g-index

98
ext. papers

9,213
ext. citations

6.1
avg, IF

5.35
L-index

#	Paper	IF	Citations
95	Endothelial-specific depletion of TGF- β signaling affects lymphatic function. <i>Inflammation and Regeneration</i> , 2021 , 41, 35	10.9	1
94	Narciclasine is a novel YAP inhibitor that disturbs interaction between YAP and TEAD4. <i>BBA Advances</i> , 2021 , 1, 100008		0
93	Involvement of miR-3180-3p and miR-4632-5p in palmitic acid-induced insulin resistance. <i>Molecular and Cellular Endocrinology</i> , 2021 , 534, 111371	4.4	2
92	Dissociation of the AhR/ARNT complex by TGF- β /Smad signaling represses gene expression and inhibits benze[a]pyrene-mediated cytotoxicity. <i>Journal of Biological Chemistry</i> , 2020 , 295, 9033-9051	5.4	10
91	Smad7 Enhances TGF- β -Induced Transcription of c-Jun and HDAC6 Promoting Invasion of Prostate Cancer Cells. <i>IScience</i> , 2020 , 101470	6.1	6
90	PDZK1-interacting protein 1 (PDZK1IP1) traps Smad4 protein and suppresses transforming growth factor- β (TGF- β) signaling. <i>Journal of Biological Chemistry</i> , 2019 , 294, 4966-4980	5.4	9
89	TMEPAI family: involvement in regulation of multiple signalling pathways. <i>Journal of Biochemistry</i> , 2018 , 164, 195-204	3.1	16
88	TMED10 Protein Interferes with Transforming Growth Factor (TGF)- β Signaling by Disrupting TGF- β Receptor Complex Formation. <i>Journal of Biological Chemistry</i> , 2017 , 292, 4099-4112	5.4	11
87	Targeted Degradation of Proteins Localized in Subcellular Compartments by Hybrid Small Molecules. <i>Molecular Pharmacology</i> , 2017 , 91, 159-166	4.3	38
86	TGF- β Promotes PI3K-AKT signaling and prostate cancer cell migration through the TRAF6-mediated ubiquitylation of p85. <i>Science Signaling</i> , 2017 , 10,	8.8	111
85	Regulation of the TMEPAI promoter by TCF7L2: the C-terminal tail of TCF7L2 is essential to activate the TMEPAI gene. <i>Journal of Biochemistry</i> , 2016 , 159, 27-30	3.1	8
84	Arf6 regulates tumour angiogenesis and growth through HGF-induced endothelial β 1 integrin recycling. <i>Nature Communications</i> , 2015 , 6, 7925	17.4	39
83	Transforming growth factor- β signaling enhancement by long-term exposure to hypoxia in a tumor microenvironment composed of Lewis lung carcinoma cells. <i>Cancer Science</i> , 2015 , 106, 1524-33	6.9	20
82	Active TGF- β signaling in hypoxic area. <i>Cancer Science</i> , 2015 , 106, November cover-November cover	6.9	78
81	TMEPAI/PMEPA1 enhances tumorigenic activities in lung cancer cells. <i>Cancer Science</i> , 2014 , 105, 334-41	6.9	47
80	C18 ORF1, a novel negative regulator of transforming growth factor- β signaling. <i>Journal of Biological Chemistry</i> , 2014 , 289, 12680-92	5.4	35
79	TGF- β Signaling Cooperates with AT Motif-Binding Factor-1 for Repression of the β 2-Microglobulin Promoter. <i>Journal of Signal Transduction</i> , 2014 , 2014, 970346		6

78	Smad2/Smad3 in endothelium is indispensable for vascular stability via S1PR1 and N-cadherin expressions. <i>Blood</i> , 2012 , 119, 5320-8	2.2	49
77	Implication of TGF- β s as a survival factor during tumour development. <i>Journal of Biochemistry</i> , 2012 , 151, 559-62	3.1	13
76	Delayed cutaneous wound healing in Fam129b/Minerva-deficient mice. <i>Journal of Biochemistry</i> , 2012 , 152, 549-55	3.1	4
75	Interference of E2-2-mediated effect in endothelial cells by FAM96B through its limited expression of E2-2. <i>Cancer Science</i> , 2011 , 102, 1808-14	6.9	7
74	Inhibitory machinery for the TGF- β -family signaling pathway. <i>Growth Factors</i> , 2011 , 29, 163-73	1.6	19
73	Requirement of TCF7L2 for TGF-beta-dependent transcriptional activation of the TMEPAI gene. <i>Journal of Biological Chemistry</i> , 2010 , 285, 38023-33	5.4	39
72	TMEPAI, a transmembrane TGF-beta-inducible protein, sequesters Smad proteins from active participation in TGF-beta signaling. <i>Molecular Cell</i> , 2010 , 37, 123-34	17.6	111
71	Flk1-GFP BAC Tg mice: an animal model for the study of blood vessel development. <i>Experimental Animals</i> , 2010 , 59, 615-22	1.8	31
70	Inhibition of endothelial cell activation by bHLH protein E2-2 and its impairment of angiogenesis. <i>Blood</i> , 2010 , 115, 4138-47	2.2	29
69	Poor vessel formation in embryos from knock-in mice expressing ALK5 with L45 loop mutation defective in Smad activation. <i>Laboratory Investigation</i> , 2009 , 89, 800-10	5.9	17
68	TAL1/SCL relieves the E2-2-mediated repression of VEGFR2 promoter activity. <i>Journal of Biochemistry</i> , 2009 , 145, 129-35	3.1	11
67	Negative Regulation of the TGF- β -Family Signal Pathway by Inhibitory Smads and Their Involvement in Cancer and Fibrosis 2008 , 649-661		
66	Negative regulation of TGF-beta receptor/Smad signal transduction. <i>Current Opinion in Cell Biology</i> , 2007 , 19, 176-84	9	314
65	Compensatory signalling induced in the yolk sac vasculature by deletion of TGFbeta receptors in mice. <i>Journal of Cell Science</i> , 2007 , 120, 4269-77	5.3	94
64	Methylation of Smad6 by protein arginine N-methyltransferase 1. <i>FEBS Letters</i> , 2006 , 580, 6603-11	3.8	26
63	Smad7 and protein phosphatase 1alpha are critical determinants in the duration of TGF-beta/ALK1 signaling in endothelial cells. <i>BMC Cell Biology</i> , 2006 , 7, 16		43
62	Synergy and antagonism between Notch and BMP receptor signaling pathways in endothelial cells. <i>EMBO Journal</i> , 2004 , 23, 541-51	13	208
61	Activin receptor-like kinase (ALK)1 is an antagonistic mediator of lateral TGFbeta/ALK5 signaling. <i>Molecular Cell</i> , 2003 , 12, 817-28	17.6	561

60	Elucidation of Smad requirement in transforming growth factor-beta type I receptor-induced responses. <i>Journal of Biological Chemistry</i> , 2003 , 278, 3751-61	5.4	171
59	Regulation of cell proliferation by Smad proteins. <i>Journal of Cellular Physiology</i> , 2002 , 191, 1-16	7	368
58	The FYVE domain in Smad anchor for receptor activation (SARA) is sufficient for localization of SARA in early endosomes and regulates TGF-beta/Smad signalling. <i>Genes To Cells</i> , 2002 , 7, 321-31	2.3	126
57	Balancing the activation state of the endothelium via two distinct TGF-beta type I receptors. <i>EMBO Journal</i> , 2002 , 21, 1743-53	13	847
56	Stimulation of Id1 expression by bone morphogenetic protein is sufficient and necessary for bone morphogenetic protein-induced activation of endothelial cells. <i>Circulation</i> , 2002 , 106, 2263-70	16.7	264
55	Signal transduction mechanisms for members of the TGF- β family 2001 , 11-40		3
54	Signaling of transforming growth factor-beta family members through Smad proteins. <i>FEBS Journal</i> , 2000 , 267, 6954-67		398
53	Functional consequences of tumorigenic missense mutations in the amino-terminal domain of Smad4. <i>Oncogene</i> , 2000 , 19, 4396-404	9.2	81
52	Smad7 mediates apoptosis induced by transforming growth factor beta in prostatic carcinoma cells. <i>Current Biology</i> , 2000 , 10, 535-8	6.3	141
51	Smad and AML proteins synergistically confer transforming growth factor beta1 responsiveness to human germ-line IgA genes. <i>Journal of Biological Chemistry</i> , 2000 , 275, 3552-60	5.4	125
50	Molecular cloning and characterization of a novel human STE20-like kinase, hSLK. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2000 , 1495, 250-62	4.9	23
49	Hedgehog creates a gradient of DPP activity in Drosophila wing imaginal discs. <i>Molecular Cell</i> , 2000 , 5, 59-71	17.6	324
48	Xenopus Smad4beta is the co-Smad component of developmentally regulated transcription factor complexes responsible for induction of early mesodermal genes. <i>Developmental Biology</i> , 1999 , 214, 354-69	3.1	85
47	Molecular cloning and regulation of a novel guinea pig cytochrome P450 (CYP3A20) which differs from guinea pig CYP3A14 in only two amino acid residues. <i>IUBMB Life</i> , 1998 , 44, 1245-53	4.7	
46	Intracellular signaling of osteogenic protein-1 through Smad5 activation. <i>Journal of Cellular Physiology</i> , 1998 , 177, 355-63	7	67
45	The L45 loop in type I receptors for TGF-beta family members is a critical determinant in specifying Smad isoform activation. <i>FEBS Letters</i> , 1998 , 434, 83-7	3.8	327
44	Induction of inhibitory Smad6 and Smad7 mRNA by TGF-beta family members. <i>Biochemical and Biophysical Research Communications</i> , 1998 , 249, 505-11	3.4	306
43	Effect of 26,26,26,27,27,27-Hexafluoro-1,25-dihydroxyvitamin D3 on the expression of vitamin-D-responsive genes in vitamin-D-deficient mice. <i>Pharmacology</i> , 1998 , 57, 286-94	2.3	4

42	Identification and functional characterization of a Smad binding element (SBE) in the JunB promoter that acts as a transforming growth factor-beta, activin, and bone morphogenetic protein-inducible enhancer. <i>Journal of Biological Chemistry</i> , 1998 , 273, 21145-52	5.4	482
41	Transforming growth factor beta1 induces nuclear export of inhibitory Smad7. <i>Journal of Biological Chemistry</i> , 1998 , 273, 29195-201	5.4	198
40	Inhibition of the transcription of CYP1A1 gene by the upstream stimulatory factor 1 in rabbits. Competitive binding of USF1 with AhR.Arnt complex. <i>Journal of Biological Chemistry</i> , 1997 , 272, 30025-31	5.4	39
39	Gene structure of mouse Cyp3a11: evidence for an enhancer element within its 5'flanking sequences. <i>Archives of Biochemistry and Biophysics</i> , 1997 , 338, 43-9	4.1	9
38	Determination of FAD-binding domain in flavin-containing monooxygenase 1 (FMO1). <i>Archives of Biochemistry and Biophysics</i> , 1997 , 345, 271-7	4.1	23
37	Regulation of CYP1A and CYP3A mRNAs by ascorbic acid in guinea pigs. <i>Archives of Biochemistry and Biophysics</i> , 1997 , 348, 268-77	4.1	22
36	Molecular cloning and characterization of a novel putative STE20-like kinase in guinea pigs. <i>Archives of Biochemistry and Biophysics</i> , 1997 , 340, 201-7	4.1	21
35	Mouse cytochrome P450 (Cyp3a11): predominant expression in liver and capacity to activate aflatoxin B1. <i>Archives of Biochemistry and Biophysics</i> , 1997 , 340, 215-8	4.1	64
34	Upstream stimulatory factor 1 (USF1) suppresses induction of CYP1A1 mRNA by 3-methylcholanthrene (MC) in HepG2 cells. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 240, 293-7	3.4	15
33	Simultaneous expression of ferredoxin, ferredoxin reductase and P450 in COS7 cells. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1997 , 1318, 284-90	4.6	1
32	Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. <i>Nature</i> , 1997 , 389, 631-5	50.4	1555
31	Isolation of a promoter region in mouse cytochrome P450 3A (Cyp3A16) gene and its transcriptional control. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1997 , 1350, 155-8		6
30	Opposite effects of isoniazid and fasting on the expression of CYP2E1 protein and mRNA in house musk shrew (<i>Suncus murinus</i>). <i>IUBMB Life</i> , 1997 , 41, 293-301	4.7	
29	Expression of aryl hydrocarbon receptor (AhR) and aryl hydrocarbon receptor nuclear translocator (Arnt) in adult rabbits known to be non-responsive to cytochrome P-450 1A1 (CYP1A1) inducers. <i>FEBS Journal</i> , 1996 , 242, 512-8		20
28	Molecular cloning of 25-hydroxyvitamin D-3 24-hydroxylase (Cyp-24) from mouse kidney: its inducibility by vitamin D-3. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1995 , 1264, 26-8		19
27	cDNA cloning of mouse ferredoxin reductase from kidney. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1995 , 1264, 159-62		3
26	N-oxygenation of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine by the rat liver flavin-containing monooxygenase expressed in yeast cells. <i>European Journal of Pharmacology - Environmental Toxicology and Pharmacology Section</i> , 1995 , 293, 97-100		6
25	Simultaneous expression of human CYP3A7 and N-acetyltransferase in Chinese hamster CHL cells results in high cytotoxicity for carcinogenic heterocyclic amines. <i>Archives of Biochemistry and Biophysics</i> , 1995 , 320, 323-9	4.1	16

24	Evidence for the lack of hepatic N-acetyltransferase in suncus (<i>Suncus murinus</i>). <i>Biochemical Pharmacology</i> , 1995 , 50, 1165-70	6	11
23	Pharmacological activity of chemically modified subfragment from human serum IgG. XIV. Inhibitory effect of carboxamide-methylated light chain (G1L) on tyrosine phosphorylation and tumor necrosis factor- α production from murine macrophages stimulated by lipopolysaccharide. <i>Biological and Pharmaceutical Bulletin</i> , 1995 , 18, 1377-81	2.3	
22	Adhesion molecules on eosinophils in acute eosinophilic pneumonia. <i>American Journal of Respiratory and Critical Care Medicine</i> , 1995 , 151, 1259-62	10.2	28
21	A novel form of mouse cytochrome P450 3A (Cyp3a-16). Its cDNA cloning and expression in fetal liver. <i>FEBS Journal</i> , 1994 , 226, 877-82		38
20	Molecular cloning and functional expression of a mouse cytochrome P-450 (Cyp3a-13): examination of Cyp3a-13 enzyme to activate aflatoxin B1 (AFB1). <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1994 , 1201, 405-10	4	27
19	Studies on thermophile products. X. Further biological properties of isofatty acid-containing phosphatidylglycerol that enhances the induction of suppressor T cells. <i>Biological and Pharmaceutical Bulletin</i> , 1994 , 17, 1446-50	2.3	
18	Studies on thermophile products. VII. Effect of 1,3-di-14-methylpentadecanoyl glycerol and its related isofatty acids on T cell proliferation in vitro. <i>Biological and Pharmaceutical Bulletin</i> , 1994 , 17, 850-2 ³		
17	Inhibitory effect of tuna peptide on endothelin production in cultured endothelial cells. <i>Biological and Pharmaceutical Bulletin</i> , 1994 , 17, 886-8	2.3	4
16	Studies on thermophile products. VIII. Isolation of <i>Bacillus stearothermophilus</i> UBT8038, a component that inhibits antigen presentation on mouse macrophages. <i>Biological and Pharmaceutical Bulletin</i> , 1994 , 17, 889-93	2.3	
15	Studies on thermophile products. IX. Isofatty acid-containing phosphatidylglycerol that enhances the induction of concanavalin A-activated suppressor T cells. <i>Biological and Pharmaceutical Bulletin</i> , 1994 , 17, 1171-5	2.3	1
14	Characterization of Ah receptor promoter in human liver cell line, HepG2. <i>Pharmacogenetics and Genomics</i> , 1994 , 4, 219-22		18
13	Human Ah receptor cDNA: analysis for highly conserved sequences. <i>Nucleic Acids Research</i> , 1993 , 21, 3578	20.1	17
12	Gene structure of CYP3A4, an adult-specific form of cytochrome P450 in human livers, and its transcriptional control. <i>FEBS Journal</i> , 1993 , 218, 585-95		158
11	Rat liver flavin-containing monooxygenase (FMO): cDNA cloning and expression in yeast. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1993 , 1173, 165-71		37
10	Stable expression of cytochrome P450III A7 cDNA in human breast cancer cell line MCF-7 and its application to cytotoxicity testing. <i>Archives of Biochemistry and Biophysics</i> , 1992 , 292, 136-40	4.1	25
9	Decrease in the content of cytochrome P450IIE by fasting in liver microsomes of house musk shrew (<i>Suncus murinus</i>). <i>Biochemical Pharmacology</i> , 1992 , 43, 1907-10	6	8
8	Mouse liver cytochrome P-450 (P-450IIIA M1): its cDNA cloning and inducibility by dexamethasone. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1992 , 1130, 329-32		54
7	Genomic organization of human fetal specific P-450III A7 (cytochrome P-450HFLa)-related gene(s) and interaction of transcriptional regulatory factor with its DNA element in the 5' flanking region. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1992 , 1130, 133-8		36

- | | | | |
|---|--|-----|----|
| 6 | Assignment of the human interferon regulatory factor-1 (IRF1) gene to chromosome 5q23-q31. <i>Genomics</i> , 1991 , 10, 1097-9 | 4.3 | 45 |
| 5 | Ligand-dependent selection of the receptor gene: segregation of IL-2 binding activity and anti-Tac reactivity by a single amino acid alteration in the Tac antigen (p55). <i>Immunology Letters</i> , 1989 , 20, 139-47 ^{4.1} | | 2 |
| 4 | Inhibitory effect of bis[2-(E-2-octenylamino)ethyl] disulfide and 2-(E-octenylamino)ethyl carbamoylmethyl sulfide on various inflammation models. <i>Chemical and Pharmaceutical Bulletin</i> , 1987 , 35, 4585-91 | 1.9 | |
| 3 | Studies on biological activities of melanin from marine animals. V. Anti-inflammatory activity of low-molecular-weight melanoprotein from squid (Fr. SM II). <i>Chemical and Pharmaceutical Bulletin</i> , 1987 , 35, 1144-50 | 1.9 | 13 |
| 2 | Inhibitory effect of bis[2-(E-2-alkenylaminoethyl] disulfides and 2-(E-octenylamino)ethyl carbamoylmethyl sulfides on carrageenin-induced paw edema in rats. <i>Chemical and Pharmaceutical Bulletin</i> , 1987 , 35, 4579-84 | 1.9 | |
| 1 | Studies on pharmacological activation of human serum immunoglobulin G (IgG) by chemical modification and active subfragments. VI. Anti-allergic activity of carboxamidemethylated Fc (CM-Fc) fragment from human serum IgG. <i>Chemical and Pharmaceutical Bulletin</i> , 1987 , 35, 4935-9 | 1.9 | |