
Ismail Omer Yilmaz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6208191/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Cenomanian – Turonian Oceanic Anoxic Event (OAE2) in the Sakarya Zone, northwestern Turkey: Sedimentological, cyclostratigraphic, and geochemical records. Cretaceous Research, 2010, 31, 207-226.	1.4	34

A 2800-year multi-proxy sedimentary record of climate change from Lake \tilde{A} [‡]ubuk (G \tilde{A} ¶yn \tilde{A} ¹/4k, Bolu, NW) Tj ETQq0000 rgBT $\frac{10}{26}$ verlock 1

3	Cyclic palaeokarst surfaces in Aptian peritidal carbonate successions (Taurides, southwest Turkey): internal structure and response to mid-Aptian sea-level fall. Cretaceous Research, 2006, 27, 814-827.	1.4	24
4	Foraminiferal biostratigraphy and sequence stratigraphy across the mid-Carboniferous boundary in the Central Taurides, Turkey. Facies, 2011, 57, 705-730.	1.4	18
5	Chronology of subduction and collision along the İzmir-Ankara suture in Western Anatolia: records from the Central Sakarya Basin. International Geology Review, 2019, 61, 1244-1269.	2.1	15
6	Use of Sedimentary Structures in the Recognition of Sequence Boundaries in Upper Jurassic–Upper Cretaceous Peritidal Carbonates of the Western Taurides, Turkey. International Geology Review, 2001, 43, 736-753.	2.1	13
7	Roveacrinid microfacial assemblages (Roveacrinida, Crinoidea) from the Lower-Middle Cenomanian of the Adıyaman area (SE Turkey). Arabian Journal of Geosciences, 2018, 11, 1.	1.3	5
8	Holocene paleoenvironmental evolution of the Acıgöl paleo maar lake (Nevşehir, Central Anatolia). Mediterranean Geoscience Reviews, 2019, 1, 255-269.	1.2	5
9	Microfacies and stable isotope analyses (δ13C and δ18O) of the upper Serpukhovian-Moscovian carbonates in the Hadim Nappe, southern Turkey: an approach to document the Late Paleozoic Ice Age (LPIA). Facies, 2022, 68, .	1.4	1