

Tayo Katano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6209405/publications.pdf>

Version: 2024-02-01

15
papers

523
citations

933447

10
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

759
citing authors

#	ARTICLE	IF	CITATIONS
1	Efficacy of Nanofiber Sheets Incorporating Lenvatinib in a Hepatocellular Carcinoma Xenograft Model. <i>Nanomaterials</i> , 2022, 12, 1364.	4.1	3
2	Combination therapy with lenvatinib and radiation significantly inhibits thyroid cancer growth by uptake of tyrosine kinase inhibitor. <i>Experimental Cell Research</i> , 2021, 398, 112390.	2.6	13
3	Diabetic neuropathy research: from mouse models to targets for treatment. <i>Neural Regeneration Research</i> , 2019, 14, 1870.	3.0	19
4	Impaired peripheral nerve regeneration in type 2 diabetic mouse model. <i>European Journal of Neuroscience</i> , 2018, 47, 126-139.	2.6	25
5	Distribution of Caskin1 protein and phenotypic characterization of its knockout mice using a comprehensive behavioral test battery. <i>Molecular Brain</i> , 2018, 11, 63.	2.6	28
6	Two isoforms of cyclic GMP-dependent kinase-I exhibit distinct expression patterns in the adult mouse dorsal root ganglion. <i>Molecular Pain</i> , 2018, 14, 174480691879640.	2.1	5
7	Na ⁺ /K ⁺ -ATPase coupled to endothelin receptor type B stimulates peripheral nerve regeneration via lactate signalling. <i>European Journal of Neuroscience</i> , 2017, 46, 2096-2107.	2.6	9
8	Role of c-Jun N-terminal kinase in late nerve regeneration monitored by <i>in vivo</i> imaging of thyl ¹ yellow fluorescent protein transgenic mice. <i>European Journal of Neuroscience</i> , 2016, 43, 548-560.	2.6	12
9	Involvement of Brain-Enriched Guanylate Kinase-Associated Protein (BEGAIN) in Chronic Pain after Peripheral Nerve Injury. <i>ENeuro</i> , 2016, 3, ENEURO.0110-16.2016.	1.9	10
10	Involvement of endothelin B receptor in peripheral nerve regeneration using sciatic nerve transection-regeneration model. <i>Pain Research</i> , 2015, 30, 167-172.	0.1	3
11	Involvement of spinal phosphorylation cascade of Tyr1472-NR2B, Thr286-CaMKII, and Ser831-GluR1 in neuropathic pain. <i>Neuropharmacology</i> , 2011, 60, 609-616.	4.1	63
12	Impairment of CaMKII activation and attenuation of neuropathic pain in mice lacking NR2B phosphorylated at Tyr1472. <i>European Journal of Neuroscience</i> , 2010, 32, 798-810.	2.6	57
13	<i>N</i> -ethylmaleimide-sensitive fusion protein (NSF) is involved in central sensitization in the spinal cord through GluR2 subunit composition switch after inflammation. <i>European Journal of Neuroscience</i> , 2008, 27, 3161-3170.	2.6	77
14	Proteomic identification of a novel isoform of collapsin response mediator protein-2 in spinal nerves peripheral to dorsal root ganglia. <i>Proteomics</i> , 2006, 6, 6085-6094.	2.2	41
15	Fyn kinase-mediated phosphorylation of NMDA receptor NR2B subunit at Tyr1472 is essential for maintenance of neuropathic pain. <i>European Journal of Neuroscience</i> , 2005, 22, 1445-1454.	2.6	158