## Flavia Trettel

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6215779/publications.pdf

Version: 2024-02-01

185998 205818 3,232 48 28 48 citations h-index g-index papers 49 49 49 4361 docs citations times ranked citing authors all docs

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Dominant phenotypes produced by the HD mutation in STHdhQ111 striatal cells. Human Molecular Genetics, 2000, 9, 2799-2809.   | 1.4 | 556       |
| 2  | Episodic Ataxia Type 2 (EA2) and Spinocerebellar Ataxia Type 6 (SCA6) Due to CAG Repeat Expansion in the CACNA1A Gene on Chromosome 19p. Human Molecular Genetics, 1997, 6, 1973-1978.                                   | 1.4 | 264       |
| 3  | Specific progressive cAMP reduction implicates energy deficit in presymptomatic Huntington's disease knock-in mice. Human Molecular Genetics, 2003, 12, 497-508.   | 1.4 | 250       |
| 4  | Huntingtin: an iron-regulated protein essential for normal nuclear and perinuclear organelles.<br>Human Molecular Genetics, 2000, 9, 2789-2797.  | 1.4 | 193       |
| 5  | Complete Loss of P/Q Calcium Channel Activity Caused by a CACNA1A Missense Mutation Carried by Patients with Episodic Ataxia Type 2. American Journal of Human Genetics, 2001, 68, 759-764.                              | 2.6 | 147       |
| 6  | TBX-3, the Gene Mutated in Ulnar-Mammary Syndrome, Is a Negative Regulator of p19 and Inhibits Senescence. Journal of Biological Chemistry, 2002, 277, 6567-6572.  | 1.6 | 140       |
| 7  | Chemokine Fractalkine/CX3CL1 Negatively Modulates Active Glutamatergic Synapses in Rat Hippocampal Neurons. Journal of Neuroscience, 2006, 26, 10488-10498.  | 1.7 | 116       |
| 8  | Adenosine A1 Receptors and Microglial Cells Mediate CX3CL1-Induced Protection of Hippocampal Neurons Against Glu-Induced Death. Neuropsychopharmacology, 2010, 35, 1550-1559.  | 2.8 | 104       |
| 9  | Fractalkine in the nervous system: neuroprotective or neurotoxic molecule?. Annals of the New York Academy of Sciences, 2015, 1351, 141-148.   | 1.8 | 98        |
| 10 | Ligand-independent CXCR2 Dimerization. Journal of Biological Chemistry, 2003, 278, 40980-40988.  | 1.6 | 97        |
| 11 | Expression of functional neurotransmitter receptors in Xenopus oocytes after injection of human brain membranes. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 13238-13242. | 3.3 | 80        |
| 12 | LTP impairment by fractalkine/CX3CL1 in mouse hippocampus is mediated through the activity of adenosine receptor type 3 (A3R). Journal of Neuroimmunology, 2009, 215, 36-42.   | 1.1 | 75        |
| 13 | Two Exon-Skipping Mutations as the Molecular Basis of Succinic Semialdehyde Dehydrogenase Deficiency (4-Hydroxybutyric Aciduria). American Journal of Human Genetics, 1998, 63, 399-408.                                 | 2.6 | 73        |
| 14 | CXCL16/CXCR6 Axis Drives Microglia/Macrophages Phenotype in Physiological Conditions and Plays a Crucial Role in Glioma. Frontiers in Immunology, 2018, 9, 2750.   | 2.2 | 71        |
| 15 | BDNF modulates GABAA receptors microtransplanted from the human epileptic brain to Xenopus oocytes. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 1667-1672.               | 3.3 | 64        |
| 16 | CXCL16 Orchestrates Adenosine A <sub>3</sub> Receptor and MCP-1/CCL2 Activity to Protect Neurons from Excitotoxic Cell Death in the CNS. Journal of Neuroscience, 2012, 32, 3154-3163.                                   | 1.7 | 60        |
| 17 | Mutant Huntingtin Forms in Vivo Complexes with Distinct Context-Dependent Conformations of the Polyglutamine Segment. Neurobiology of Disease, 1999, 6, 364-375.   | 2.1 | 57        |
| 18 | Chemokines: Key Molecules that Orchestrate Communication among Neurons, Microglia and Astrocytes to Preserve Brain Function. Neuroscience, 2020, 439, 230-240.   | 1.1 | 57        |

| #  | Article  | IF           | CITATIONS |
|----|--|--------------|-----------|
| 19 | Trasmembrane chemokines CX3CL1 and CXCL16 drive interplay between neurons, microglia and astrocytes to counteract pMCAO and excitotoxic neuronal death. Frontiers in Cellular Neuroscience, 2014, 8, 193.  | 1.8          | 52        |
| 20 | Phosphatase inhibitors remove the run-down of Â-aminobutyric acid type A receptors in the human epileptic brain. Proceedings of the National Academy of Sciences of the United States of America, 2004, 101, 10183-10188.                                | 3.3          | 50        |
| 21 | Signalling pathways involved in the chemotactic activity of CXCL12 in cultured rat cerebellar neurons and CHP100 neuroepithelioma cells. Journal of Neuroimmunology, 2003, 135, 38-46.   | 1.1          | 49        |
| 22 | Microtransplantation of membranes from cultured cells to Xenopus oocytes: A method to study neurotransmitter receptors embedded in native lipids. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 2896-2900. | 3 <b>.</b> 3 | 49        |
| 23 | Chemokine receptor CXCR2 regulates the functional properties of AMPA-type glutamate receptor GluR1 in HEK cells. Journal of Neuroimmunology, 2002, 129, 66-73.   | 1.1          | 45        |
| 24 | The Chemokine CX3CL1 Reduces Migration and Increases Adhesion of Neurons with Mechanisms Dependent on the $\hat{l}^21$ Integrin Subunit. Journal of Immunology, 2006, 177, 7599-7606.  | 0.4          | 45        |
| 25 | Rare missense variants of neuronal nicotinic acetylcholine receptor altering receptor function are associated with sporadic amyotrophic lateral sclerosis. Human Molecular Genetics, 2009, 18, 3997-4006.  | 1.4          | 42        |
| 26 | Expression of human epileptic temporal lobe neurotransmitter receptors in Xenopus oocytes: An innovative approach to study epilepsy. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 15078-15083.             | 3.3          | 40        |
| 27 | The chemokine CXCL16 modulates neurotransmitter release in hippocampal CA1 area. Scientific Reports, 2016, 6, 34633.   | 1.6          | 34        |
| 28 | Fractalkine/CX3CL1 engages different neuroprotective responses upon selective glutamate receptor overactivation. Frontiers in Cellular Neuroscience, 2014, 8, 472.   | 1.8          | 31        |
| 29 | Acetazolamide-responsive episodic ataxia in an Italian family refines gene mapping on chromosome<br>19p13. Brain, 1997, 120, 805-812.  | 3.7          | 24        |
| 30 | Human Succinic Semialdehyde Dehydrogenase. Advances in Experimental Medicine and Biology, 1996, , 253-260.   | 0.8          | 24        |
| 31 | The Glycoside Oleandrin Reduces Glioma Growth with Direct and Indirect Effects on Tumor Cells. Journal of Neuroscience, 2017, 37, 3926-3939.   | 1.7          | 23        |
| 32 | Role of Infiltrating Microglia/Macrophages in Glioma. Advances in Experimental Medicine and Biology, 2020, 1202, 281-298.  | 0.8          | 23        |
| 33 | Construction of a YAC Contig Covering Human Chromosome 6p22. Genomics, 1996, 36, 399-407.  | 1.3          | 19        |
| 34 | Expression of AMPA-type glutamate receptors in HEK cells and cerebellar granule neurons impairs CXCL2-mediated chemotaxis. Journal of Neuroimmunology, 2003, 134, 61-71.   | 1.1          | 19        |
| 35 | Adenosine A2A receptor induces protein kinase Aâ€dependent functional modulation of human α3β4 nicotinic receptor. Journal of Physiology, 2011, 589, 2755-2766.  | 1.3          | 18        |
| 36 | Basal adenosine modulates the functional properties of AMPA receptors in mouse hippocampal neurons through the activation of A1R A2AR and A3R. Frontiers in Cellular Neuroscience, 2015, 9, 409.   | 1.8          | 16        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | Co-occurring WARS2 and CHRNA6 mutations in a child with a severe form of infantile parkinsonism. Parkinsonism and Related Disorders, 2020, 72, 75-79.   | 1.1 | 16        |
| 38 | Short-chain fatty acids promote the effect of environmental signals on the gut microbiome and metabolome in mice. Communications Biology, 2022, 5, .  | 2.0 | 16        |
| 39 | A fine physical map of the CACNA1A gene region on 19p13.1–p13.2 chromosome. Gene, 2000, 241, 45-50.   | 1.0 | 15        |
| 40 | Cysteine residues are critical for chemokine receptor CXCR2 functional properties. Experimental Cell Research, 2005, 307, 65-75.  | 1.2 | 15        |
| 41 | Neuro-Signals from Gut Microbiota: Perspectives for Brain Glioma. Cancers, 2021, 13, 2810.  | 1.7 | 14        |
| 42 | Human succinic semialdehyde dehydrogenase. Molecular cloning and chromosomal localization. Advances in Experimental Medicine and Biology, 1997, 414, 253-60.                                    | 0.8 | 14        |
| 43 | Chemokine CXCL8 modulates GluR1 phosphorylation. Journal of Neuroimmunology, 2008, 198, 75-81.  | 1.1 | 10        |
| 44 | Mutant human $\hat{1}^24$ subunit identified in amyotrophic lateral sclerosis patients impairs nicotinic receptor function. Pflugers Archiv European Journal of Physiology, 2011, 461, 225-233. | 1.3 | 8         |
| 45 | Localization and genomic structure of human deoxyhypusine synthase gene on chromosome 19p13.2-distal 19p13.1. Gene, 1998, 215, 153-157.   | 1.0 | 7         |
| 46 | Editorial Research Topic "Chemokines and chemokine receptors in brain homeostasis― Frontiers in Cellular Neuroscience, 2015, 9, 132.  | 1.8 | 7         |
| 47 | Chemokines and chemokine receptors in the nervous system. Journal of Neuroimmunology, 2008, 198, 1-8.   | 1.1 | 4         |
| 48 | Ordering of 44 Genetic Markers in the 6p22 Cytogenetic Band. DNA Sequence, 1996, 7, 51-52.  | 0.7 | 0         |