

Llorenç Caballeria

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6219024/publications.pdf>

Version: 2024-02-01

60
papers

4,827
citations

172207

29
h-index

110170

64
g-index

71
all docs

71
docs citations

71
times ranked

4046
citing authors

#	ARTICLE	IF	CITATIONS
1	Excellent Long-Term Survival in Patients With Primary Biliary Cirrhosis and Biochemical Response to Ursodeoxycholic Acid. <i>Gastroenterology</i> , 2006, 130, 715-720.	0.6	649
2	S-Adenosylmethionine in alcoholic liver cirrhosis: a randomized, placebo-controlled, double-blind, multicenter clinical trial. <i>Journal of Hepatology</i> , 1999, 30, 1081-1089.	1.8	428
3	Levels of Alkaline Phosphatase and Bilirubin Are Surrogate End Points of Outcomes of Patients With Primary Biliary Cirrhosis: An International Follow-up Study. <i>Gastroenterology</i> , 2014, 147, 1338-1349.e5.	0.6	365
4	Development and Validation of a Scoring System to Predict Outcomes of Patients With Primary Biliary Cirrhosis Receiving Ursodeoxycholic Acid Therapy. <i>Gastroenterology</i> , 2015, 149, 1804-1812.e4.	0.6	330
5	Long-term effects of ursodeoxycholic acid in primary biliary cirrhosis: results of a double-blind controlled multicentric trial. <i>Journal of Hepatology</i> , 2000, 32, 561-566.	1.8	254
6	Hepatitis C virus antibodies in chronic alcoholic patients: Association with severity of liver injury. <i>Hepatology</i> , 1990, 12, 1295-1299.	3.6	242
7	Prevalence and factors associated with the presence of nonalcoholic fatty liver disease in an adult population in Spain. <i>European Journal of Gastroenterology and Hepatology</i> , 2010, 22, 24-32.	0.8	196
8	High Prevalence of Liver Fibrosis Among European Adults With Unknown Liver Disease: A Population-Based Study. <i>Clinical Gastroenterology and Hepatology</i> , 2018, 16, 1138-1145.e5.	2.4	180
9	Severity of cholestasis and advanced histological stage but not menopausal status are the major risk factors for osteoporosis in primary biliary cirrhosis. <i>Journal of Hepatology</i> , 2005, 42, 573-577.	1.8	163
10	HLA class II haplotypes in primary sclerosing cholangitis patients from five European populations. <i>Tissue Antigens</i> , 1999, 53, 459-469.	1.0	151
11	Stratification of hepatocellular carcinoma risk in primary biliary cirrhosis: a multicentre international study. <i>Gut</i> , 2016, 65, 321-329.	6.1	139
12	Screening for liver fibrosis in the general population: a call for action. <i>The Lancet Gastroenterology and Hepatology</i> , 2016, 1, 256-260.	3.7	131
13	Low Bone Mass and Severity of Cholestasis Affect Fracture Risk in Patients With Primary Biliary Cirrhosis. <i>Gastroenterology</i> , 2010, 138, 2348-2356.	0.6	115
14	Extracorporeal Albumin Dialysis: A Procedure for Prolonged Relief of Intractable Pruritus in Patients with Primary Biliary Cirrhosis. <i>American Journal of Gastroenterology</i> , 2004, 99, 1105-1110.	0.2	111
15	Transient elastography for screening of liver fibrosis: Cost-effectiveness analysis from six prospective cohorts in Europe and Asia. <i>Journal of Hepatology</i> , 2019, 71, 1141-1151.	1.8	104
16	Hepatocellular Carcinoma in Primary Biliary Cirrhosis: Similar Incidence To That in Hepatitis C Virus-Related Cirrhosis. <i>American Journal of Gastroenterology</i> , 2001, 96, 1160-1163.	0.2	96
17	Incidence, risk factors, and survival of hepatocellular carcinoma in primary biliary cirrhosis: Comparative analysis from two centers. <i>Hepatology</i> , 2009, 50, 1162-1168.	3.6	93
18	Low Accuracy of FIB-4 and NAFLD Fibrosis Scores for Screening for Liver Fibrosis in the Population. <i>Clinical Gastroenterology and Hepatology</i> , 2022, 20, 2567-2576.e6.	2.4	80

#	ARTICLE	IF	CITATIONS
19	Time-dependent Cox regression model is superior in prediction of prognosis in primary sclerosing cholangitis. <i>Hepatology</i> , 2002, 35, 652-657.	3.6	77
20	New ELISA for Detecting Primary Biliary Cirrhosis's Specific Antimitochondrial Antibodies. <i>Clinical Chemistry</i> , 2009, 55, 978-985.	1.5	71
21	Documento de consenso. Manejo de la enfermedad hepática grasa no alcohólica (EHGNA). Guía de práctica clínica. <i>Gastroenterología Y Hepatología</i> , 2018, 41, 328-349.	0.2	71
22	Metadoxine accelerates fatty liver recovery in alcoholic patients: results of a randomized double-blind, placebo-control trial. <i>Journal of Hepatology</i> , 1998, 28, 54-60.	1.8	68
23	Collagen type I±1 and vitamin D receptor gene polymorphisms and bone mass in primary biliary cirrhosis. <i>Hepatology</i> , 2001, 33, 554-560.	3.6	64
24	Disease-specific cross-reactivity between mimicking peptides of heat shock protein of mycobacterium gordonae and dominant epitope of E2 subunit of pyruvate dehydrogenase is common in Spanish but not British patients with primary biliary cirrhosis. <i>Journal of Autoimmunity</i> , 2004, 22, 353-362.	3.0	64
25	Anti-gp210 antibody mirrors disease severity in primary biliary cirrhosis. <i>Hepatology</i> , 2007, 45, 1583-1583.	3.6	51
26	Fatty liver index is a predictor of incident diabetes in patients with prediabetes: The PREDAPS study. <i>PLoS ONE</i> , 2018, 13, e0198327.	1.1	38
27	Prevalence and factors associated with the presence of non alcoholic fatty liver disease in an apparently healthy adult population in primary care units. <i>BMC Gastroenterology</i> , 2007, 7, 41.	0.8	35
28	Thyroid Dysfunction in Primary Biliary Cholangitis: A Comparative Study at Two European Centers. <i>American Journal of Gastroenterology</i> , 2017, 112, 114-119.	0.2	34
29	Alerts in electronic medical records to promote a colorectal cancer screening programme: a cluster randomised controlled trial in primary care. <i>British Journal of General Practice</i> , 2016, 66, e483-e490.	0.7	30
30	Serum hyaluronate reflects hepatic fibrogenesis in alcoholic liver disease and is useful as a marker of fibrosis. <i>Hepatology</i> , 1996, 24, 1399-1403.	3.6	27
31	Hepatocellular carcinoma in primary biliary cirrhosis: similar incidence to that in hepatitis C virus-related cirrhosis. <i>American Journal of Gastroenterology</i> , 2001, 96, 1160-1163.	0.2	26
32	Alendronate is more effective than etidronate for increasing bone mass in osteopenic patients with primary biliary cirrhosis. <i>American Journal of Gastroenterology</i> , 2003, 98, 2268-2274.	0.2	24
33	Metabolic syndrome and nonalcoholic fatty liver disease in a Spanish population. <i>European Journal of Gastroenterology and Hepatology</i> , 2012, 24, 1007-1011.	0.8	24
34	Carbohydrate-Deficient Transferrin as a Marker of Alcohol Consumption in Male Patients with Liver Disease. <i>Alcoholism: Clinical and Experimental Research</i> , 1997, 21, 923-927.	1.4	23
35	Using Transient Elastography to Detect Chronic Liver Diseases in a Primary Care Nurse Consultancy. <i>Nursing Research</i> , 2013, 62, 450-454.	0.8	21
36	High osteoprotegerin serum levels in primary biliary cirrhosis are associated with disease severity but not with the mRNA gene expression in liver tissue. <i>Journal of Bone and Mineral Metabolism</i> , 2009, 27, 347-354.	1.3	19

#	ARTICLE	IF	CITATIONS
37	Extrahepatic Malignancies in Primary Biliary Cirrhosis: A Comparative Study at Two European Centers. <i>Clinical Reviews in Allergy and Immunology</i> , 2015, 48, 254-262.	2.9	19
38	Recomendaciones para la detección, diagnóstico y seguimiento de los pacientes con enfermedad por hígado graso no alcohólico en atención primaria y hospitalaria. <i>Medicina Clínica</i> , 2019, 153, 169-177.	0.3	18
39	Risk factors associated with non-alcoholic fatty liver disease in subjects from primary care units. A case-control study. <i>BMC Gastroenterology</i> , 2008, 8, 44.	0.8	17
40	Relación entre el hipotiroidismo y el hígado graso no alcohólico en una población española. <i>Medicina Clínica</i> , 2020, 154, 1-6.	0.3	13
41	Marcadores de fibrosis hepática. <i>Medicina Clínica</i> , 2018, 150, 310-316.	0.3	12
42	Atherogenic dyslipidemia, but not hyperglycemia, is an independent factor associated with liver fibrosis in subjects with type 2 diabetes and NAFLD: a population-based study. <i>European Journal of Endocrinology</i> , 2021, 184, 587-596.	1.9	12
43	TSH Levels as an Independent Risk Factor for NAFLD and Liver Fibrosis in the General Population. <i>Journal of Clinical Medicine</i> , 2021, 10, 2907.	1.0	12
44	Zinc Administration Improves Gastric Alcohol Dehydrogenase Activity and First-Pass Metabolism of Ethanol in Alcohol-Fed Rats. <i>Alcoholism: Clinical and Experimental Research</i> , 1997, 21, 1619-1622.	1.4	9
45	Relationship between hypothyroidism and non-alcoholic fatty liver disease in the Spanish population. <i>Medicina Clínica (English Edition)</i> , 2020, 154, 1-6.	0.1	9
46	Predicting and preventing autoimmunity: the case of anti-mitochondrial antibodies. <i>Autoimmunity Highlights</i> , 2012, 3, 105-112.	3.9	7
47	Consensus document. Management of non-alcoholic fatty liver disease (NAFLD). Clinical practice guideline. <i>Gastroenterología Y Hepatología (English Edition)</i> , 2018, 41, 328-349.	0.0	7
48	Recommendations for the detection, diagnosis and follow-up of patients with non-alcoholic fatty liver disease in primary and hospital care. <i>Medicina Clínica (English Edition)</i> , 2019, 153, 169-177.	0.1	7
49	Estrategias para la detección de infección por virus de la hepatitis C en población general. <i>Revista Clínica Española</i> , 2014, 214, 242-246.	0.2	6
50	Adequacy and quality of abdominal echographies requested by primary care professionals. <i>BMC Gastroenterology</i> , 2010, 10, 101.	0.8	5
51	Nonalcoholic steatohepatitis and diabetes. <i>Endocrinología Y Nutrición (English Edition)</i> , 2016, 63, 377-379.	0.5	5
52	Esteatohepatitis no alcohólica y diabetes. <i>Endocrinología Y Nutrición: Órgano De La Sociedad Española De Endocrinología Y Nutrición</i> , 2016, 63, 377-379.	0.8	5
53	Markers of hepatic fibrosis. <i>Medicina Clínica (English Edition)</i> , 2018, 150, 310-316.	0.1	5
54	Strategies for the detection of hepatitis C viral infection in the general population. <i>Revista Clínica Española</i> , 2014, 214, 242-246.	0.3	4

#	ARTICLE	IF	CITATIONS
55	Risk of Liver Fibrosis According to TSH Levels in Euthyroid Subjects. Journal of Clinical Medicine, 2021, 10, 1350.	1.0	4
56	Prevalence of Early Chronic Kidney Disease and Main Associated Factors in Spanish Population: Populational Study. Journal of Clinical Medicine, 2019, 8, 1384.	1.0	3
57	Reply:. Hepatology, 2010, 52, 2239-2239.	3.6	1
58	Acceptability and quality of abdominal ultrasound studies requested by medical professionals. Gastroenterología Y Hepatología (English Edition), 2016, 39, 516-525.	0.0	1
59	Non-alcoholic fatty liver disease. Role of primary care physicians. Medicina Clínica (English Edition), 2015, 145, 112-114.	0.1	0
60	Screening studies of transient elastography and FibroTest in the general population – Authors' reply. The Lancet Gastroenterology and Hepatology, 2017, 2, 246-247.	3.7	0