

Josune Orbe

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/622476/josune-orbe-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

93
papers

2,454
citations

31
h-index

46
g-index

109
ext. papers

2,855
ext. citations

5.5
avg, IF

4.37
L-index

#	Paper	IF	Citations
93	Role of Extracellular Vesicles as Potential Diagnostic and/or Therapeutic Biomarkers in Chronic Cardiovascular Diseases.. <i>Frontiers in Cell and Developmental Biology</i> , 2022 , 10, 813885	5.7	3
92	Hemostatic Biomarkers and Volumetry Help to Identify High-Risk Abdominal Aortic Aneurysms. <i>Life</i> , 2022 , 12, 823	3	
91	The Role of Circulating Biomarkers in Peripheral Arterial Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	7
90	Urinary Extracellular Vesicles for Diabetic Kidney Disease Diagnosis. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	6
89	Association of SDF1 and MMP12 with Atherosclerosis and Inflammation: Clinical and Experimental Study. <i>Life</i> , 2021 , 11,	3	1
88	Molecular and Cellular Mechanisms of Delayed Fracture Healing in Mmp10 (Stromelysin 2) Knockout Mice. <i>Journal of Bone and Mineral Research</i> , 2021 , 36, 2203-2213	6.3	0
87	Inside the Thrombus: Association of Hemostatic Parameters With Outcomes in Large Vessel Stroke Patients. <i>Frontiers in Neurology</i> , 2021 , 12, 599498	4.1	1
86	Association of calprotectin with other inflammatory parameters in the prediction of mortality for ischemic stroke. <i>Journal of Neuroinflammation</i> , 2021 , 18, 3	10.1	5
85	A Role For MMP-10 (Matrix Metalloproteinase-10) in Calcific Aortic Valve Stenosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2020 , 40, 1370-1382	9.4	15
84	Circulating TIMP-1 is associated with hematoma volume in patients with spontaneous intracranial hemorrhage. <i>Scientific Reports</i> , 2020 , 10, 10329	4.9	1
83	NADPH Oxidase Overactivity Underlies Telomere Shortening in Human Atherosclerosis. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	7
82	Functional and transcriptomic analysis of extracellular vesicles identifies calprotectin as a new prognostic marker in peripheral arterial disease (PAD). <i>Journal of Extracellular Vesicles</i> , 2020 , 9, 1729646	16.4	17
81	Matrix Metalloproteinases in Diabetic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	35
80	MMP-10 is Increased in Early Stage Diabetic Kidney Disease and can be Reduced by Renin-Angiotensin System Blockade. <i>Scientific Reports</i> , 2020 , 10, 26	4.9	8
79	Persistently high circulating tissue inhibitor of matrix metalloproteinase-1 levels in non-survivor brain trauma injury patients. <i>Journal of Critical Care</i> , 2019 , 51, 117-121	4	3
78	MMP10 Promotes Efficient Thrombolysis After Ischemic Stroke in Mice with Induced Diabetes. <i>Translational Stroke Research</i> , 2019 , 10, 389-401	7.8	11
77	High serum levels of tissue inhibitor of matrix metalloproteinase-1 during the first week of a malignant middle cerebral artery infarction in non-surviving patients. <i>BMC Neurology</i> , 2019 , 19, 167	3.1	3

76	Trimethylamine-N-Oxide (TMAO) Predicts Cardiovascular Mortality in Peripheral Artery Disease. <i>Scientific Reports</i> , 2019 , 9, 15580	4.9	49
75	Combined sustained release of BMP2 and MMP10 accelerates bone formation and mineralization of calvaria critical size defect in mice. <i>Drug Delivery</i> , 2018 , 25, 750-756	7	15
74	Phenotypic Screening To Discover Novel Chemical Series as Efficient Antihemorrhagic Agents. <i>ACS Medicinal Chemistry Letters</i> , 2018 , 9, 428-433	4.3	2
73	Selective increase of cardiomyocyte derived extracellular vesicles after experimental myocardial infarction and functional effects on the endothelium. <i>Thrombosis Research</i> , 2018 , 170, 1-9	8.2	9
72	Matrix metalloproteinase-10 deficiency delays atherosclerosis progression and plaque calcification. <i>Atherosclerosis</i> , 2018 , 278, 124-134	3.1	20
71	Recent progress in translational research on neurovascular and neurodegenerative disorders. <i>Restorative Neurology and Neuroscience</i> , 2017 , 35, 87-103	2.8	14
70	CM352 Reduces Brain Damage and Improves Functional Recovery in a Rat Model of Intracerebral Hemorrhage. <i>Journal of the American Heart Association</i> , 2017 , 6,	6	14
69	New thrombolytic strategy providing neuroprotection in experimental ischemic stroke: MMP10 alone or in combination with tissue-type plasminogen activator. <i>Cardiovascular Research</i> , 2017 , 113, 1219-1229 ¹⁰	8.9	29
68	Reduced high-density lipoprotein cholesterol: A valuable, independent prognostic marker in peripheral arterial disease. <i>Journal of Vascular Surgery</i> , 2017 , 66, 1527-1533.e1	3.5	11
67	Efficacy of Alteplase in a Mouse Model of Acute Ischemic Stroke: A Retrospective Pooled Analysis. <i>Stroke</i> , 2016 , 47, 1312-1318	6.7	25
66	High triglyceride-low HDL cholesterol lipid profile is associated with a dysregulated gene expression in mononuclear leukocyte from hypercholesterolemic patients. <i>International Journal of Cardiology</i> , 2015 , 178, 102-4	3.2	3
65	Lack of TAFI increases brain damage and microparticle generation after thrombolytic therapy in ischemic stroke. <i>Thrombosis Research</i> , 2015 , 136, 445-50	8.2	12
64	Serum tissue inhibitor of matrix metalloproteinase-1 levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>BMC Neurology</i> , 2015 , 15, 111	3.1	10
63	Functional MMP-10 is required for efficient tissue repair after experimental hind limb ischemia. <i>FASEB Journal</i> , 2015 , 29, 960-72	0.9	14
62	Activated protein C glycoform promotes enhanced noncanonical PAR1 proteolysis and superior resistance to ischemic injury. <i>Blood</i> , 2015 , 126, 915-9	2.2	17
61	Matrix metalloproteinase 10 contributes to hepatocarcinogenesis in a novel crosstalk with the stromal derived factor 1/C-X-C chemokine receptor 4 axis. <i>Hepatology</i> , 2015 , 62, 166-78	11.2	44
60	Targeting βsecretases protect against angiotensin II-induced cardiac hypertrophy. <i>Journal of Hypertension</i> , 2015 , 33, 843-50; discussion 850	1.9	5
59	Matrix metalloproteinase 10 is associated with disease severity and mortality in patients with peripheral arterial disease. <i>Journal of Vascular Surgery</i> , 2015 , 61, 428-35	3.5	31

58	Discovery and safety profiling of a potent preclinical candidate, (4-[4-[[[(3R)-3-(hydroxycarbamoyl)-8-azaspiro[4.5]decan-3-yl]sulfonyl]phenoxy]-N-methylbenzamide) (CM-352), for the prevention and treatment of hemorrhage. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 2941-57	8.3	10
57	Design, synthesis, and biological evaluation of novel matrix metalloproteinase inhibitors as potent antihemorrhagic agents: from hit identification to an optimized lead. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 2465-88	8.3	17
56	The CXCR4/SDF1 axis improves muscle regeneration through MMP-10 activity. <i>Stem Cells and Development</i> , 2014 , 23, 1417-27	4.4	24
55	MMP-10 is required for efficient muscle regeneration in mouse models of injury and muscular dystrophy. <i>Stem Cells</i> , 2014 , 32, 447-61	5.8	23
54	Matrix metalloproteinase-10 expression is induced during hepatic injury and plays a fundamental role in liver tissue repair. <i>Liver International</i> , 2014 , 34, e257-70	7.9	37
53	Altered atherosclerotic-related gene expression signature in circulating mononuclear leukocytes from hypercholesterolemic patients with low HDL cholesterol levels. <i>International Journal of Cardiology</i> , 2014 , 173, 337-8	3.2	4
52	Association of sepsis-related mortality with early increase of TIMP-1/MMP-9 ratio. <i>PLoS ONE</i> , 2014 , 9, e94318	3.7	43
51	Association between serum tissue inhibitor of matrix metalloproteinase-1 levels and mortality in patients with severe brain trauma injury. <i>PLoS ONE</i> , 2014 , 9, e94370	3.7	27
50	"tissue factor expressed by microparticles is associated with mortality but not with thrombosis in cancer patients." reply to a comment by Geddings and Mackman. <i>Thrombosis and Haemostasis</i> , 2014 , 111, 182	7	1
49	Signature of subclinical femoral artery atherosclerosis in peripheral blood mononuclear cells. <i>European Journal of Clinical Investigation</i> , 2014 , 44, 539-48	4.6	5
48	The 372 T/C genetic polymorphism of TIMP-1 is associated with serum levels of TIMP-1 and survival in patients with severe sepsis. <i>Critical Care</i> , 2013 , 17, R94	10.8	22
47	proMetalloproteinase-10 is associated with brain damage and clinical outcome in acute ischemic stroke. <i>Journal of Thrombosis and Haemostasis</i> , 2013 , 11, 1464-73	15.4	38
46	Tissue factor expressed by microparticles is associated with mortality but not with thrombosis in cancer patients. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 598-608	7	45
45	Synergistic effect of thrombin and CD40 ligand on endothelial matrix metalloproteinase-10 expression and microparticle generation in vitro and in vivo. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2012 , 32, 1477-87	9.4	46
44	Adipose stromal vascular fraction improves cardiac function in chronic myocardial infarction through differentiation and paracrine activity. <i>Cell Transplantation</i> , 2012 , 21, 1023-37	4	38
43	Association between serum soluble CD40 ligand levels and mortality in patients with severe sepsis. <i>Critical Care</i> , 2011 , 15, R97	10.8	43
42	Matrix metalloproteinase-10 effectively reduces infarct size in experimental stroke by enhancing fibrinolysis via a thrombin-activatable fibrinolysis inhibitor-mediated mechanism. <i>Circulation</i> , 2011 , 124, 2909-19	16.7	45
41	CCL20 is increased in hypercholesterolemic subjects and is upregulated by LDL in vascular smooth muscle cells: role of NF- κ B. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 2733-41	9.4	34

40	Serum levels of matrix metalloproteinase-10 are associated with the severity of atherosclerosis in patients with chronic kidney disease. <i>Kidney International</i> , 2010 , 78, 1275-80	9.9	31
39	Corrigendum to Preliminary characterisation of the promoter of the human p22phox gene: Identification of a new polymorphism associated with hypertension [FEBS Lett. 542 (2003) 27B1]. <i>FEBS Letters</i> , 2010 , 584, 4709-4709	3.8	
38	Tissue-specific PAI-1 gene expression and glycosylation pattern in insulin-resistant old rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2009 , 297, R1563-9	3.2	23
37	Matrix metalloproteinase-10 is upregulated by thrombin in endothelial cells and increased in patients with enhanced thrombin generation. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 2109-16	9.4	36
36	Matrix metalloproteinase-9, -10, and tissue inhibitor of matrix metalloproteinases-1 blood levels as biomarkers of severity and mortality in sepsis. <i>Critical Care</i> , 2009 , 13, R158	10.8	86
35	The CD163-expressing macrophages recognize and internalize TWEAK: potential consequences in atherosclerosis. <i>Atherosclerosis</i> , 2009 , 207, 103-10	3.1	108
34	[Not Available]. <i>Revista Espanola De Cardiologia</i> , 2008 , 61, 327-8	1.5	4
33	Association between matrix metalloproteinase-10 concentration and smoking in individuals without cardiovascular disease. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2008 , 61, 1267-73	0.7	2
32	Increased thrombin generation after acute versus chronic coronary disease as assessed by the thrombin generation test. <i>Thrombosis and Haemostasis</i> , 2008 , 99, 382-7	7	45
31	Asociaci3n de la metaloproteinasas-10 y el tabaquismo en sujetos sin enfermedad cardiovascular. <i>Revista Espanola De Cardiologia</i> , 2008 , 61, 1267-1273	1.5	6
30	Metalloproteinases and atherothrombosis: MMP-10 mediates vascular remodeling promoted by inflammatory stimuli. <i>Frontiers in Bioscience - Landmark</i> , 2008 , 13, 2916-21	2.8	70
29	Independent association of matrix metalloproteinase-10, cardiovascular risk factors and subclinical atherosclerosis. <i>Journal of Thrombosis and Haemostasis</i> , 2007 , 5, 91-7	15.4	51
28	Folic acid and B vitamins improve hyperhomocysteinemia-induced cardiovascular risk profile in renal transplant recipients. <i>Journal of Thrombosis and Haemostasis</i> , 2007 , 5, 1072-6	15.4	3
27	Phagocytic NADPH oxidase-dependent superoxide production stimulates matrix metalloproteinase-9: implications for human atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 587-93	9.4	71
26	Metaloproteasa-10 (estromelisina-2): un nuevo marcador de aterosclerosis subcl3nica. <i>Cl3nica E Investigaci3n En Arteriosclerosis</i> , 2007 , 19, 122-128	1.4	
25	Identification of soluble tumor necrosis factor-like weak inducer of apoptosis (sTWEAK) as a possible biomarker of subclinical atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 916-22	9.4	107
24	Integrating soluble biomarkers and imaging technologies in the identification of vulnerable atherosclerotic patients. <i>Biomarker Insights</i> , 2007 , 1, 165-73	3.5	3
23	A comparison between percutaneous and surgical transplantation of autologous skeletal myoblasts in a swine model of chronic myocardial infarction. <i>Cardiovascular Research</i> , 2006 , 71, 744-53	9.9	47

22	C-reactive protein induces matrix metalloproteinase-1 and -10 in human endothelial cells: implications for clinical and subclinical atherosclerosis. <i>Journal of the American College of Cardiology</i> , 2006 , 47, 1369-78	15.1	147
21	Protective effect of the G-765C COX-2 polymorphism on subclinical atherosclerosis and inflammatory markers in asymptomatic subjects with cardiovascular risk factors. <i>Clinica Chimica Acta</i> , 2006 , 368, 138-43	6.2	48
20	Integrating Soluble Biomarkers and Imaging Technologies in the Identification of Vulnerable Atherosclerotic Patients. <i>Biomarker Insights</i> , 2006 , 1, 117727190600100	3.5	1
19	The 4G/5G PAI-1 polymorphism influences the endothelial response to IL-1 and the modulatory effect of pravastatin. <i>Journal of Thrombosis and Haemostasis</i> , 2006 , 4, 1798-803	15.4	13
18	Independent association of von Willebrand factor with surrogate markers of atherosclerosis in middle-aged asymptomatic subjects. <i>Journal of Thrombosis and Haemostasis</i> , 2005 , 3, 662-4	15.4	31
17	Monocyte cyclooxygenase-2 overactivity: a new marker of subclinical atherosclerosis in asymptomatic subjects with cardiovascular risk factors?. <i>European Heart Journal</i> , 2005 , 26, 153-8	9.5	34
16	Prothrombin fragment 1+2 is associated with carotid intima-media thickness in subjects free of clinical cardiovascular disease. <i>Stroke</i> , 2004 , 35, 1085-9	6.7	43
15	Influence of the 4G/5G PAI-1 genotype on angiotensin II-stimulated human endothelial cells and in patients with hypertension. <i>Cardiovascular Research</i> , 2004 , 63, 176-85	9.9	15
14	Validation of plasma fibrinogen as a marker of carotid atherosclerosis in subjects free of clinical cardiovascular disease. <i>Haematologica</i> , 2004 , 89, 1226-31	6.6	25
13	The A1166C polymorphism of the AT1 receptor gene is associated with collagen type I synthesis and myocardial stiffness in hypertensives. <i>Journal of Hypertension</i> , 2003 , 21, 2085-2092	1.9	31
12	Antioxidant vitamins increase the collagen content and reduce MMP-1 in a porcine model of atherosclerosis: implications for plaque stabilization. <i>Atherosclerosis</i> , 2003 , 167, 45-53	3.1	48
11	Different expression of MMPs/TIMP-1 in human atherosclerotic lesions. Relation to plaque features and vascular bed. <i>Atherosclerosis</i> , 2003 , 170, 269-76	3.1	85
10	Preliminary characterisation of the promoter of the human p22(phox) gene: identification of a new polymorphism associated with hypertension. <i>FEBS Letters</i> , 2003 , 542, 27-31	3.8	73
9	Independent association of fibrinogen with carotid intima-media thickness in asymptomatic subjects. <i>Cerebrovascular Diseases</i> , 2003 , 16, 356-62	3.2	24
8	The A1166C polymorphism of the AT1 receptor gene is associated with collagen type I synthesis and myocardial stiffness in hypertensives. <i>Journal of Hypertension</i> , 2003 , 21, 2085-92	1.9	10
7	Regulation by Nitric Oxide of Endotoxin-Induced Tissue Factor and Plasminogen Activator Inhibitor-1 in Endothelial Cells. <i>Thrombosis and Haemostasis</i> , 2002 , 88, 1060-1065	7	24
6	Involvement of leptin in the association between percentage of body fat and cardiovascular risk factors. <i>Clinical Biochemistry</i> , 2002 , 35, 315-20	3.5	79
5	Hemostasis, inflammation and cardiovascular disease. <i>Clinical Laboratory</i> , 2002 , 48, 463-70	2	3

4	Vitamins C and E attenuate plasminogen activator inhibitor-1 (PAI-1) expression in a hypercholesterolemic porcine model of angioplasty. <i>Cardiovascular Research</i> , 2001 , 49, 484-92	9.9	12
3	Effect of the administration of recombinant hirudin and/or tissue-plasminogen activator (t-PA) on endotoxin-induced disseminated intravascular coagulation model in rabbits. <i>British Journal of Haematology</i> , 1999 , 105, 117-121	4.5	30
2	Markers of fibrinolytic potency and clotting activation in stable angina pectoris: role of urokinase, assessment of atrioventricular differences and correlation with coronary patency. <i>Fibrinolysis and Proteolysis</i> , 1999 , 13, 133-138		
1	Evidence that heparin but not hirudin reduces PAI-1 expression in cultured human endothelial cells. <i>Thrombosis Research</i> , 1999 , 94, 137-45	8.2	8