
## **Zbigniew Siudak**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6239772/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Impact of Multivessel Coronary Artery Disease and Noninfarct-Related Artery Revascularization on Outcome of Patients With ST-Elevation Myocardial Infarction Transferred for Primary Percutaneous Coronary Intervention (from the EUROTRANSFER Registry). American Journal of Cardiology, 2010, 106, 342-347.	0.7	109
2	A Prospective, Multicenter Study of aÂNovel Mesh-Covered Carotid Stent. JACC: Cardiovascular Interventions, 2015, 8, 1229-1234.	1,1	108
3	Plasma Homocysteine Affects Fibrin Clot Permeability and Resistance to Lysis in Human Subjects. Arteriosclerosis, Thrombosis, and Vascular Biology, 2006, 26, 1397-1404.	1.1	107
4	Thrombus aspiration followed by direct stenting: A novel strategy of primary percutaneous coronary intervention in ST-segment elevation myocardial infarction. Results of the Polish-Italian-Hungarian RAndomized ThrombEctomy Trial (PIHRATE Trial). American Heart Journal, 2010, 160, 966-972.	1.2	83
5	European registry on patients with ST-elevation myocardial infarction transferred for mechanical reperfusion with a special focus on early administration of abciximab—EUROTRANSFER Registry. American Heart Journal, 2008, 156, 1147-1154.	1.2	60
6	Altered Plasma Fibrin Clot Properties Are Associated With In-Stent Thrombosis. Arteriosclerosis, Thrombosis, and Vascular Biology, 2010, 30, 276-282.	1.1	55
7	Early abciximab administration before transfer for primary percutaneous coronary interventions for ST-elevation myocardial infarction reduces 1-year mortality in patients with high-risk profile. Results from EUROTRANSFER Registry. American Heart Journal, 2009, 158, 569-575.	1.2	35
8	Impact of smoking status on outcome in patients with ST-segment elevation myocardial infarction treated with primary percutaneous coronary intervention. Journal of Thrombosis and Thrombolysis, 2012, 34, 397-403.	1.0	33
9	Mesh covered stent in ST-segment elevation myocardial infarction. EuroIntervention, 2010, 6, 582-589.	1.4	33
10	Transradial approach in patients with ST-elevation myocardial infarction treated with abciximab results in fewer bleeding complications: data from EUROTRANSFER registry. Coronary Artery Disease, 2010, 21, 292-297.	0.3	31
11	Admission glucose level and in-hospital outcomes in diabetic and non-diabetic patients with acute myocardial infarction. Clinical Research in Cardiology, 2010, 99, 715-721.	1.5	27
12	Impact of direct stenting on outcome of patients with STâ€elevation myocardial infarction transferred for primary percutaneous coronary intervention (from the EUROTRANSFER registry). Catheterization and Cardiovascular Interventions, 2014, 84, 925-931.	0.7	27
13	Characteristics of patients presenting with myocardial infarction with non-obstructive coronary arteries (MINOCA) in Poland: data from the ORPKI national registry. Journal of Thrombosis and Thrombolysis, 2019, 47, 462-466.	1.0	27
14	Age-related differences in treatment strategies and clinical outcomes in unselected cohort of patients with ST-segment elevation myocardial infarction transferred for primary angioplasty. Journal of Thrombosis and Thrombolysis, 2012, 34, 214-221.	1.0	26
15	Morphologic variability of the mitral valve leaflets. Journal of Thoracic and Cardiovascular Surgery, 2017, 154, 1927-1935.	0.4	26
16	Clinical and procedural characteristics of <scp>COVID</scp> â€19 patients treated with percutaneous coronary interventions. Catheterization and Cardiovascular Interventions, 2020, 96, E568-E575.	0.7	26
17	Out-of-hospital cardiac arrest in patients treated with primary PCI for STEMI. Long-term follow up data from EUROTRANSFER registry. Resuscitation, 2012, 83, 303-306.	1.3	24
18	Assessment of quality of life in patients after surgical and transcatheter aortic valve replacement. Catheterization and Cardiovascular Interventions, 2016, 88, E80-8.	0.7	23

#	Article	IF	CITATIONS
19	Anatomy of the mitral subvalvular apparatus. Journal of Thoracic and Cardiovascular Surgery, 2018, 155, 2002-2010.	0.4	23
20	Predictors of inâ€hospital effectiveness and complications of rotational atherectomy (from the ORPKI) Tj ETQqC E278-E287.	0 0 rgBT 0.7	Overlock 107 23
21	Bioresorbable vascular scaffolds in patients with acute coronary syndromes : the POLAR ACS study. Polish Archives of Internal Medicine, 2014, 124, 669-677.	0.3	22
22	Reduced periprocedural mortality and bleeding rates of radial approach in ST-segment elevation myocardial infarction. Propensity score analysis of data from the ORPKI Polish National Registry. EuroIntervention, 2017, 13, 843-850.	1.4	22
23	Radial Approach Expertise and Clinical Outcomes of Percutanous Coronary Interventions Performed Using Femoral Approach. Journal of Clinical Medicine, 2019, 8, 1484.	1.0	20
24	Percutaneous coronary intervention during on- and off-hours in patients with ST-segment elevation myocardial infarction. Hellenic Journal of Cardiology, 2021, 62, 212-218.	0.4	20
25	Use of bioresorbable vascular scaffolds in patients with stable angina and acute coronary syndromes. Polish National Registry. Kardiologia Polska, 2014, 72, 1394-1399.	0.3	18
26	Prevalence and Predictors of Coronary Artery Perforation During Percutaneous Coronary Interventions (from the ORPKI National Registry in Poland). American Journal of Cardiology, 2019, 124, 1186-1189.	0.7	17
27	Determinants of stroke following percutaneous coronary intervention in acute myocardial infarction (from ORPKI Polish National Registry). International Journal of Cardiology, 2016, 223, 236-238.	0.8	16
28	Prognostic significance of new onset atrial fibrillation in acute coronary syndrome patients treated conservatively. Cardiology Journal, 2010, 17, 57-64.	0.5	16
29	Association between the mortality rate and operator volume in patients undergoing emergency or elective percutaneous coronary interventions. Kardiologia Polska, 2020, 78, 138-146.	0.3	15
30	Impact of Admission Glucose Level and Presence of Diabetes Mellitus on Mortality in Patients With Non–ST-Segment Elevation Acute Coronary Syndrome Treated Conservatively. American Journal of Cardiology, 2009, 103, 954-958.	0.7	14
31	Simvastatin administration reduces thromboxane production in subjects taking aspirin: Links between aspirin resistance and thrombin generation. International Journal of Cardiology, 2012, 154, 59-64.	0.8	13
32	Borderline trend towards longâ€ŧerm mortality benefit from drug eluting stents implantation in STâ€elevation myocardial infarction patients in Poland—data from NRDES registry. Catheterization and Cardiovascular Interventions, 2014, 83, 436-442.	0.7	13
33	Contemporary use of P2Y12 inhibitors in patients with ST-segment elevation myocardial infarction referred to primary percutaneous coronary interventions in Poland: Data from ORPKI national registry. Journal of Thrombosis and Thrombolysis, 2018, 45, 151-157.	1.0	13
34	Predictive utility of NT-pro BNP for infarct size and left ventricle function after acute myocardial infarction in long-term follow-up. Disease Markers, 2013, 34, 199-204.	0.6	13
35	Poland: coronary and structural heart interventions from 2010 to 2015. EuroIntervention, 2017, 13, Z51-Z54.	1.4	13
36	More aggressive pharmacological treatment may improve clinical outcome in patients with non-ST-elevation acute coronary syndromes treated conservatively. Coronary Artery Disease, 2007, 18, 299-303.	0.3	12

#	Article	IF	CITATIONS
37	Predictors and in-hospital outcomes of cardiogenic shock on admission in patients with acute coronary syndromes admitted to hospitals without on-site invasive facilities. Acute Cardiac Care, 2010, 12, 3-9.	0.2	12
38	Drug-eluting versus bare-metal stents in ST-segment elevation myocardial infarction: a mortality analysis from the EUROTRANSFER Registry. Clinical Research in Cardiology, 2011, 100, 139-145.	1.5	12
39	Predictive Utility of NT-pro BNP for Infarct Size and Left Ventricle Function after Acute Myocardial Infarction in Long-Term Follow-Up. Disease Markers, 2013, 34, 199-204.	0.6	12

The network of invasive cardiology facilities in Poland in 2016 (data from the ORPKI Polish National) Tj ETQq000 rgBT /Overlock 10 Tf 5 0.3 /Overlock 10 Tf 5

41	Management and mortality in patients with non-ST-segment elevation vs. ST-segment elevation myocardial infarction. Data from the Malopolska Registry of Acute Coronary Syndromes. Kardiologia Polska, 2009, 67, 115-20; discussion 121-2.	0.3	12
42	ST-segment resolution assessed immediately after primary percutaneous coronary intervention correlates with infarct size and left ventricular function in cardiac magnetic resonance at 1-year follow-up. Journal of Electrocardiology, 2009, 42, 152-156.	0.4	11
43	Percutaneous interventions in cardiology in Poland in the year 2014. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society AISN PTK. Postepy W Kardiologii Interwencyjnej, 2015, 3, 177-181.	0.1	11
44	Twelve months clinical outcome after bioresorbable vascular scaffold implantation in patients with stable angina and acute coronary syndrome. Data from the Polish National Registry. Postepy W Kardiologii Interwencyjnej, 2016, 2, 108-115.	0.1	11
45	Interventional cardiology in Poland in 2020 – impact of the COVID-19 pandemic. Annual summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society and Jagiellonian University Medical College*. Postepy W Kardiologii Interwencyjnej, 2021, 17, 131-134.	0.1	11
46	Chronic obstructive pulmonary disease affects the angiographic presentation and outcomes of patients with coronary artery disease treated with percutaneous coronary interventions. Polish Archives of Internal Medicine, 2017, 128, 24-34.	0.3	11
47	Treatment Delay and Clinical Outcomes in Patients with ST-Segment Elevation Myocardial Infarction during the COVID-19 Pandemic. Journal of Clinical Medicine, 2021, 10, 3920.	1.0	10
48	Prevalence and clinical presentation of myocardial bridge on the basis of the National Polish Percutaneous Interventions Registry and the Classification of Rare Cardiovascular Diseases. Kardiologia Polska, 2019, 77, 465-470.	0.3	10
49	Early abciximab use in ST-elevation myocardial infarction treated with primary percutaneous coronary intervention improves long-term outcome. Data from EUROTRANSFER Registry. Kardiologia Polska, 2010, 68, 539-43.	0.3	10
50	Early administration of abciximab reduces mortality in female patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention (from the EUROTRANSFER) Tj ETQq0 0 0 rgB <sup>-</sup>	⊺/Otwoerlocl	k 19 Tf 50 2
51	Patient profile and periprocedural outcomes of bioresorbable vascular scaffold implantation in comparison with drug-eluting and bare-metal stent implantation. Experience from ORPKI Polish National Registry 2014–2015. Postepy W Kardiologii Interwencyjnej, 2016, 4, 321-328.	0.1	9
52	Interventional cardiology procedures in Poland in 2018. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society (AISN PTK) and Jagiellonian University Medical College. Postepy W Kardiologii Interwencyjnej, 2019, 15, 391-393.	0.1	9
53	Long-term clinical outcomes in patients with unstable angina undergoing percutaneous coronary interventions in a contemporary registry data from Poland. Coronary Artery Disease, 2020, 31, 215-221.	0.3	9
54	Long-term follow-up of mesh-covered stent implantation in patients with ST-segment elevation myocardial infarction. Kardiologia Polska, 2014, 72, 140-145.	0.3	9

#	Article	IF	CITATIONS
55	The Polish Interventional Cardiology TAVI Survey (PICTS): adoption and practice of transcatheter aortic valve implantation in Poland. Postepy W Kardiologii Interwencyjnej, 2017, 1, 10-17.	0.1	8
56	Percutaneous interventions in cardiology in Poland in the year 2017. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society AISN PTK and Jagiellonian University Medical College. Postepy W Kardiologii Interwencyjnej, 2018, 14, 422-424.	0.1	8
57	Clinical outcomes in nonagenarians undergoing a percutaneous coronary intervention. Coronary Artery Disease, 2018, 29, 573-578.	0.3	8
58	Intima-media thickness and ankle-brachial index are correlated with the extent of coronary artery disease measured by the SYNTAX score. Postepy W Kardiologii Interwencyjnej, 2018, 14, 52-58.	0.1	8
59	Interventional cardiology in Poland in 2019. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society (AISN PTK) and Jagiellonian University Medical College*. Postepy W Kardiologii Interwencyjnej, 2020, 16, 123-126.	0.1	8
60	The relationship between increased air pollution expressed as PM10 concentration and the frequency of percutaneous coronary interventions in patients with acute coronary syndromes—a seasonal differences. Environmental Science and Pollution Research, 2020, 27, 21320-21330.	2.7	8
61	Radial approach reduces mortality in ST-segment elevation myocardial infarction with cardiogenic shock. Polish Archives of Internal Medicine, 2021, 131, 421-428.	0.3	8
62	Predictors of periprocedural complications in patients undergoing percutaneous coronary interventions within coronary artery bypass grafts. Cardiology Journal, 2020, 26, 633-644.	0.5	8
63	The Most Relevant Factors Affecting the Perioperative Death Rate in Patients with Acute Coronary Syndrome and COVID-19, Based on Annual Follow-Up in the ORPKI Registry. Biomedicines, 2021, 9, 1813.	1.4	8
64	Renal insufficiency increases mortality in acute coronary syndromes regardless of TIMI risk score. Kardiologia Polska, 2008, 66, 28-34; discussion 35-6.	0.3	8
65	In-hospital management and mortality in elderly patients with non-ST-segment elevation acute coronary syndromes treated in centers without on-site invasive facilities. Cardiology Journal, 2008, 15, 451-7.	0.5	8
66	Primary percutaneous coronary intervention during on- vs off-hours in patients with ST-elevation myocardial infarction. Results from EUROTRANSFER Registry. Kardiologia Polska, 2011, 69, 1017-22.	0.3	8
67	Early abciximab administration before primary percutaneous coronary intervention improves clinical outcome in elderly patients transferred with ST-elevation myocardial infarction. International Journal of Cardiology, 2010, 143, 147-153.	0.8	7
68	Bailout rotational atherectomy in patients with myocardial infarction is not associated with an increased periprocedural complication rate or poorer angiographic outcomes in comparison to elective procedures (from the ORPKI Polish National Registry 2015–2016). Postepy W Kardiologii Interwencyjnei, 2018, 14, 135-143.	0.1	7
69	Impact of On-Site Surgical Backup on Periprocedural Outcomes of Primary Percutaneous Interventions in Patients Presenting With ST-Segment Elevation Myocardial Infarction (From the ORPKI) Tj ETQq	1 1007784	31 <b>4</b> rgBT /Ov
70	Aspiration Thrombectomy in Patients with Acute Myocardial Infarction—5-Year Analysis Based on a Large National Registry (ORPKI). Journal of Clinical Medicine, 2020, 9, 3610.	1.0	7
71	Knowledge and prevalence of risk factors for coronary artery disease in patients after the first and repeated percutaneous coronary intervention. Kardiologia Polska, 2020, 78, 147-153.	0.3	7
72	Temporal trends and patterns in percutaneous treatment of coronary artery disease in Poland in the years 2005–2011. Kardiologia Polska, 2015, 73, 485-492.	0.3	7

#	Article	IF	CITATIONS
73	The efficacy of an education-based secondary outpatient prevention programme after acute coronary syndrome hospitalisations and treatment in Poland. The Patient Club initiative. Kardiologia Polska, 2016, 74, 185-191.	0.3	7
74	Impact of intra-aortic balloon pump on long-term mortality of unselected patients with ST-segment elevation myocardial infarction complicated by cardiogenic shock. Postepy W Kardiologii Interwencyjnej, 2014, 3, 175-180.	0.1	6
75	No longâ€ŧerm clinical benefit from manual aspiration thrombectomy in STâ€elevation myocardial infarction patients. Data from NRDES registry. Catheterization and Cardiovascular Interventions, 2015, 85, E16-22.	0.7	6
76	Chronic obstructive pulmonary disease and periprocedural complications in patients undergoing percutaneous coronary interventions. PLoS ONE, 2018, 13, e0204257.	1.1	6
77	Current trends and procedural outcomes in the era of rotational atherectomy expansion in Poland in the period 2014–2017 (based on the nationwide ORPKI registry). Postepy W Kardiologii Interwencyjnej, 2019, 15, 158-166.	0.1	6
78	Clinical presentation and 3-year outcomes of patients with acute coronary syndromes and non-obstructive coronary arteries on angiography. PLoS ONE, 2020, 15, e0234735.	1.1	6
79	Impact of acute total occlusion of the culprit artery on outcome in NSTEMI based on the results of a large national registry. BMC Cardiovascular Disorders, 2021, 21, 297.	0.7	6
80	Management of myocardial infarction with ST-segment elevation in district hospitals without catheterisation laboratory-Acute Coronary Syndromes Registry of MaÅ,opolska 2002-2003. Kardiologia Polska, 2006, 64, 1053-60; discussion 1061-2.	0.3	6
81	Local hospital networks for STEMI treatment for a population of half a million inhabitants increase the use of invasive treatment of acute coronary syndromes to the European recommended level. The MaÅ,opolska Registry of Acute Coronary Syndromes 2005-2006. Kardiologia Polska, 2008, 66, 489-97, discussion 498-9.	0.3	6
82	Early abciximab administration before primary percutaneous coronary intervention improves clinical outcome in diabetic patients with ST-segment elevation myocardial infarction (EUROTRANSFER) Tj ETQq0 0 0 rg	gBT¢Qaverlo	ock510 Tf 50 3
83	The ACEF (age, creatinine, ejection fraction) score predicts ischemic and bleeding outcomes of patients with acute coronary syndromes treated conservatively. Postepy W Kardiologii Interwencyjnej, 2017, 2, 160-164.	0.1	5
84	Comparison of patient comfort after coronary angiography by standard arterial access approaches. Kardiologia Polska, 2016, 74, 68-74.	0.3	5
85	The Effect of Periprocedural Clinical Factors Related to the Course of STEMI in Men and Women Based on the National Registry of Invasive Cardiology Procedures (ORPKI) between 2014 and 2019. Journal of Clinical Medicine, 2021, 10, 5716.	1.0	5
86	Cost-effectiveness of a photopethysmographic procedure for screening for atrial fibrillation in 6 European countries. Health Economics Review, 2022, 12, 17.	0.8	5
87	Prevention of infective endocarditis during dental extractions among Polish dentists - a contemporary nationwide survey. Folia Medica Cracoviensia, 2019, 59, 5-12.	0.3	5
88	Spontaneous closure of aorta-to-right atrium fistula after septal occluder implantation. Journal of Cardiovascular Medicine, 2008, 9, 744-746.	0.6	4
89	Thr715Pro P-selectin polymorphism and P-selectin release in blood obtained from the bleeding time wounds in patients with deep-vein thrombosis. Thrombosis Research, 2009, 124, 248-250.	0.8	4
90	Transportation with very long transfer delays (>90 min) for facilitated PCI with reduced-dose fibrinolysis in patients with ST-segment elevation myocardial infarction. International Journal of Cardiology, 2010, 139, 218-227.	0.8	4

#	Article	IF	CITATIONS
91	Prothrombinase formation at the site of microvascular injury and aspirin resistance: The effect of simvastatin. Thrombosis Research, 2010, 125, 283-285.	0.8	4
92	Knowledge of chronic total occlusion among Polish interventional cardiologists. Postepy W Kardiologii Interwencyjnej, 2015, 2, 89-94.	0.1	4
93	Long-term quality of life and clinical outcomes in patients with resistant hypertension treated with renal denervation. Postepy W Kardiologii Interwencyjnej, 2016, 4, 329-333.	0.1	4
94	Diabetes and periprocedural outcomes in patients treated with rotablation during percutaneous coronary interventions. Cardiology Journal, 2020, 27, 152-161.	0.5	4
95	What do Polish interventional cardiologists know about indications and qualification for recanalisation of chronic total coronary artery occlusions?. Kardiologia Polska, 2015, 73, 722-729.	0.3	4
96	"Heart without smoke―educational campaign — the role of patient education in secondary prevention of cardiovascular disease. Kardiologia Polska, 2018, 76, 125-129.	0.3	4
97	Presence and characteristics of coronary artery fistulas among patients undergoing coronary angiography. Kardiologia Polska, 2019, 77, 1034-1039.	0.3	4
98	Comparación de seguridad y efectividad entre los accesos radiales derecho e izquierdo en la intervención coronaria percutánea. Revista Espanola De Cardiologia, 2022, 75, 119-128.	0.6	4
99	Annual operator volume among patients treated using percutaneous coronary interventions with rotational atherectomy and procedural outcomes: Analysis based on a large national registry. Catheterization and Cardiovascular Interventions, 2022, , .	0.7	4
100	Safety of dental extractions in patients on dual antiplatelet therapy – a meta-analysis. Postepy W Kardiologii Interwencyjnej, 2019, 15, 68-73.	0.1	3
101	Comparison of safety and effectiveness between the right and left radial artery approach in percutaneous coronary intervention. Revista Espanola De Cardiologia (English Ed ), 2020, 75, 119-119.	0.4	3
102	Cardiovascular risk in patients with plaque psoriasis and psoriatic arthritis without a clinically overt cardiovascular disease: the role of endothelial progenitor cells. Postepy Dermatologii I Alergologii, 2020, 37, 299-305.	0.4	3
103	Long-term outcomes of percutaneous coronary interventions within coronary artery bypass grafts. Archives of Medical Science, 2021, 17, 628-637.	0.4	3
104	Similar outcome of ST-elevation myocardial infarction patients treated with primary percutaneous coronary intervention regardless of presence of cardiac surgery on-site. Kardiologia Polska, 2014, 72, 949-953.	0.3	3
105	Five-year report from the Polish national registry on percutaneous coronary interventions with a focus on coronary artery perforations within chronic total occlusions. Postepy W Kardiologii Interwencyjnej, 2020, 16, 399-409.	0.1	3
106	Prehospital Clopidogrel Administration in Patients With ST-Segment Elevation Myocardial Infarction Treated With Primary PCI: Real-Life Experience From the Multicenter NRDES Registry. Journal of Invasive Cardiology, 2016, 28, E56-8.	0.4	3
107	Angiographic perfusion score assessed in patients with acute myocardial infarction is correlated with cardiac magnetic resonance infarct size and N-terminal pro-brain natriuretic peptide in 6-month follow-up. Journal of Thrombosis and Thrombolysis, 2010, 30, 441-445.	1.0	2
108	Introduction of new oral antiplatelet drugs in myocardial infarction hospital network: initial experience. Journal of Thrombosis and Thrombolysis, 2014, 37, 243-245.	1.0	2

#	Article	IF	CITATIONS
109	Transradial and Transfemoral Approach in Patients with Prior Coronary Artery Bypass Grafting. Journal of Clinical Medicine, 2020, 9, 764.	1.0	2
110	Nurse-managed education: the effectiveness of secondary prevention after acute coronary syndromes and the prevalence and predictors of dropout from aÂcardiac rehabilitation programme. Postepy W Kardiologii Interwencyjnej, 2021, 17, 46-53.	0.1	2
111	ST-segment elevation myocardial infarction with non-obstructive coronary arteries: Score derivation for prediction based on a large national registry. PLoS ONE, 2021, 16, e0254427.	1.1	2
112	New model of secondary cardiovascular prevention for patients after acute coronary syndromes in Poland with regard to Norwegian experiences. Kardiologia Polska, 2016, 74, 101-103.	0.3	2
113	Chronic obstructive pulmonary disease affects angiographic presentation and outcomes. Authors' reply Polish Archives of Internal Medicine, 2018, 128, 195-196.	0.3	2
114	Psoriasis is an independent predictor of increased risk of allergic reaction during percutaneous coronary interventions. Big data analysis from the Polish National PCI Registry (ORPKI). Cardiology Journal, 2020, 27, 278-284.	0.5	2
115	Impact of infarct related artery patency after early abciximab administration on one-year mortality in patients with ST-segment elevation myocardial infarction (data from the EUROTRANSFER Registry). Kardiologia Polska, 2012, 70, 215-21.	0.3	2
116	Recanalization of peripheral arteries by interventional cardiologists: Rationale and results. International Journal of Cardiology, 2008, 129, 304-306.	0.8	1
117	Long-term follow-up of percutaneous peripheral interventions in lower limb arteries in patients with acute coronary syndrome and diabetes. Postepy W Kardiologii Interwencyjnej, 2010, 3, 117-121.	0.1	1
118	Interventional cardiology in Poland in the year 2011. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society. Postepy W Kardiologii Interwencyjnej, 2012, 2, 102-113.	0.1	1
119	ST-elevation myocardial infarction with local infusion of abciximab using thrombectomy catheter in a patient with very late stent thrombosis. Postepy W Kardiologii Interwencyjnej, 2012, 4, 338-341.	0.1	1
120	Complete infarct-related artery revascularization in acute myocardial infarction patients. CORAMI Registry. Postepy W Kardiologii Interwencyjnej, 2015, 2, 84-88.	0.1	1
121	Transradial access and the risk of periprocedural stroke. American Heart Journal, 2017, 186, e5-e6.	1.2	1
122	Single and dual chamber pacemaker implantation in patients with left superior vena cava persistence – own experiences. Postepy W Kardiologii Interwencyjnej, 2017, 2, 170-172.	0.1	1
123	Comparison of clinical and echocardiographic outcomes and quality of life in patients with severe mitral regurgitation treated by MitraClip implantation or treated conservatively. Postepy W Kardiologii Interwencyjnej, 2018, 14, 291-298.	0.1	1
124	Impact of percutaneous invasive coronary procedures using a radial approach on endothelial function of radial artery. Postepy W Kardiologii Interwencyjnej, 2018, 14, 95-98.	0.1	1
125	TCT-250 Prevalence and Predictors of Coronary Artery Perforation During Percutaneous Coronary Interventions: Data From the ORPKI National Registry inÂPoland. Journal of the American College of Cardiology, 2019, 74, B249.	1.2	1
126	TCT-839 The Relationship Between Winter Time and Increased Air Pollution Expressed as PM10 Concentration and the Frequency of Percutaneous Coronary Interventions in Patients With Acute Coronary Syndromes. Journal of the American College of Cardiology, 2019, 74, B822.	1.2	1

#	Article	IF	CITATIONS
127	Concomitant multi-vessel disease is associated with a lower procedural death rate in patients treated with percutaneous coronary interventions within the left main coronary artery (from the ORPKI) Tj ETQq1 1 0.78	43 <b>0.4</b> rgBT	- / <b>Q</b> verlock 10
128	From pharmacologically assisted early transfer to a universal primary angioplasty service: the experience of the Malopolska region. EuroIntervention, 2012, 8, P51-P54.	1.4	1
129	Statistics regarding interventional cardiology in Poland in 2013. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society (AISN PTK). Kardiologia Polska, 2014, 72, 1402-1407.	0.3	1
130	The impact of multiple stent implantation in the infarct-related artery on one-year clinical outcomes of patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. Data from the Polish NRDES Registry. Kardiologia Polska, 2016, 74, 717-725.	0.3	1
131	Predictors of infarct-related artery patency following combined lytic therapy in patients with ST-segment elevation myocardial infarction treated with immediate percutaneous coronary intervention. Kardiologia Polska, 2011, 69, 452-7.	0.3	1
132	Procedural Outcomes in Patients Treated with Percutaneous Coronary Interventions within Chronic Total Occlusions Stratified by Gender. Journal of Clinical Medicine, 2022, 11, 1419.	1.0	1
133	No clinical benefit from manual thrombus aspiration in patients with non-ST-elevation myocardial infarction. Postepy W Kardiologii Interwencyjnej, 2016, 1, 32-40.	0.1	0
134	TCT CONNECT-233 Coronary Artery Perforations in Patients Treated Using Percutaneous Coronary Interventions Within Chronic Total Occlusions: Analysis Based on a Large National Registry. Journal of the American College of Cardiology, 2020, 76, B102-B103.	1.2	0
135	How to Organize Networks for Invasive Treatment of STEMI. , 2010, , 30-35.		0
136	Authors' response. Kardiologia Polska, 2014, 72, 476-477.	0.3	0