

Sang-Won Lee

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6242133/sang-won-lee-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

193
papers

1,538
citations

21
h-index

30
g-index

224
ext. papers

1,937
ext. citations

3.4
avg, IF

4.9
L-index

#	Paper	IF	Citations
193	Gallic acid, a natural polyphenolic acid, induces apoptosis and inhibits proinflammatory gene expressions in rheumatoid arthritis fibroblast-like synoviocytes. <i>Joint Bone Spine</i> , 2013 , 80, 274-9	2.9	89
192	Measurement of the serum leptin level could assist disease activity monitoring in rheumatoid arthritis. <i>Rheumatology International</i> , 2007 , 27, 537-40	3.6	77
191	Differences in clinical manifestations and outcomes between adult and child patients with Henoch-Schönlein purpura. <i>Journal of Korean Medical Science</i> , 2014 , 29, 198-203	4.7	76
190	The role of (18) F-fluorodeoxyglucose-positron emission tomography in the assessment of disease activity in patients with takayasu arteritis. <i>Arthritis and Rheumatism</i> , 2012 , 64, 866-75		74
189	Leflunomide increases the risk of silent liver fibrosis in patients with rheumatoid arthritis receiving methotrexate. <i>Arthritis Research and Therapy</i> , 2012 , 14, R232	5.7	34
188	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017 , 44, 996-1003	4.1	31
187	Treatment of Collagen-Induced Arthritis Using Immune Modulatory Properties of Human Mesenchymal Stem Cells. <i>Cell Transplantation</i> , 2016 , 25, 1057-72	4	31
186	Adipokines, inflammation, insulin resistance, and carotid atherosclerosis in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2013 , 15, R194	5.7	31
185	Usefulness of serum leucine-rich alpha-2 glycoprotein as a disease activity biomarker in patients with rheumatoid arthritis. <i>Journal of Korean Medical Science</i> , 2014 , 29, 1199-204	4.7	28
184	ROR γ -specific transcriptional interactomic inhibition suppresses autoimmunity associated with TH17 cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 18673-8	11.5	28
183	Malignancies in Korean patients with inflammatory myopathy. <i>Yonsei Medical Journal</i> , 2006 , 47, 519-23	3	28
182	Systemic immune-inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019 , 24, 711-717	2.2	27
181	Alleviation of rheumatoid arthritis by cell-transducible methotrexate upon transcutaneous delivery. <i>Biomaterials</i> , 2012 , 33, 1563-72	15.6	26
180	C-Reactive Protein to Serum Albumin Ratio Is an Independent Predictor of All-Cause Mortality in Patients with ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 865-871	3	25
179	Evaluation of Spleen Glucose Metabolism Using F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 507-513	8.9	24
178	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1028-1035	2.3	24
177	The mid-range of the adjusted level of ferritin can predict the chronic course in patients with adult onset Still's disease. <i>Journal of Rheumatology</i> , 2009 , 36, 156-62	4.1	23

176	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 216-221	5.3	21
175	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018 , 19, 187	2.7	21
174	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018 , 38, 1865-1871	3.6	21
173	Inhibition of glycogen synthase kinase-3 β suppresses inflammatory responses in rheumatoid arthritis fibroblast-like synoviocytes and collagen-induced arthritis. <i>Joint Bone Spine</i> , 2014 , 81, 240-6	2.9	21
172	Soluble receptor for advanced glycation end products alleviates nephritis in (NZB/NZW)F1 mice. <i>Arthritis and Rheumatism</i> , 2013 , 65, 1902-12		21
171	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 1147-1154	3.9	19
170	Cell-penetrating interactomic inhibition of nuclear factor-kappa B in a mouse model of postoperative cognitive dysfunction. <i>Scientific Reports</i> , 2017 , 7, 13482	4.9	17
169	Increased serum interleukin-32 levels in patients with Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2018 , 21, 2167-2174	2.3	17
168	E/E β ratio is more sensitive than E/A ratio for detection of left ventricular diastolic dysfunction in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2010 , 28, S12-7	2.2	17
167	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018 , 59, 452-456	3	16
166	The prevalence and the clinical relevance of anti-Ro52 in Korean patients with primary Sjögren's syndrome. <i>Rheumatology International</i> , 2012 , 32, 491-5	3.6	16
165	Anti-Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016 , 19, 897-902	2.3	16
164	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , 2015 , 17, 77	5.7	15
163	Delta neutrophil index as an early marker for differential diagnosis of adult-onset Still's disease and sepsis. <i>Yonsei Medical Journal</i> , 2014 , 55, 753-9	3	15
162	Inflammatory polyarthritis in a patient with psoriasis: is it psoriatic arthritis or rheumatoid arthritis?. <i>Korean Journal of Internal Medicine</i> , 2010 , 25, 224-6	2.5	14
161	Patients with early arthritis who fulfil the 1987 ACR classification criteria for rheumatoid arthritis but not the 2010 ACR/EULAR criteria. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1097-8	2.4	14
160	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 111-118	2.2	13
159	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 397-405	3	12

158	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1593-1605	2.3	12
157	Retroperitoneal fibrosis in 27 Korean patients: single center experience. <i>Journal of Korean Medical Science</i> , 2011 , 26, 985-90	4.7	12
156	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019 , 23, 223-230	2.5	12
155	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 85-87	2.2	12
154	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017 , 37, 685-694	3.6	11
153	The antibody atliximab attenuates collagen-induced arthritis by neutralizing AIMP1, an inflammatory cytokine that enhances osteoclastogenesis. <i>Biomaterials</i> , 2015 , 44, 45-54	15.6	11
152	Extravascular manifestations of Takayasu arteritis: focusing on the features shared with spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2018 , 20, 142	5.7	11
151	Secreted frizzled-related protein 5 suppresses inflammatory response in rheumatoid arthritis fibroblast-like synoviocytes through down-regulation of c-Jun N-terminal kinase. <i>Rheumatology</i> , 2014 , 53, 1704-11	3.9	11
150	Activation of liver X receptors suppresses inflammatory gene expressions and transcriptional corepressor clearance in rheumatoid arthritis fibroblast like synoviocytes. <i>Journal of Clinical Immunology</i> , 2013 , 33, 190-9	5.7	11
149	Usefulness of FDG PET/CT in the diagnosis of eosinophilic fasciitis. <i>Clinical Nuclear Medicine</i> , 2014 , 39, 801-2	1.7	11
148	Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , 2020 , 30, 70-77	3.3	11
147	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018 , 37, 2133-2141	3.9	10
146	Attenuation of nephritis in lupus-prone mice by thalidomide. <i>Rheumatology</i> , 2012 , 51, 2131-40	3.9	10
145	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 47-54	2.2	10
144	Five factor score of more than 1 is associated with relapse during the first 2 year-follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1261-1268	2.3	9
143	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017 , 19, 193	5.7	9
142	Active tuberculosis risk with tumor necrosis factor inhibitors after treating latent tuberculosis. <i>Journal of Clinical Rheumatology</i> , 2014 , 20, 68-73	1.1	9
141	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 487-494	3.9	9

140	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017 , 36, 2751-2759	3.9	8
139	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019 , 19, 90	3	8
138	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 797-804	2.3	8
137	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017 , 84, 583-587	2.9	8
136	Antineutrophil Cytoplasmic Antibody-Associated Vasculitis in Korea: A Narrative Review. <i>Yonsei Medical Journal</i> , 2019 , 60, 10-21	3	8
135	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018 , 13, e0205651	3.7	8
134	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 65-72	2.2	8
133	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 261-268	2.3	7
132	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019 , 39, 1937-1944	3.6	7
131	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	7
130	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018 , 38, 1095-1101	3.6	7
129	Atalizumab, a humanized anti-aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , 2019 , 220, 119408	15.6	7
128	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017 , 12, e0181590	3.7	7
127	Eosinophilic Granulomatosis with Polyangiitis: Experiences in Korean Patients. <i>Yonsei Medical Journal</i> , 2019 , 60, 705-712	3	7
126	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017 , 56, 795-801	3.9	6
125	Classification of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2019 , 26, 156	1.2	6
124	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019 , 60, 454-460	3	6
123	Intranuclear delivery of the transcription modulation domain of Tbet improved lupus nephritis in (NZB/NZW) F1 lupus-prone mice. <i>Kidney International</i> , 2018 , 93, 1118-1130	9.9	6

122	Serum leucine-rich α -glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018 , 486, 253-258	6.2	6
121	Concurrent use of methotrexate and celecoxib increases risk of silent liver fibrosis in rheumatoid arthritis patients with subclinical reduced kidney function. <i>Clinical Rheumatology</i> , 2014 , 33, 1415-23	3.9	6
120	Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017 , 96, e7431	1.8	6
119	Two cases of refractory thrombocytopenia in systemic lupus erythematosus that responded to intravenous low-dose cyclophosphamide. <i>Journal of Korean Medical Science</i> , 2013 , 28, 472-5	4.7	6
118	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	6
117	Rheumatoid factor false positivity in patients with ANCA-associated vasculitis not having medical conditions producing rheumatoid factor. <i>Clinical Rheumatology</i> , 2018 , 37, 2771-2779	3.9	6
116	Comparison of therapeutic efficacy between bortezomib and combination treatment of prednisolone and mycophenolate mofetil on nephritis in NZB/WF1 mice. <i>Clinical and Experimental Rheumatology</i> , 2010 , 28, 393-6	2.2	6
115	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017 , 58, 1128-1134	3	5
114	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019 , 2019, 7508240	3	5
113	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019 , 11,	6.7	5
112	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020 , 34, e23022	3	5
111	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018 , 37, 1333-1340	3.9	5
110	Glycated albumin increases with disease activity in rheumatoid factor positive rheumatoid arthritis patients with normal fasting glucose and HbA. <i>Joint Bone Spine</i> , 2017 , 84, 115-118	2.9	4
109	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019 , 38, 1685-1690	3.9	4
108	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 294-302	3	4
107	The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017 , 96, e7901	1.8	4
106	The concomitant use of meloxicam and methotrexate does not clearly increase the risk of silent kidney and liver damages in patients with rheumatoid arthritis. <i>Rheumatology International</i> , 2014 , 34, 833-40	3.6	4
105	Clinical images: osteoporotic calcaneal stress fractures mistaken for aggravation of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 2983		4

104	Typical 18-FDG-PET/CT Findings of Polymyalgia Rheumatica: A Case Report. <i>Journal of Rheumatic Diseases</i> , 2013 , 20, 113	1.2	4
103	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019 , 60, 1164-1173 ³		4
102	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020 , 19, 184	4.4	4
101	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020 , 40, 1089-1099	3.6	4
100	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 73	5.7	4
99	HBsAg-negative and anti-HBc-positive in eosinophilic granulomatosis with polyangiitis: a retrospective pilot study. <i>Rheumatology International</i> , 2018 , 38, 1531-1538	3.6	4
98	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 88-93	2.2	4
97	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019 , 39, 1859-1866	3.6	3
96	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019 , 39, 879-884	3.6	3
95	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017 , 12, e0184337	3.7	3
94	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019 , 38, 3485-3491	3.9	3
93	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010-2018. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
92	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013 , 23, 509-515	3.3	3
91	Clinical Manifestations of Korean Adult Patients with Henoch-Schleien Purpura. <i>The Journal of the Korean Rheumatism Association</i> , 2010 , 17, 133		3
90	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 251, 117-123	2.4	3
89	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020 , 61, 712-719	3	3
88	Lipid Profiles in Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis: A Cross-sectional Analysis. <i>Journal of Rheumatic Diseases</i> , 2020 , 27, 261-269	1.2	3
87	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020 , 40, 1063-1070	3.6	3

86	Serological Biomarkers and Indices for the Current Activity and Prognosis of ANCA-Associated Vasculitis: Experience in a Single Centre in Korea. <i>Yonsei Medical Journal</i> , 2021 , 62, 279-287	3	3
85	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 73-77	2.2	3
84	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019 , 39, 1359-1367	3.6	2
83	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020 , 39, 3009-3015	3.9	2
82	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020 , 20, 147	3.5	2
81	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 138	5.7	2
80	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020 , 21, 258	2.7	2
79	Clinical significance of ANCA positivity in patients with IgA vasculitis: a retrospective monocentric study. <i>Rheumatology International</i> , 2019 , 39, 1927-1936	3.6	2
78	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019 , 39, 1899-1905	3.6	2
77	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. <i>Clinical Rheumatology</i> , 2021 , 1	3.9	2
76	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020 , 61, 951-957	3	2
75	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020 , 61, 720-725	3	2
74	Red blood cell distribution width is useful in discriminating adult onset Still's disease and sepsis within 24 hours after hospitalization. <i>Korean Journal of Internal Medicine</i> , 2018 , 33, 1234-1240	2.5	2
73	Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Monocentric Study. <i>Journal of Clinical Rheumatology</i> , 2021 , 27, e330-e335	1.1	2
72	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients: Results From the Korean College of Rheumatology Biologics Registry. <i>Journal of Clinical Rheumatology</i> , 2020 , Publish Ahead of Print,	1.1	2
71	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13431	2.9	2
70	Unclassifiable repeated antineutrophil cytoplasmic antibody (ANCA) positivity in diseases other than ANCA-associated vasculitis. <i>Zeitschrift Fur Rheumatologie</i> , 2021 , 1	1.9	2
69	Increased prevalence rate of metabolic syndrome is an independent predictor of cardiovascular disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Rheumatology International</i> , 2021 , 1	3.6	2

68	Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , 2016 , 26, 576-82	3.3	2
67	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020 , 50, 1414-1420	5.3	2
66	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23731	3	2
65	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 165-166	2.2	2
64	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 115-120	2.2	2
63	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 533-539	2.2	2
62	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37 Suppl 117, 116-121	2.2	2
61	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 236-237	2.2	2
60	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis.. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
59	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019 , 39, 885-892	3.6	1
58	The Presence of Anti-ribonucleoprotein at Diagnosis Is Associated with the Flare during the First Follow-up Year in Korean Patients with Systemic Lupus Erythematosus. <i>Journal of Rheumatic Diseases</i> , 2016 , 23, 154	1.2	1
57	Clinical characteristics and long-term outcomes of Libman-Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020 , 29, 1115-1120	2.6	1
56	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis.. <i>Journal of Clinical Laboratory Analysis</i> , 2022 , e24237	3	1
55	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e13929	2.9	1
54	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 1016-1022	3	1
53	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e24048	3	1
52	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020 , 61, 970-975	3	1
51	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013 , 23, 509-15	3.3	1

50	Brain meningioma in a patient with systemic lupus erythematosus. <i>Yeungnam University Journal of Medicine</i> , 2016 , 33, 159	1.3	1
49	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020 , 39, 831-840	3.9	1
48	Total Haemolytic Complement Activity at Diagnosis as an Indicator of the Baseline Activity of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2021 , 28, 85-93 ^{1,2}	1.2	1
47	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 731-741	2.5	1
46	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23885	3	1
45	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019 , 39, 395-401	1.5	1
44	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021 , 53, 1631-1638	2.3	1
43	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021 , 21, 399-406	4.9	1
42	Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018 , 25, 131	1.2	1
41	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018 , 21, 1838-1843	2.3	1
40	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 1221-1232	2.5	1
39	Antineutrophil Cytoplasmic Antibody Positivity Is Associated with Vascular Involvement in Behçet's Disease. <i>Yonsei Medical Journal</i> , 2021 , 62, 149-158	3	1
38	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021 , 36, e120	4.7	1
37	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 494-502	3	1
36	Subclinical but significant liver fibrosis in patients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37 Suppl 117, 26-31	2.2	1
35	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39 Suppl 129, 88-93	2.2	1
34	Thyroid Dysfunction in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis: A Monocentric Retrospective Study. <i>Journal of Rheumatology</i> , 2019 , 46, 1248-1250	4.1	0
33	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019 , 20, e299-e308	4.9	0

32	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020 , 76, 407-412	3	○
31	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019 , 14, e0219970	3.7	○
30	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020 , 2020, 8847306	3.2	○
29	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 77	5.7	○
28	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021 , 40, 3703-3710	3.9	○
27	Risk of Stroke in Systemic Necrotizing Vasculitis: A Nationwide Study Using the National Claims Database. <i>Frontiers in Immunology</i> , 2021 , 12, 629902	8.4	○
26	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 441-446	2.5	○
25	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e14512	2.9	○
24	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021 , 40, 1423-1430	3.9	○
23	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021 , 35, 1761-1768	2.3	○
22	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021 , 2021, 6668884	4.3	○
21	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021 , 16, e0254226	3.7	○
20	New body mass index for predicting prognosis in patients with antineutrophil cytoplasmic antibody-associated vasculitis.. <i>Journal of Clinical Laboratory Analysis</i> , 2022 , e24357	3	○
19	Incidence and Patterns of Interstitial Lung Disease and Their Clinical Impact on Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Korean Single-Centre Observational Study. <i>Journal of Immunology Research</i> , 2022 , 2022, 1-7	4.5	○
18	Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017 , 24, 143	1.2	
17	Red nail folds and GottronB sign in a patient with leukemic infiltration of pelvic muscles. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015 , 13, 581-2	1.2	
16	Hepatic Sarcoidosis in a Patient with Chronic Hepatitis B Virus Infection. <i>Journal of Rheumatic Diseases</i> , 2015 , 22, 200	1.2	
15	Dermatomyositis Associated with Gallbladder Cancer. <i>Journal of Rheumatic Diseases</i> , 2014 , 21, 261	1.2	

- 14 A Case of Ankylosing Spondylitis with Follicular Lymphoma. *Journal of Rheumatic Diseases*, **2011**, 18, 315-1.2
- 13 Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. *Journal of Investigative Medicine*, **2021**, 69, 91-95 2.9
- 12 Association Between Idiopathic Cutaneous Leukocytoclastic Angiitis and ANCA-negative Microscopic Polyangiitis. *Journal of Rheumatic Diseases*, **2022**, 29, 40-45 1.2
- 11 Kikuchi-Fujimoto Disease with Adult Onset Still Disease. *The Journal of the Korean Rheumatism Association*, **2008**, 15, 58
- 10 A Case of Systemic Lupus Erythematosus Manifested as Anemia and Thrombocytopenia during Antithyroid Drug Treatment. *Korean Journal of Medicine*, **2012**, 82, 232 0.5
- 9 Incidence of Tuberculosis in Systemic Necrotizing Vasculitides: A Population-Based Study From an Intermediate-Burden Country. *Frontiers in Medicine*, **2020**, 7, 550004 4.9
- 8 Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. *Chinese Medical Journal*, **2021**, 134, 1168-1174 2.9
- 7 Efficacy of tacrolimus as maintenance therapy after cyclophosphamide for treating antineutrophil cytoplasmic antibody-associated vasculitis. *Medicine (United States)*, **2021**, 100, e26956 1.8
- 6 The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. *Clinics*, **2021**, 76, e2501 2.3
- 5 D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. *Clinical and Experimental Rheumatology*, **2021**, 39 Suppl 129, 94-100 2.2
- 4 Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. *Clinical and Experimental Rheumatology*, **2021**, 39 Suppl 131, 111-118 2.2
- 3 D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. *Clinical and Experimental Rheumatology*, **2021**, 39, 94-100 2.2
- 2 Clinical characteristics and long-term outcomes in patients with mixed Class III/IV + V and pure proliferative lupus nephritis: A single-center experience.. *Lupus*, **2022**, 9612033221088437 2.6
- 1 Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. *Clinical and Experimental Rheumatology*, **2021**, 39, 111-118 2.2