

Jose C Reyes

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6244089/publications.pdf>

Version: 2024-02-01

58
papers

4,618
citations

101543

36
h-index

155660

55
g-index

60
all docs

60
docs citations

60
times ranked

6038
citing authors

#	ARTICLE	IF	CITATIONS
1	Altered control of cellular proliferation in the absence of mammalian brahma (SNF2 ^{Δ±}). EMBO Journal, 1998, 17, 6979-6991.	7.8	400
2	The GATA Family of Transcription Factors in Arabidopsis and Rice. Plant Physiology, 2004, 134, 1718-1732.	4.8	331
3	Cyanobacteria Perceive Nitrogen Status by Sensing Intracellular 2-Oxoglutarate Levels. Journal of Biological Chemistry, 2001, 276, 38320-38328.	3.4	283
4	Ammonium assimilation in cyanobacteria. Photosynthesis Research, 2005, 83, 135-150.	2.9	241
5	Components of the Human SWI/SNF Complex Are Enriched in Active Chromatin and Are Associated with the Nuclear Matrix. Journal of Cell Biology, 1997, 137, 263-274.	5.2	216
6	Histone H2A.Z and homologues of components of the SWR1 complex are required to control immunity in Arabidopsis. Plant Journal, 2008, 53, 475-487.	5.7	209
7	Arsenic Sensing and Resistance System in the Cyanobacterium Synechocystis sp. Strain PCC 6803. Journal of Bacteriology, 2003, 185, 5363-5371.	2.2	165
8	The Arabidopsis thaliana SNF2 homolog AtBRM controls shoot development and flowering. Development (Cambridge), 2004, 131, 4965-4975.	2.5	152
9	CHD3 Proteins and Polycomb Group Proteins Antagonistically Determine Cell Identity in Arabidopsis. PLoS Genetics, 2009, 5, e1000605.	3.5	141
10	SUMO association with repressor complexes, emerging routes for transcriptional control. Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms, 2009, 1789, 451-459.	1.9	134
11	The Beauty of Being a Variant: H2A.Z and the SWR1 Complex in Plants. Molecular Plant, 2009, 2, 565-577.	8.3	130
12	The putative SWI/SNF complex subunit BRAHMA activates flower homeotic genes in Arabidopsis thaliana. Plant Molecular Biology, 2006, 62, 291-304.	3.9	121
13	SEF, a New Protein Required for Flowering Repression in Arabidopsis, Interacts with PIE1 and ARP6. Plant Physiology, 2007, 143, 893-901.	4.8	119
14	A two-component signal transduction system involved in nickel sensing in the cyanobacterium Synechocystis sp. PCC 6803. Molecular Microbiology, 2002, 43, 247-256.	2.5	113
15	NtcA represses transcription of gifA and gifB, genes that encode inhibitors of glutamine synthetase type I from Synechocystis sp. PCC 6803. Molecular Microbiology, 2000, 35, 1192-1201.	2.5	110
16	Chromatin-Remodeling and Memory Factors. New Regulators of Plant Development. Plant Physiology, 2002, 130, 1090-1101.	4.8	100
17	High expression of SMARCA4 or SMARCA2 is frequently associated with an opposite prognosis in cancer. Scientific Reports, 2018, 8, 2043.	3.3	100
18	A Gene Cluster Involved in Metal Homeostasis in the Cyanobacterium Synechocystis sp. Strain PCC 6803. Journal of Bacteriology, 2000, 182, 1507-1514.	2.2	97

#	ARTICLE	IF	CITATIONS
19	The Arabidopsis BRAHMA Chromatin-Remodeling ATPase Is Involved in Repression of Seed Maturation Genes in Leaves. <i>Plant Physiology</i> , 2008, 147, 1143-1157.	4.8	97
20	ras transformation is associated with decreased expression of the brm/SNF2alpha ATPase from the mammalian SWI-SNF complex. <i>EMBO Journal</i> , 1998, 17, 223-231.	7.8	95
21	The Glucocorticoid Receptor Interacting Protein 1 (GRIP1) Localizes in Discrete Nuclear Foci That Associate with ND10 Bodies and Are Enriched in Components of the 26S Proteasome. <i>Molecular Endocrinology</i> , 2001, 15, 485-500.	3.7	90
22	The CopRS Two-Component System Is Responsible for Resistance to Copper in the Cyanobacterium <i>Synechocystis</i> sp. PCC 6803. <i>Plant Physiology</i> , 2012, 159, 1806-1818.	4.8	88
23	The chromatin remodeller CHD8 is required for E2F-dependent transcription activation of S-phase genes. <i>Nucleic Acids Research</i> , 2014, 42, 2185-2196.	14.5	72
24	Chromatin modifiers that control plant development. <i>Current Opinion in Plant Biology</i> , 2006, 9, 21-27.	7.1	71
25	Control of neuronal differentiation by sumoylation of BRAF35, a subunit of the LSD1-CoREST histone demethylase complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 8085-8090.	7.1	68
26	Defective histone supply causes changes in RNA polymerase II elongation rate and cotranscriptional pre-mRNA splicing. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 14840-14845.	7.1	68
27	The PHD Domain of Plant PIAS Proteins Mediates Sumoylation of Bromodomain GTE Proteins. <i>Journal of Biological Chemistry</i> , 2008, 283, 21469-21477.	3.4	63
28	Brahma Is Required for Proper Expression of the Floral Repressor FLC in Arabidopsis. <i>PLoS ONE</i> , 2011, 6, e17997.	2.5	50
29	BRG1 helps RNA polymerase II to overcome a nucleosomal barrier during elongation, <i>in vivo</i> . <i>EMBO Reports</i> , 2010, 11, 751-757.	4.5	49
30	Analysis of the relationship between coexpression domains and chromatin 3D organization. <i>PLoS Computational Biology</i> , 2017, 13, e1005708.	3.2	49
31	Diverse functions of Polycomb group proteins during plant development. <i>Seminars in Cell and Developmental Biology</i> , 2003, 14, 77-84.	5.0	47
32	A positioned +1 nucleosome enhances promoter-proximal pausing. <i>Nucleic Acids Research</i> , 2015, 43, 3068-3078.	14.5	46
33	The Chromatin Remodeler CHD8 Is Required for Activation of Progesterone Receptor-Dependent Enhancers. <i>PLoS Genetics</i> , 2015, 11, e1005174.	3.5	44
34	p21 as a Transcriptional Co-Repressor of S-Phase and Mitotic Control Genes. <i>PLoS ONE</i> , 2012, 7, e37759.	2.5	42
35	Purification and Characterization of A New Type of Glutamine Synthetase from Cyanobacteria. <i>FEBS Journal</i> , 1997, 244, 258-264.	0.2	41
36	The inactivating factor of glutamine synthetase, IF7, is a natively unfolded protein. <i>Protein Science</i> , 2003, 12, 1443-1454.	7.6	39

#	ARTICLE	IF	CITATIONS
37	The Many Faces of Plant SWI/SNF Complex. <i>Molecular Plant</i> , 2014, 7, 454-458.	8.3	38
38	The type 2 diabetes-associated HMG20A gene is mandatory for islet beta cell functional maturity. <i>Cell Death and Disease</i> , 2018, 9, 279.	6.3	36
39	The Cornelia de Lange Syndrome-associated factor NIPBL interacts with BRD4 ET domain for transcription control of a common set of genes. <i>Cell Death and Disease</i> , 2019, 10, 548.	6.3	35
40	Expression of TDRD9 in a subset of lung carcinomas by CpG island hypomethylation protects from DNA damage. <i>Oncotarget</i> , 2018, 9, 9618-9631.	1.8	29
41	Histone availability as a strategy to control gene expression. <i>RNA Biology</i> , 2017, 14, 281-286.	3.1	27
42	The ammonium-inactivated cyanobacterial glutamine synthetase I is reactivated in vivo by a mechanism involving proteolytic removal of its inactivating factors. <i>Molecular Microbiology</i> , 2007, 65, 166-179.	2.5	23
43	PML and COP1 are two proteins with much in common. <i>Trends in Biochemical Sciences</i> , 2001, 26, 18-20.	7.5	22
44	TGF β 2 promotes widespread enhancer chromatin opening and operates on genomic regulatory domains. <i>Nature Communications</i> , 2020, 11, 6196.	12.8	21
45	The Glucocorticoid Receptor Interacting Protein 1 (GRIP1) Localizes in Discrete Nuclear Foci That Associate with ND10 Bodies and Are Enriched in Components of the 26S Proteasome. <i>Molecular Endocrinology</i> , 2001, 15, 485-500.	3.7	20
46	Tracking Components of the Transcription Apparatus in Living Cells. <i>Methods</i> , 1999, 19, 353-361.	3.8	17
47	The metabesity factor HMG20A potentiates astrocyte survival and reactive astrogliosis preserving neuronal integrity. <i>Theranostics</i> , 2021, 11, 6983-7004.	10.0	16
48	Chromatin structure and pre-mRNA processing work together. <i>Transcription</i> , 2016, 7, 63-68.	3.1	11
49	Dissecting the Brain/Islet Axis in Metabesity. <i>Genes</i> , 2019, 10, 350.	2.4	11
50	To cross or not to cross the nucleosome, that is the elongation question. <i>RNA Biology</i> , 2011, 8, 389-393.	3.1	8
51	TBL1 is required for the mesenchymal phenotype of transformed breast cancer cells. <i>Cell Death and Disease</i> , 2019, 10, 95.	6.3	6
52	Human prefoldin modulates co-transcriptional pre-mRNA splicing. <i>Nucleic Acids Research</i> , 2021, 49, 6267-6280.	14.5	5
53	Co-transcriptional splicing efficiency is a gene-specific feature that can be regulated by TGF β 2. <i>Communications Biology</i> , 2022, 5, 277.	4.4	4
54	Ammonium assimilation in cyanobacteria. The Regulation of the GS-GOGAT Pathway. , 1998, , 3607-3612.		2

#	ARTICLE	IF	CITATIONS
55	SENP7 overexpression protects cancer cells from oxygen and glucose deprivation and associates with poor prognosis in colon cancer. <i>Genes and Diseases</i> , 2022, 9, 1419-1422.	3.4	2
56	Regulation of Ammonium Assimilation in Cyanobacteria. , 2002, , 93-113.		1
57	Deciphering CHFR Role in Pancreatic Ductal Adenocarcinoma. <i>Frontiers in Medicine</i> , 2021, 8, 720128.	2.6	1
58	213-OR: Obesity-Induced Astrogliosis Is Regulated by the Diabesity Factor HMG20A. <i>Diabetes</i> , 2020, 69, .	0.6	0