

Maria Criscuolo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6251363/publications.pdf>

Version: 2024-02-01

11
papers

422
citations

933447

10
h-index

1281871

11
g-index

11
all docs

11
docs citations

11
times ranked

849
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Molecular analysis of Fanconi anemia: the experience of the Bone Marrow Failure Study Group of the Italian Association of Pediatric Onco-Hematology. <i>Haematologica</i> , 2014, 99, 1022-1031. | 3.5 | 44 |
| 2 | p57Kip2 and Cancer: Time for a Critical Appraisal. <i>Molecular Cancer Research</i> , 2011, 9, 1269-1284. | 3.4 | 81 |
| 3 | Targeting p27 ^{Kip1} protein: its relevance in the therapy of human cancer. <i>Expert Opinion on Therapeutic Targets</i> , 2011, 15, 677-693. | 3.4 | 52 |
| 4 | Histone deacetylase inhibitors upregulate p57Kip2 level by enhancing its expression through Sp1 transcription factor. <i>Carcinogenesis</i> , 2007, 29, 560-567. | 2.8 | 25 |
| 5 | A novel Leu153Ser mutation of the Fanconi anemia FANCD2 gene is associated with severe chemotherapy toxicity in a pediatric T-cell acute lymphoblastic leukemia. <i>Leukemia</i> , 2007, 21, 72-78. | 7.2 | 21 |
| 6 | Retinoic Acid Induces p27Kip1 Nuclear Accumulation by Modulating Its Phosphorylation. <i>Cancer Research</i> , 2006, 66, 4240-4248. | 0.9 | 37 |
| 7 | Retinoic acid inhibits the growth of bone marrow mesenchymal stem cells and induces p27Kip1 and p16INK4A up-regulation. <i>Molecular and Cellular Biochemistry</i> , 2003, 247, 55-60. | 3.1 | 17 |
| 8 | Spectrum of FANCA mutations in Italian Fanconi anemia patients: Identification of six novel alleles and phenotypic characterization of the S858R variant. <i>Human Mutation</i> , 2003, 22, 338-339. | 2.5 | 35 |
| 9 | Infant hypervitaminosis A causes severe anemia and thrombocytopenia: evidence of a retinol-dependent bone marrow cell growth inhibition. <i>Blood</i> , 2002, 99, 2017-2022. | 1.4 | 38 |
| 10 | Cell division cycle control in embryonal and alveolar rhabdomyosarcomas. <i>European Journal of Cancer</i> , 2002, 38, 2290-2299. | 2.8 | 9 |
| 11 | p27Kip1 accumulation is associated with retinoic-induced neuroblastoma differentiation: evidence of a decreased proteasome-dependent degradation. <i>Oncogene</i> , 2000, 19, 51-60. | 5.9 | 63 |