

M. Azizur Rahman

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6254876/publications.pdf>

Version: 2024-02-01

56
papers

3,299
citations

218381
26
h-index

182168
51
g-index

56
all docs

56
docs citations

56
times ranked

3847
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of droughts on child mortality: a case study in Southern African countries. <i>Natural Hazards</i> , 2021, 108, 2211-2224.	1.6	7
2	Freshwater phytoplankton: biotransformation of inorganic arsenic to methylarsenic and organoarsenic. <i>Scientific Reports</i> , 2019, 9, 12074.	1.6	27
3	Permeable pavement as a stormwater best management practice: a review and discussion. <i>Environmental Earth Sciences</i> , 2019, 78, 1.	1.3	54
4	Assessing the ecological health status using macrobenthic communities of tropical coastal water. <i>Human and Ecological Risk Assessment (HERA)</i> , 2018, 24, 1761-1785.	1.7	2
5	Human health risks and socio-economic perspectives of arsenic exposure in Bangladesh: A scoping review. <i>Ecotoxicology and Environmental Safety</i> , 2018, 150, 335-343.	2.9	127
6	Effect of biodegradable chelating ligands on Fe uptake in and growth of marine microalgae. <i>Journal of Applied Phycology</i> , 2018, 30, 2215-2225.	1.5	4
7	The Response of Macrobenthic Communities to Environmental Variability in Tropical Coastal Waters. <i>Estuaries and Coasts</i> , 2018, 41, 1178-1192.	1.0	5
8	Inorganic arsenic in rice and rice-based diets: Health risk assessment. <i>Food Control</i> , 2017, 82, 196-202.	2.8	66
9	Phytoremediation of Toxic Metals in Soils and Wetlands: Concepts and Applications. , 2016, , 161-195.		26
10	Ecotoxicological Effects of an Arsenic Remediation Method on Three Freshwater Organisms— <i>Lemna disperma</i> , <i>Chlorella</i> sp. CE-35 and <i>Ceriodaphnia</i> cf. <i>dubia</i> . <i>Water, Air, and Soil Pollution</i> , 2015, 226, 1.	1.1	1
11	Integrated ecological risk assessment of dioxin compounds. <i>Environmental Science and Pollution Research</i> , 2015, 22, 11193-11208.	2.7	29
12	A Fluorescent-Based HPLC Assay Using 4-Chloro-7-nitrobenzo-2-oxa-1, 3-diazole as Derivatization Agent for the Determination of Iron Bioavailability to Red Tide Phytoplankton. <i>Chromatographia</i> , 2015, 78, 65-72.	0.7	0
13	Economic efficiency of different light wavelengths and intensities using LEDs for the cultivation of green microalga <i>Botryococcus braunii</i> (NIES-836) for biofuel production. <i>Environmental Progress and Sustainable Energy</i> , 2015, 34, 269-275.	1.3	22
14	Arsenic in Rice. , 2014, , 365-375.		4
15	Distribution of trace element in Japanese red coral <i>Paracorallium japonicum</i> by μ -XRF and sulfur speciation by XANES: Linkage between trace element distribution and growth ring formation. <i>Geochimica Et Cosmochimica Acta</i> , 2014, 127, 1-9.	1.6	24
16	New Citrate-Bicarbonate-Ethylenediaminetetraacetate (CBE) Method for Chemical Extraction of Hydrous Iron Oxides from Plant Root Surfaces. <i>Communications in Soil Science and Plant Analysis</i> , 2014, 45, 1760-1771.	0.6	1
17	Arsenic Speciation in Australian-Grown and Imported Rice on Sale in Australia: Implications for Human Health Risk. <i>Journal of Agricultural and Food Chemistry</i> , 2014, 62, 6016-6024.	2.4	78
18	Is arsenic biotransformation a detoxification mechanism for microorganisms?. <i>Aquatic Toxicology</i> , 2014, 146, 212-219.	1.9	108

#	ARTICLE	IF	CITATIONS
19	Human health risk of polycyclic aromatic hydrocarbons from consumption of blood cockle and exposure to contaminated sediments and water along the Klang Strait, Malaysia. <i>Marine Pollution Bulletin</i> , 2014, 84, 268-279.	2.3	33
20	Toxicity of arsenic species to three freshwater organisms and biotransformation of inorganic arsenic by freshwater phytoplankton (<i>Chlorella</i> sp. CE-35). <i>Ecotoxicology and Environmental Safety</i> , 2014, 106, 126-135.	2.9	64
21	Heavy metals in Australian grown and imported rice and vegetables on sale in Australia: Health hazard. <i>Ecotoxicology and Environmental Safety</i> , 2014, 100, 53-60.	2.9	195
22	A marine phytoplankton (<i>Prymnesium parvum</i>) up-regulates ABC transporters and several other proteins to acclimatize with Fe-limitation. <i>Chemosphere</i> , 2014, 95, 213-219.	4.2	4
23	Effect of nitrate on the determination of iron concentration in phytoplankton culture medium by liquid scintillation counting (LSC) method using ⁵⁵ Fe as radioisotope tracer. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , 2013, 296, 1295-1302.	0.7	1
24	Growth characteristics and growth rate estimation of Japanese precious corals. <i>Journal of Experimental Marine Biology and Ecology</i> , 2013, 441, 117-125.	0.7	20
25	Effect of Iron (Fe ²⁺) Concentration in Soil on Arsenic Uptake in Rice Plant (<i>Oryza sativa</i> L.) when Grown with Arsenate [As(V)] and Dimethylarsinate (DMA). <i>Water, Air, and Soil Pollution</i> , 2013, 224, 1.	1.1	5
26	Arsenic in freshwater systems: Influence of eutrophication on occurrence, distribution, speciation, and bioaccumulation. <i>Applied Geochemistry</i> , 2012, 27, 304-314.	1.4	83
27	Bioaccumulation, biotransformation and trophic transfer of arsenic in the aquatic food chain. <i>Environmental Research</i> , 2012, 116, 118-135.	3.7	290
28	Significance of the concentration of chelating ligands on Fe ³⁺ -solubility, bioavailability, and uptake in rice plant. <i>Plant Physiology and Biochemistry</i> , 2012, 58, 205-211.	2.8	14
29	The significance of biodegradable methylglycinediacetic acid (MGDA) for iron and arsenic bioavailability and uptake in rice plant. <i>Soil Science and Plant Nutrition</i> , 2012, 58, 627-636.	0.8	2
30	Phytotoxicity of Arsenate and Salinity on Early Seedling Growth of Rice (<i>Oryza sativa</i> L.): A Threat to Sustainable Rice Cultivation in South and South-East Asia. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2012, 88, 695-702.	1.3	9
31	Trace elements in <i>Corallium</i> spp. as indicators for origin and habitat. <i>Journal of Experimental Marine Biology and Ecology</i> , 2012, 414-415, 1-5.	0.7	19
32	Arsenic-Induced Straighthead: An Impending Threat to Sustainable Rice Production in South and South-East Asia!. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2012, 88, 311-315.	1.3	15
33	High levels of inorganic arsenic in rice in areas where arsenic-contaminated water is used for irrigation and cooking. <i>Science of the Total Environment</i> , 2011, 409, 4645-4655.	3.9	196
34	Transport of DMAA and MMAA into rice (<i>Oryza sativa</i> L.) roots. <i>Environmental and Experimental Botany</i> , 2011, 72, 41-46.	2.0	42
35	Influence of chelating ligands on bioavailability and mobility of iron in plant growth media and their effect on radish growth. <i>Environmental and Experimental Botany</i> , 2011, 71, 345-351.	2.0	22
36	Influence of aggregated particles on biodegradation activities for dimethylarsinic acid (DMA) in Lake Kahokugata. <i>Chemosphere</i> , 2011, 83, 1486-1492.	4.2	0

#	ARTICLE	IF	CITATIONS
37	Aquatic arsenic: Phytoremediation using floating macrophytes. <i>Chemosphere</i> , 2011, 83, 633-646.	4.2	310
38	Effect of external iron and arsenic species on chelant-enhanced iron bioavailability and arsenic uptake in rice (<i>Oryza sativa</i> L.). <i>Chemosphere</i> , 2011, 84, 439-445.	4.2	20
39	Stagnant surface water bodies (SSWBs) as an alternative water resource for the Chittagong metropolitan area of Bangladesh: physicochemical characterization in terms of water quality indices. <i>Environmental Monitoring and Assessment</i> , 2011, 173, 669-684.	1.3	16
40	Seasonal changes of arsenic speciation in lake waters in relation to eutrophication. <i>Science of the Total Environment</i> , 2010, 408, 1684-1690.	3.9	72
41	EFFECT OF BIODEGRADABLE CHELATING LIGAND ON IRON BIOAVAILABILITY AND RADISH GROWTH. <i>Journal of Plant Nutrition</i> , 2010, 33, 933-942.	0.9	3
42	Effect of eutrophication on the distribution of arsenic species in eutrophic and mesotrophic lakes. <i>Science of the Total Environment</i> , 2009, 407, 1418-1425.	3.9	52
43	PHYSICOCHEMICAL PROPERTIES OF <i>MORINGA OLEIFERA</i> LAM. SEED OIL OF THE INDIGENOUS CULTIVAR OF BANGLADESH. <i>Journal of Food Lipids</i> , 2009, 16, 540-553.	0.9	43
44	Seasonal dynamics of biodegradation activities for dimethylarsinic acid (DMA) in Lake Kahokugata. <i>Chemosphere</i> , 2009, 77, 36-42.	4.2	10
45	Hydroxyiminodisuccinic acid (HIDS): A novel biodegradable chelating ligand for the increase of iron bioavailability and arsenic phytoextraction. <i>Chemosphere</i> , 2009, 77, 207-213.	4.2	22
46	Influence of Chelating Ligands on Arsenic Uptake by Hydroponically Grown Rice Seedlings (<i>Oryza</i>) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i>	0.7	4
47	Arsenic uptake by aquatic macrophyte <i>Spirodela polyrhiza</i> L.: Interactions with phosphate and iron. <i>Journal of Hazardous Materials</i> , 2008, 160, 356-361.	6.5	67
48	Influence of phosphate and iron ions in selective uptake of arsenic species by water fern (<i>Salvinia</i>) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i>	6.6	49
49	Straighthead disease of rice (<i>Oryza sativa</i> L.) induced by arsenic toxicity. <i>Environmental and Experimental Botany</i> , 2008, 62, 54-59.	2.0	75
50	Arsenic accumulation in rice (<i>Oryza sativa</i> L.): Human exposure through food chain. <i>Ecotoxicology and Environmental Safety</i> , 2008, 69, 317-324.	2.9	186
51	Influence of EDTA and chemical species on arsenic accumulation in <i>Spirodela polyrhiza</i> L. (duckweed). <i>Ecotoxicology and Environmental Safety</i> , 2008, 70, 311-318.	2.9	25
52	Effect of arsenic on photosynthesis, growth and yield of five widely cultivated rice (<i>Oryza sativa</i> L.) varieties in Bangladesh. <i>Chemosphere</i> , 2007, 67, 1072-1079.	4.2	228
53	Arsenic accumulation in duckweed (<i>Spirodela polyrhiza</i> L.): A good option for phytoremediation. <i>Chemosphere</i> , 2007, 69, 493-499.	4.2	120
54	Accumulation of arsenic in tissues of rice plant (<i>Oryza sativa</i> L.) and its distribution in fractions of rice grain. <i>Chemosphere</i> , 2007, 69, 942-948.	4.2	268

#	ARTICLE	IF	CITATIONS
55	Arsenic Accumulation in Rice (<i>Oryza sativa</i> L.) Varieties of Bangladesh: A Glass House Study. <i>Water, Air, and Soil Pollution</i> , 2007, 185, 53-61.	1.1	31
56	Influence of cooking method on arsenic retention in cooked rice related to dietary exposure. <i>Science of the Total Environment</i> , 2006, 370, 51-60.	3.9	99