

# Michael Ailion

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6257331/michael-ailion-publications-by-year.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

43  
papers

2,547  
citations

23  
h-index

50  
g-index

56  
ext. papers

3,057  
ext. citations

6.9  
avg, IF

4.65  
L-index

#	Paper	IF	Citations
43	Dopamine receptor DOP-1 engages a sleep pathway to modulate swimming in. <i>IScience</i> , <b>2021</b> , 24, 102247.1	4.1	2
42	EIPR1 controls dense-core vesicle cargo retention and EARP complex localization in insulin-secreting cells. <i>Molecular Biology of the Cell</i> , <b>2020</b> , 31, 59-79	3.5	5
41	Casein Kinase 1 $\beta$ Stabilizes Mature Axons by Inhibiting Transcription Termination of Ankyrin. <i>Developmental Cell</i> , <b>2020</b> , 52, 88-103.e18	10.2	3
40	Hybridization promotes asexual reproduction in Caenorhabditis nematodes. <i>PLoS Genetics</i> , <b>2019</b> , 15, e1008520	6	2
39	Modulation of Gq-Rho Signaling by the ERK MAPK Pathway Controls Locomotion in. <i>Genetics</i> , <b>2018</b> , 209, 523-535	4	4
38	Pristionchus nematodes occur frequently in diverse rotting vegetal substrates and are not exclusively necromenic, while Panagrellus redivivoides is found specifically in rotting fruits. <i>PLoS ONE</i> , <b>2018</b> , 13, e0200851	3.7	17
37	The NCA-1 and NCA-2 Ion Channels Function Downstream of G and Rho To Regulate Locomotion in. <i>Genetics</i> , <b>2017</b> , 206, 265-282	4	17
36	The SEK-1 p38 MAP Kinase Pathway Modulates Gq Signaling in. <i>G3: Genes, Genomes, Genetics</i> , <b>2017</b> , 7, 2979-2989	3.2	5
35	Genetics: Master Regulator or Master of Disguise?. <i>Current Biology</i> , <b>2017</b> , 27, R844-R847	6.3	1
34	The dense-core vesicle maturation protein CCCP-1 binds RAB-2 and membranes through its C-terminal domain. <i>Traffic</i> , <b>2017</b> , 18, 720-732	5.7	10
33	Cytoplasmic-Nuclear Incompatibility Between Wild Isolates of. <i>G3: Genes, Genomes, Genetics</i> , <b>2017</b> , 7, 823-834	3.2	9
32	Dopamine negatively modulates the NCA ion channels in C. elegans. <i>PLoS Genetics</i> , <b>2017</b> , 13, e1007032	6	15
31	The Conserved VPS-50 Protein Functions in Dense-Core Vesicle Maturation and Acidification and Controls Animal Behavior. <i>Current Biology</i> , <b>2016</b> , 26, 862-71	6.3	18
30	The EARP Complex and Its Interactor EIPR-1 Are Required for Cargo Sorting to Dense-Core Vesicles. <i>PLoS Genetics</i> , <b>2016</b> , 12, e1006074	6	29
29	Two Rab2 interactors regulate dense-core vesicle maturation. <i>Neuron</i> , <b>2014</b> , 82, 167-80	13.9	41
28	The membrane-associated proteins FCHO and SGIP are allosteric activators of the AP2 clathrin adaptor complex. <i>ELife</i> , <b>2014</b> , 3,	8.9	49
27	Improved Mos1-mediated transgenesis in C. elegans. <i>Nature Methods</i> , <b>2012</b> , 9, 117-8	21.6	266

26	Parallel evolution of domesticated <i>Caenorhabditis</i> species targets pheromone receptor genes. <i>Nature</i> , <b>2011</b> , 477, 321-5	50.4	182
25	A phylogeny and molecular barcodes for <i>Caenorhabditis</i> , with numerous new species from rotting fruits. <i>BMC Evolutionary Biology</i> , <b>2011</b> , 11, 339	3	239
24	Neuron-specific proteotoxicity of mutant ataxin-3 in <i>C. elegans</i> : rescue by the DAF-16 and HSF-1 pathways. <i>Human Molecular Genetics</i> , <b>2011</b> , 20, 2996-3009	5.6	74
23	A novel sperm-delivered toxin causes late-stage embryo lethality and transmission ratio distortion in <i>C. elegans</i> . <i>PLoS Biology</i> , <b>2011</b> , 9, e1001115	9.7	108
22	Genetics of extracellular matrix remodeling during organ growth using the <i>Caenorhabditis elegans</i> pharynx model. <i>Genetics</i> , <b>2010</b> , 186, 969-82	4	18
21	<i>C. elegans</i> anaplastic lymphoma kinase ortholog SCD-2 controls dauer formation by modulating TGF-beta signaling. <i>Current Biology</i> , <b>2008</b> , 18, 1101-9	6.3	51
20	Ammonium-acetate is sensed by gustatory and olfactory neurons in <i>Caenorhabditis elegans</i> . <i>PLoS ONE</i> , <b>2008</b> , 3, e2467	3.7	15
19	UNC-31 (CAPS) is required for dense-core vesicle but not synaptic vesicle exocytosis in <i>Caenorhabditis elegans</i> . <i>Journal of Neuroscience</i> , <b>2007</b> , 27, 6150-62	6.6	191
18	TrioB Rho-specific GEF domain is the missing Galpha q effector in <i>C. elegans</i> . <i>Genes and Development</i> , <b>2007</b> , 21, 2731-46	12.6	61
17	Functional genomics and biochemical characterization of the <i>C. elegans</i> orthologue of the Machado-Joseph disease protein ataxin-3. <i>FASEB Journal</i> , <b>2007</b> , 21, 1126-36	0.9	53
16	Genetic analysis of dauer formation in <i>Caenorhabditis briggsae</i> . <i>Genetics</i> , <b>2007</b> , 177, 809-18	4	21
15	NCR-1 and NCR-2, the <i>C. elegans</i> homologs of the human Niemann-Pick type C1 disease protein, function upstream of DAF-9 in the dauer formation pathways. <i>Development (Cambridge)</i> , <b>2004</b> , 131, 5741-52	6.6	60
14	Isolation and characterization of high-temperature-induced Dauer formation mutants in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , <b>2003</b> , 165, 127-44	4	57
13	egl-4 acts through a transforming growth factor-beta/SMAD pathway in <i>Caenorhabditis elegans</i> to regulate multiple neuronal circuits in response to sensory cues. <i>Genetics</i> , <b>2000</b> , 156, 123-41	4	86
12	Dauer formation induced by high temperatures in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , <b>2000</b> , 156, 1047-67	4	121
11	Neurosecretory control of aging in <i>Caenorhabditis elegans</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>1999</b> , 96, 7394-7	11.5	97
10	A PDK1 homolog is necessary and sufficient to transduce AGE-1 PI3 kinase signals that regulate diapause in <i>Caenorhabditis elegans</i> . <i>Genes and Development</i> , <b>1999</b> , 13, 1438-52	12.6	307
9	Genetic characterization of the pdu operon: use of 1,2-propanediol in <i>Salmonella typhimurium</i> . <i>Journal of Bacteriology</i> , <b>1997</b> , 179, 1013-22	3.5	48

8	Repression of the cob operon of <i>Salmonella typhimurium</i> by adenosylcobalamin is influenced by mutations in the pdu operon. <i>Journal of Bacteriology</i> , <b>1997</b> , 179, 6084-91	3.5	11
7	Five promoters integrate control of the cob/pdu regulon in <i>Salmonella typhimurium</i> . <i>Journal of Bacteriology</i> , <b>1995</b> , 177, 5401-10	3.5	41
6	The end of the cob operon: evidence that the last gene (cobT) catalyzes synthesis of the lower ligand of vitamin B12, dimethylbenzimidazole. <i>Journal of Bacteriology</i> , <b>1995</b> , 177, 1461-9	3.5	26
5	Two global regulatory systems (Crp and Arc) control the cobalamin/propanediol regulon of <i>Salmonella typhimurium</i> . <i>Journal of Bacteriology</i> , <b>1993</b> , 175, 7200-8	3.5	70
4	A single regulatory gene integrates control of vitamin B12 synthesis and propanediol degradation. <i>Journal of Bacteriology</i> , <b>1992</b> , 174, 2253-66	3.5	109
3	EIPR1 controls dense-core vesicle cargo retention and EARP complex localization in insulin-secreting cells		1
2	Hybridization promotes asexual reproduction in <i>Caenorhabditis</i> nematodes		1
1	CCDC186 controls dense-core vesicle cargo sorting by exit		4