

Adam J Noble

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6257478/publications.pdf>

Version: 2024-02-01

67
papers

1,345
citations

304368

22
h-index

377514

34
g-index

67
all docs

67
docs citations

67
times ranked

1715
citing authors

#	ARTICLE	IF	CITATIONS
1	POSTTRAUMATIC STRESS DISORDER EXPLAINS REDUCED QUALITY OF LIFE IN SUBARACHNOID HEMORRHAGE PATIENTS IN BOTH THE SHORT AND LONG TERM. <i>Neurosurgery</i> , 2008, 63, 1095-1105.	0.6	103
2	The VASOGRADE. <i>Stroke</i> , 2015, 46, 1826-1831.	1.0	97
3	Measuring patient experience: a systematic review to evaluate psychometric properties of patient reported experience measures (PREMs) for emergency care service provision. <i>International Journal for Quality in Health Care</i> , 2017, 29, 314-326.	0.9	79
4	Characteristics associated with quality of life among people with drug-resistant epilepsy. <i>Journal of Neurology</i> , 2017, 264, 1174-1184.	1.8	77
5	Characteristics of people with epilepsy who attend emergency departments: Prospective study of metropolitan hospital attendees. <i>Epilepsia</i> , 2012, 53, 1820-1828.	2.6	58
6	Decision-making by ambulance clinicians in London when managing patients with epilepsy: a qualitative study. <i>Emergency Medicine Journal</i> , 2013, 30, 236-240.	0.4	51
7	Increased Prevalence of Posttraumatic Stress Disorder in Patients After Transient Ischemic Attack. <i>Stroke</i> , 2014, 45, 3360-3366.	1.0	50
8	Which Variables Help Explain the Poor Health-Related Quality of Life After Subarachnoid Hemorrhage? A Meta-analysis. <i>Neurosurgery</i> , 2010, 66, 772-783.	0.6	46
9	Explanations given by people with epilepsy for using emergency medical services: A qualitative study. <i>Epilepsy and Behavior</i> , 2012, 25, 529-533.	0.9	38
10	A nurse-led self-management intervention for people who attend emergency departments with epilepsy: the patients' view. <i>Journal of Neurology</i> , 2013, 260, 1022-1030.	1.8	37
11	Which outcomes should we measure in adult epilepsy trials? The views of people with epilepsy and informal carers. <i>Epilepsy and Behavior</i> , 2016, 59, 105-110.	0.9	34
12	Cognitive-behavioural therapy does not meaningfully reduce depression in most people with epilepsy: a systematic review of clinically reliable improvement. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2018, 89, 1129-1137.	0.9	34
13	Clinical- and Cost-Effectiveness of a Nurse Led Self-Management Intervention to Reduce Emergency Visits by People with Epilepsy. <i>PLoS ONE</i> , 2014, 9, e90789.	1.1	34
14	Anxiety and depression in people with epilepsy: The contribution of metacognitive beliefs. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2017, 50, 153-159.	0.9	32
15	Self-Management education for adults with poorly controlled epILEpsy (SMILE (UK)): a randomised controlled trial protocol. <i>BMC Neurology</i> , 2014, 14, 69.	0.8	30
16	A pilot study of cognitive behavioural therapy and relaxation for migraine headache: a randomised controlled trial. <i>Journal of Neurology</i> , 2015, 262, 2764-2772.	1.8	29
17	The Subarachnoid Hemorrhage International Trialists (SAHIT) Repository: Advancing Clinical Research in Subarachnoid Hemorrhage. <i>Neurocritical Care</i> , 2014, 21, 551-559.	1.2	28
18	Posttraumatic stress disorder in the family and friends of patients who have suffered spontaneous subarachnoid hemorrhage. <i>Journal of Neurosurgery</i> , 2008, 109, 1027-1033.	0.9	27

#	ARTICLE	IF	CITATIONS
19	Subarachnoid Hemorrhage Patients' Fears of Recurrence Are Related to the Presence of Posttraumatic Stress Disorder. <i>Neurosurgery</i> , 2011, 69, 323-333.	0.6	27
20	Outcomes Reported After Surgery for Cauda Equina Syndrome. <i>Spine</i> , 2018, 43, E1005-E1013.	1.0	26
21	What is the cause of post-traumatic stress disorder following subarachnoid haemorrhage? Post-ictal events are key. <i>Acta Neurochirurgica</i> , 2011, 153, 913-922.	0.9	24
22	Qualitative study of paramedics' experiences of managing seizures: a national perspective from England. <i>BMJ Open</i> , 2016, 6, e014022.	0.8	23
23	Self-Management education for adults with poorly controlled epILEpsy [SMILE (UK)]: a randomised controlled trial. <i>Health Technology Assessment</i> , 2018, 22, 1-142.	1.3	23
24	Clinical utility of the Metacognitions Questionnaire 30 in people with epilepsy. <i>Epilepsy and Behavior</i> , 2016, 57, 185-191.	0.9	21
25	Neuropsychologic Impact of Treatment Modalities in Subarachnoid Hemorrhage: Clipping Is No Different from Coiling. <i>World Neurosurgery</i> , 2010, 74, 129-138.	0.7	20
26	The effectiveness of a group self-management education course for adults with poorly controlled epilepsy, SMILE (UK): A randomized controlled trial. <i>Epilepsia</i> , 2018, 59, 1048-1061.	2.6	20
27	Should we stop saying "epileptic"? A comparison of the effect of the terms "epileptic" and "person with epilepsy". <i>Epilepsy and Behavior</i> , 2016, 59, 21-27.	0.9	19
28	"Epileptic", "epileptic person" or "person with epilepsy"? Bringing quantitative and qualitative evidence on the views of UK patients and carers to the terminology debate. <i>Epilepsy and Behavior</i> , 2017, 67, 20-27.	0.9	19
29	Predictors of Delayed Cerebral Ischemia in Patients with Aneurysmal Subarachnoid Hemorrhage with Asymptomatic Angiographic Vasospasm on Admission. <i>World Neurosurgery</i> , 2017, 97, 199-204.	0.7	19
30	Diagnosis of Spontaneous Cervical Artery Dissection May Be Associated with Increased Prevalence of Posttraumatic Stress Disorder. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2014, 23, 335-342.	0.7	17
31	"Seizure First Aid Training" for people with epilepsy who attend emergency departments, and their family and friends: study protocol for intervention development and a pilot randomised controlled trial. <i>BMJ Open</i> , 2015, 5, e009040.	0.8	17
32	Loss of Consciousness at Onset of Aneurysmal Subarachnoid Hemorrhage is Associated with Functional Outcomes in Good-Grade Patients. <i>World Neurosurgery</i> , 2017, 98, 308-313.	0.7	17
33	Self-management education for adults with poorly controlled epilepsy (SMILE (UK)): statistical, economic and qualitative analysis plan for a randomised controlled trial. <i>Trials</i> , 2015, 16, 269.	0.7	16
34	Cauda Equina Syndrome Core Outcome Set (CESCOS): An international patient and healthcare professional consensus for research studies. <i>PLoS ONE</i> , 2020, 15, e0225907.	1.1	16
35	Frequency and factors associated with emergency department attendance for people with epilepsy in a rural UK population. <i>Epilepsy and Behavior</i> , 2017, 68, 192-195.	0.9	12
36	Paramedics' views on their seizure management learning needs: a qualitative study in England. <i>BMJ Open</i> , 2017, 7, e014024.	0.8	11

#	ARTICLE	IF	CITATIONS
37	Psychological distress after subarachnoid hemorrhage: Patient support groups can help us better detect it. <i>Journal of the Neurological Sciences</i> , 2014, 343, 125-131.	0.3	10
38	Implementation fidelity of a self-management course for epilepsy: method and assessment. <i>BMC Medical Research Methodology</i> , 2017, 17, 100.	1.4	10
39	The SAFARI Score to Assess the Risk of Convulsive Seizure During Admission for Aneurysmal Subarachnoid Hemorrhage. <i>Neurosurgery</i> , 2018, 82, 887-893.	0.6	10
40	Metacognitive beliefs and illness perceptions are associated with emotional distress in people with epilepsy. <i>Epilepsy and Behavior</i> , 2018, 86, 9-14.	0.9	9
41	Protocol for the development of a core outcome set for cauda equina syndrome: systematic literature review, qualitative interviews, Delphi survey and consensus meeting. <i>BMJ Open</i> , 2019, 9, e024002.	0.8	8
42	Developing patient-centred, feasible alternative care for adult emergency department users with epilepsy: protocol for the mixed-methods observational 'Collaborate'™ project. <i>BMJ Open</i> , 2019, 9, e031696.	0.8	8
43	Developing and assessing the acceptability of an epilepsy first aid training intervention for patients who visit UK emergency departments: A multi-method study of patients and professionals. <i>Epilepsy and Behavior</i> , 2017, 68, 177-185.	0.9	6
44	Perceptions of emergency care using a seizure care pathway for patients presenting to emergency departments in the North West of England following a seizure: a qualitative study. <i>BMJ Open</i> , 2018, 8, e021246.	0.8	6
45	Family and friends' fears of recurrence: impact on the patient's recovery after subarachnoid hemorrhage. <i>Journal of Neurosurgery</i> , 2013, 119, 948-954.	0.9	5
46	Perceptions and experiences of epilepsy among patients from black ethnic groups in South London. <i>Primary Health Care Research and Development</i> , 2015, 16, 450-460.	0.5	5
47	Are 'Theory of Mind' Skills in People with Epilepsy Related to How Stigmatised They Feel? An Exploratory Study. <i>Behavioural Neurology</i> , 2016, 2016, 1-9.	1.1	5
48	Assessing Treatment Fidelity within an Epilepsy Randomized Controlled Trial: Seizure First Aid Training for People with Epilepsy Who Visit Emergency Departments. <i>Behavioural Neurology</i> , 2019, 2019, 1-11.	1.1	5
49	Patient views on use of emergency and alternative care services for adult epilepsy: A qualitative study. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2020, 80, 56-62.	0.9	5
50	Psychosocial outcome following subarachnoid haemorrhage: An under-researched problem. <i>British Journal of Neuroscience Nursing</i> , 2009, 5, 372-380.	0.1	4
51	Clinically unnecessary and avoidable emergency health service use for epilepsy: A survey of what English services are doing to reduce it. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2020, 76, 156-160.	0.9	4
52	The impact of spontaneous subarachnoid haemorrhage on patients' families and friends. <i>British Journal of Neuroscience Nursing</i> , 2008, 4, 278-285.	0.1	2
53	The effect of noun, adjective, and possessive noun labels on perceptions of someone with epilepsy. <i>Epilepsy and Behavior</i> , 2017, 68, 236-237.	0.9	2
54	Psychological interventions for epilepsy: How good are trialists at assessing their implementation fidelity, what are the barriers, and what are journals doing to encourage it? A mixed methods study. <i>Epilepsy and Behavior</i> , 2019, 97, 174-181.	0.9	2

#	ARTICLE	IF	CITATIONS
55	Seizure first aid training for people with epilepsy attending emergency departments and their significant others: the SAFE intervention and feasibility RCT. Health Services and Delivery Research, 2020, 8, 1-190.	1.4	2
56	Unplanned hospital use by people with epilepsy: A lever by which to bring about increased self-management support?. Epilepsy and Behavior, 2015, 51, 57-59.	0.9	1
57	A disease, disorder, illness or condition: How to label epilepsy?. Acta Neurologica Scandinavica, 2017, 136, 536-540.	1.0	1
58	Seizure First Aid Training For people with Epilepsy (SAFE) frequently attending emergency departments and their significant others: results of a UK multi-centre randomised controlled pilot trial. BMJ Open, 2020, 10, e035516.	0.8	1
59	Better Dead than Alive? Quality of Life After Stroke. , 2014, , 241-255.		1
60	The limited efficacy of psychological interventions for depression in people with Type 1 or Type 2 diabetes: An Individual Participant Data Meta-Analysis (IPD-MA). Journal of Affective Disorders, 2022, 310, 25-31.	2.0	1
61	Worth the paper theyâ€™re printed on? Findings from an independent evaluation of how understandable patient information leaflets for antiseizure medications are. Epilepsia, 2022, , .	2.6	1
62	PO055â€™...Results of an rct of self-management education in epilepsy: smile-uk. Journal of Neurology, Neurosurgery and Psychiatry, 2017, 88, A26.2-A26.	0.9	0
63	Preconception care for women with epilepsy: a mixed methods review. The Cochrane Library, 2020, , .	1.5	0
64	Title is missing!. , 2020, 15, e0225907.		0
65	Title is missing!. , 2020, 15, e0225907.		0
66	Title is missing!. , 2020, 15, e0225907.		0
67	Title is missing!. , 2020, 15, e0225907.		0